

Article

Not peer-reviewed version

The Effectiveness of Applying Digital Teaching Materials in Pediatric Clinical Education

[Hsin-Shu Huang](#)* and [Bih-O Lee](#)

Posted Date: 19 August 2024

doi: 10.20944/preprints202408.1338.v1

Keywords: Artificial Intelligence (AI); Pediatric Clinical Education; National Library e-book storage and reading service



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

The Effectiveness of Applying Digital Teaching Materials in Pediatric Clinical Education

Hsin-Shu Huang ^{1,2,*} and Bih-O Lee ³

¹ Department of Nursing, Central Taiwan University of Science and Technology, Taichung 40601, Taiwan

² Host of the College Smart Integrated Care Practice Program, Workforce Development Agency, Ministry of Labor

³ College of Nursing, Kaohsiung Medical University, Kaohsiung 80708, Taiwan

* Correspondence: hshuang@ctust.edu.tw

Abstract: AI-driven tools can identify nursing students' strengths and weaknesses, thus enabling the development of customized learning plans that target specific areas for improvement. This targeted approach can enhance knowledge retention and skill development, resulting in better prepared healthcare professionals. Nursing students are required to overcome their own psychological stress during their internship in order to understand sick children's emotional reactions, as well as to be able to interact and communicate with such children. The application of AI picture-based teaching e-books helps nursing students to understand how to use therapeutic games to improve communication with sick children. A comparison of the differences between nursing teachers' self-efficacy scores and nursing students' application of AI picture-based teaching e-books (experimental group) and narrative handouts (control group) was achieved via therapeutic games to deal with sick children's fear of medical examinations and treatments. In addition, a comparison was also conducted on the differences between the impact of the self-efficacy of the experimental group and that of the control group in dealing with sick children's behavioral responses with respect to fears of medical examinations and treatments. This study is a quasi-experimental study, in which the self-efficacy scores of 30 nursing students were collected and analyzed in both the experimental group and the control group before and after intervention in dealing with sick children's behavioral responses to fears of medical examinations and treatments. It was found that the self-efficacy score of the experimental group was higher than that of the control group (β coefficient is 0.356) when assessing the self-efficacy scores and behavior of the experimental and control groups in dealing with sick children's fears of medical examinations and treatments through one-way analysis of covariance (one-way ANCOVA); moreover, a statistically significant difference ($p < 0.05$) was observed. The experimental group's score of behavioral responses to the fears of sick children to examinations and treatments was lower than that of the control group (β coefficient was -1.540), wherein a statistically significant difference ($p < 0.05$) was shown. The self-efficacy of the experimental group in dealing with sick children's fears of medical examinations and treatments had a statistically significant impact on the sick children's behavioral responses to the fear of medical examinations and treatments ($p < 0.05$). In addition, the higher the self-efficacy scores of the experimental group in dealing with sick children's fear of medical examinations and treatments, the lower the sick children's behavioral responses to fears of medical examinations and treatments (the β coefficient was -0.914); the control group's self-efficacy scores in dealing with sick children's fears of medical examinations and treatments did not have a statistically significant impact on the sick children's behavioral responses to fears of medical examinations and treatments ($p > 0.05$). This study proves that the application of AI picture-based teaching e-books by nursing teachers is more effective than narrative handouts in improving nursing students' self-efficacy when using therapeutic games to deal with, as well as reduce, sick children's fears of medical examinations and treatments.

Keywords: Artificial Intelligence (AI); pediatric clinical education; national library e-book storage and reading service

1. Introduction

The majority of pediatric care patients are in their early-childhood and preschool years; thus, they are more prone to exhibiting behavioral responses to fears, such as crying and rejection, before medical examinations and treatments as they are too young, have poor oral expression skills, have had bad medical experiences in the past, are being exposed to unfamiliar medical staff, and/or have a lack of unself-conscious personal control, which forms the biggest source of psychological stress for nursing students during pediatric internships. If sick children's fears are not handled in time, they will be prone to low self-esteem, insecurity, lacking self-confidence, and will incur negative impacts on their mental health in the development of personality formation [1]. The psychologists Evie Crotty and Alberto Magni pointed out that children's fearful emotions should be understood to help them face and overcome their fears [2]. Therefore, when sick children receive medical treatment, the timely improvement of behavioral responses to fear is the goal of pediatric care. Therapeutic games are an important and effective intervention measure that helps medical staff to observe sick children's physical and mental needs and feelings, help to mitigate fear and stress, as well as improve sick children's ability to obtain a sense of control over the treatment process [3].

Most nursing students express that, before their clinical internship in pediatrics, they lacked motivation to learn about child development, had never done it before, had a vague sense thereof, and/or could not apply it in clinical communication. As a result, they had a poor response ability to cope with preschool children's fear of medical examinations and treatments, and did not even have confidence in themselves. The higher their learning motivation, the more likely they are to use learning strategies, and the higher their learning efficiency. Thus, the clinical instructor plays a key role in stimulating learning motivation and formulating learning strategies simultaneously. The researcher applied Chat GPT to design AI picture-based teaching e-books of therapeutic games that are suitable for pediatric clinical nursing and can be used in the teaching of pediatric nursing internships based on the literature, pediatric clinical experience, and the learning needs of nursing students. Therapeutic games could be applied in nursing interventions from the beginning of internships for the purpose of helping nursing students, families, and preschool children to communicate smoothly, as well as in developing friendly relationships. It was expected that the nursing students would build confidence in the pediatric internship process and would be willing to continuously use it in future clinical scenarios.

Pediatric patients have a higher level of emotional responses when facing unfamiliar environments and medical staff than adults [4], and this is a common challenge for nursing students on internships in pediatric clinical practice. People of different ages have different desires and needs with respect to using different communication methods, or they may need assistance in the handling of other family-member problems; having said this, however, nursing students on internships in pediatric clinical practice also look forward to participating in clinical nurse jobs. Nursing students hope that they are able to communicate smoothly with children and that they can professionally face challenges from family members, as well as earning recognition and trust from instructors [5].

Going through the internship experience is the best way through which to develop nursing professionalism [6]. Clinical instructors serve as professional models, assisting the students on internships in developing skills, providing professional guidance (where academic theory and clinical practice are combined in the same way, thus becoming the factor having the most impact on students [6,7]), as well as providing a certain professional manner that will enhance the nursing students' learning motivation and attempts to strive and encourage positive behaviors [8]. In addition, the family environment will affect nursing students' concepts, behaviors, attitudes, and even personal future development. Part-time work experience will improve their self-esteem and psychological and physical adjustment, which enhances students' knowledge of future work and

helps them become more independent. Nursing students' clinical work performance, independent character, interpersonal interaction, and self-confidence may be affected by their part-time work experience [9].

Adults can have intrinsic motivation and self-directed learning in any topic. They bring life experience and knowledge into their learning according to different roles and positions, attaching importance to practical operations and goal-oriented and appropriate learning, which is the only way through which the learning effect can be truly achieved. Adult learners like to be respected; as such, the courses should be arranged, if possible, in relation to their life experiences. They can learn new knowledge, attitudes, and skills through dialogue and interaction with teachers [10]. The design of clinical nursing education should be learner-centered, where learner needs and limitations are understood, arranging and planning for pediatric learners is undertaken, and the concept of "therapeutic games" of nursing intervention can be integrated into teaching using vivid and distinctive patterns, health checks, treatments, health education, and other such teaching elements. In addition, nursing students should be taught to use therapeutic games in practical scenarios, in order to enhance their interest and experience in childcare.

Hospitalization always puts great pressure on sick children. For sick children, hospitalization means that they will feel physically uncomfortable and restricted as they need to stay in an unfamiliar environment and undergo unfamiliar medical treatments, such as intravenous injections and the measurement of vital signs, which causes sick children to temporarily lose their sense of self-control [11]; according to the Ministry of Health and Welfare's 2017 announcement, pneumonia ranked 7th among the top 10 causes of death in children and adolescents [12]. According to the 2020 announcement from the Directorate-General of Budget, Accounting and Statistics, Executive Yuan, respiratory diseases and gastrointestinal diseases were ranked as the highest and the second-highest among the top 10 diseases in children in 2018, respectively [13]. Nebulization therapy is a common treatment in pediatric care [14], and abdominal ultrasound is a key diagnostic support for unexplained acute abdomens, which also prevents the occurrence of medical disputes [15]. Less than 5% of pediatric abdominal emergencies are caused by diseases. When ruling out pathological factors, constipation caused by lifestyle is commonly seen. A small-volume enema is commonly used to help relieve constipation, which is a common examination and therapeutic activity in pediatrics [16].

Therapeutic games refer to developing a trust relationship through games, and it helps sick children express their feelings and deal with stressful moods, providing medical staff with the possibility of assessing sick children's cognition and needs in medical scenarios, as well as assisting in adjustment [17]. It assists medical staff in observing sick children's physical and mental needs and feelings, as well as helping sick children to express their fears and stress, improve their sense of security in order to face the treatment process, and to cooperate with examinations and treatments [3]. AI picture-based teaching e-books have been used in various pediatric clinical nursing practices. They can help to alleviate sick children's discomfort during treatments. For preschool children who are in the phase of picture-based cognitive development, they can be a therapeutic story and come with a theme on how to become a good guided medium for education [18]. In the systematic analysis of 16 studies by Brondani, J. P., Pedro, and E. N. R., it was found that the use of therapeutic games—which are explained through story guidance—in nursing intervention enables sick children and their family members to understand the nursing process, as well as helping to cure discomforts in examination and treatment processes. This proves that it is a highly accepted, applicable communication method, which also has low cost, for children in clinic nursing [19].

2. Materials and Methods

2.1. Study Design and Participants

This study is a quasi-experimental study that used Chat GPT to prepare AI picture-based teaching e-books as therapeutic game teaching tools for the experimental group, thereby teaching the nursing students of the experimental group's internship institution to intervene when sick children produce behavioral responses to fear. The control group's internship institution provided handouts to explain the theory and methods when using therapeutic games that might be used to intervene

when sick children produce behavioral responses to fear. The internship institution of the experimental group and the internship institution of the control group conducted evaluations of nursing students' self-efficacy in dealing with sick children's fear of examinations and treatments, as well as sick children's behavioral responses to fear of examinations and treatments before and after intervention.

The research sample estimation method used the G-Power version 3.1 software for calculation and ANOVA with an effect size of 0.25 (medium), alpha level of 0.05, and power of 0.8. The calculation result was based on a sample number of 60 subjects. In the estimated sample collection, there were 30 subjects in each of the experimental and control groups, for a total of 60 subjects. The basic information, such as age, sex, level of education, related working experience, etc., of the two groups did not present significant differences when examined using an independent *t*-test and a chi-square test at a significance level of $p > 0.05$; thus, the two groups had similar conditions prior to intervention.

2.2. Measurement

2.2.1. AI Picture-Based Teaching E-Books

A total of five therapeutic games—namely, “intravenous injection of children,” “vital signs measurement of children,” “bedside ultrasound of children,” “small volume enema of children,” and “aerosol therapy of children”—were used. These were integrated with the learning needs of nursing students, as well as in the examinations and treatments commonly seen in pediatric units and in literature reviews. The accuracy and suitability of the content and information conveyed in the AI picture-based teaching e-books were tested in terms of their validity by experts; specifically, six nursing teachers with more than three years of clinical experience (CVI value of 1.0).

2.2.2. Nursing Students' Self-Efficacy Scale for Preschool Sick Children's Fears of Medical Examinations and Treatments

This part was conducted using a revised version of the “Nursing Students' Self-Efficacy Scale for Preschool Sick Children's Fear of Medical Examination and Treatment,” which refers to the Chinese version of the General Self-Efficacy Scale [20]. The scale was rated by six nursing teachers with more than three years of clinical experience for content suitability and text clarity, where $CVI \geq 0.8$ was retained, 10 questions had a CVI value > 0.83 –1, and the overall CVI value was 0.92. Using Cronbach's α as a measure of the internal consistency reliability of the scale and a total of 10 questions, a pre-test was conducted on 30 nursing students on the internship program, and the results showed that the Cronbach's α of the total scale was 0.809. The test was scored on a 4-point Likert scale, with 1 to 4, respectively, representing “not at all correct”, “somewhat correct”, “mostly correct”, and “completely correct”. The total score scale ranged from 0 to 40, and higher scores indicate greater self-efficacy.

2.2.3. Observation Scale for Sick Children's Behavioral Response to Fear of Medical Examinations and Treatments

The content validity was assessed by experts based on a literature review and clinical experiences. Six nursing teachers with more than three years of clinical experience rated the content in terms of suitability and text clarity. A $CVI \geq 0.8$ was retained, and 8 of the questions had a CVI value ≥ 0.8 –1. The overall CVI value was 0.9. Six sick children undertook a test using the Observation Scale for Sick Children's Behavioral Response to Fear of Medical Examination and Treatment, and the consistency among the observers' observation on the behavioral responses to fear reached 100%. The internal consistency reliability of the scale was tested with Cronbach's α , and the 30 sick children's behavior was observed. Cronbach's α was 0.76, with a total of 8 scoring items. Each of the remaining items were worth 1 point, with a maximum score of 8 points. Higher scores indicate greater fear.

2.3. Research Time and Place

The study period was from June 1, 2022, to July 31, 2023. The subjects of this study were pediatric nursing students on an internship in a teaching hospital.

2.4. Ethical Considerations

This research project was reviewed and approved by the IRB of the China Medical University Hospital (no: CRREC-111-051). The research subjects could quit at any time, had the right to raise questions, and the questionnaires were anonymous. The questionnaire results were numbered anonymously in order to delink and ensure confidentiality. The names and conditions of the research subjects will never be publicized, and the results are for academic use only.

2.5. Data Collection and Analysis

After the data were collected, decoded, logged, and archived, statistical analysis was performed with the SPSS Version 26.0 statistical software package for Windows/PC. Statistical analysis was conducted based on the research purpose, and $\alpha = 0.05$ was set as the standard for significant differences.

2.5.1. One-Way Analysis of Covariance (ANCOVA)

It was required that the homogeneity of the within-group regression coefficient be tested in order to ensure that there was no interactions between the groups in the pretest before carrying out the one-way analysis of covariance (ANCOVA). This was required as differences may have existed between the control groups at the pretest stage.

2.5.2. Generalized Estimating Equation (GEE)

When the data pertained to longitudinal data, the status of the same subject at different time points was recorded, and there was also a correlation between the observation values from the same subject. As the same subjects in this study were observed at multiple time points (before and after intervention), they can be considered interdependent; therefore, this study is suitable for GEE analysis. As the dependent variable in this study is a numerical variable, it is statistically significant ($p < 0.05$) when the assumed distribution in the GEE is a normal distribution, the link function has identity, and the 95% CI of its β does not contain 0.

3. Results

3.1. Comparison between Self-Efficacy of the Experimental Group and the Self-Efficacy of the Control Group after Intervention (ANCOVA)

As can be seen from Table 1, the test results of the homogeneity of the within-group regression coefficient for self-efficacy in the group $\times$ pretest did not reach the level of a statistically significant difference ($F = 1.743, p > 0.05$), which indicates that the assumption of the homogeneity of regression coefficients was not violated. Thus, a one-way analysis of covariance (ANCOVA) could be carried out directly.

Table 1. Test of the homogeneity of the within-group regression coefficient for self-efficacy.

Source of Variation	Sum of Squares (SS)	Degrees of Freedom (df)	Mean Square (MS)	F	P value
Group	0.691	1	0.691	4.514	0.037
Pretest	9.992	1	9.992	65.303	<0.001
Group $\times$ Pretest	0.267	1	0.267	1.743	0.191
Error	11.017	72	0.153		

As can be seen from Table 2, under the self-efficacy score before the control of interventions, the self-efficacy scores of the groups (experimental group vs. control group) reached a statistically significant difference ($p < 0.05$), which means that the self-efficacy score of the experimental group was higher than that of the control group (where the β coefficient was 0.356).

Table 2. Comparison of the self-efficacy of the experimental group and the self-efficacy of the control group after intervention.

Variable Name	β	SE	P value	95% C.I.	
				Lower	Upper
Self-Efficacy before Intervention	0.805	0.097	<0.001	0.612	0.999
Group (Experimental Group vs. Control Group)	0.356	0.090	<0.001	0.176	0.537
Dependent Variable: Self-Efficacy after Intervention					

3.2. Comparison between Experimental Group and Control Group on Behavioral Response to Fear after Intervention (ANCOVA)

As can be seen from Table 3, the test results of the homogeneity of the within-group regression coefficient for fear in the group $\times$ pretest did not reach the level of a statistically significant difference ($F = 1.264, p > 0.05$), which indicates that the assumption of the homogeneity of regression coefficients was not violated. Thus, one-way analysis of covariance (ANCOVA) could be carried out directly.

Table 3. Test of the homogeneity of the within-group regression coefficient for fear.

Source of Variation	Sum of Squares (SS)	Degrees of Freedom (df)	Mean Square (MS)	F	P value
Group	0.102	1	0.102	0.040	0.842
Pretest	29.356	1	29.356	11.486	0.001
Group $\times$ Pretest	3.231	1	3.231	1.264	0.265
Error	184.021	72	2.556		

As can be seen from Table 4, regarding the fear before the control of the interventions, the fear scores of the groups (experimental group vs. control group) reached the level of a statistically significant difference ($p < 0.05$), which means that the fear score of the experimental group was lower than that of the control group (where the β coefficient was -1.540).

Table 4. Comparison of the fear of the experimental group and the fear of the control group after intervention.

Variable Name	β	SE	P value	95% C.I.	
				Lower	Upper
Fear before Intervention	0.404	0.125	0.002	0.154	0.654
Group (Experimental Group vs. Control Group)	-1.540	0.374	<0.001	-2.285	-0.794
Dependent Variable: Fear after Intervention					

3.3. Comparison of the Difference between the Experimental Group’s Impact of Self-Efficacy on the Behavioral Responses to Fear and the Control Group’s Impact of Self-Efficacy on The behavioral Responses to Fear (GEE)

Next, generalized estimating equations (GEE) was used to conduct an analysis. First, the aim was to understand whether the self-efficacy of the experimental group and the control group had a significant impact on fear; second, the aim was to then further compare whether there was a significant difference in the impact of self-efficacy on the fear between the two groups.

As can be seen from Table 5, the experimental group’s self-efficacy had a statistically significant impact on fear ($p < 0.05$), and the higher the experimental group’s self-efficacy, the lower the fear (where the β coefficient was -0.914); in addition, the control group’s self-efficacy had no statistically significant impact on fear ($p > 0.05$).

Table 5. The experimental group’s and the control group’s impact of self-efficacy on fear.

Variable Name	β	SE	P value	95% C.I.	
				Lower	Upper
Time Points					
After Intervention	3.291	0.883	<0.001	1.560	5.021
Before Intervention	5.356	0.782	<0.001	3.824	6.888
Group (Experimental Group vs. Control Group)	0.667	1.104	0.546	-1.497	2.831
Group × Self-Efficacy					
Experimental Group × Self-Efficacy	-0.914	0.351	0.009	-1.602	-0.226
Control Group × Self-Efficacy	-0.224	0.342	0.512	-0.896	0.447
Dependent Variable: Fear					

4. Discussion

The findings of this study were deemed to be consistent with the studies of Huang et al.; Chang et al.; Natalya A. et al.; Holliman, R. P., and Blanco; P. J., Bliss; and H. and Gildner, J., who showed that therapeutic games can significantly improve sick children’s behavioral responses to fear [21–25]. AI picture-based teaching e-books should be used to teach nursing students how to use therapeutic games. One-way analysis of covariance (ANCOVA) was used to find that the self-efficacy scores of the experimental group in dealing with the sick children’s fear of examinations and treatments were higher when compared with that of the control group (where the β coefficient was 0.356), and the difference was also found to be statistically significant ($p < 0.05$). The score of the sick children’s behavioral responses to fear in the experimental group was lower than that in the control group (where the β coefficient was -1.540), which reached the level of a statistically significant difference ($p < 0.05$). The self-efficacy of the experimental group in dealing with the sick children’s fear of examinations and treatments had a statistically significant impact on the sick children’s behavioral responses to fear of examinations and treatments ($p < 0.05$), and the higher self-efficacy of the experimental group in dealing with sick children’s fear of examinations and treatments led to lower sick children’s behavioral responses to fear of examinations and treatments (where the β coefficient was -0.914). The self-efficacy of the control group in dealing with the sick children’s fear of examinations and treatments did not have a statistically significant impact on the sick children’s behavioral response to fear of examinations and treatments ($p > 0.05$); therefore, this study also demonstrates the benefits of applying AI-based materials in clinical education [26].

5. Conclusions

This study proved that AI picture-based teaching e-books have a better effect than narrative handouts in improving nursing students’ self-efficacy in using therapeutic games to deal with sick children’s fear of examinations and treatments and in reducing sick children’s behavioral responses to fear of examinations and treatments. There is a need for application research at different levels in the future to explore the application effectiveness of AI-based materials in medical and nursing

education. A limitation of this study is that the research subjects were all female nursing students; as such, the learning effectiveness of male nursing students cannot be inferred.

Author Contributions: Conceptualization, H.-S.H.; data curation, H.-S.H.; methodology, H.-S.H.; resources, B.-O.L.; supervision, H.-S.H.; writing—original draft, H.-S.H.; writing—review and editing, H.-S.H. The author has read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: This study conducted data collection after receiving approval from the Central Regional Research Ethics Committee of China Medical University Hospital (IRB No: CRREC-111-051).

Informed Consent Statement: Informed consent was obtained from all of the subjects involved in this study.

Data Availability Statement: All of the relevant datasets in this study are described in the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Huoy-Jia Tzeng. A Study of Fears. *Journal of National Taipei Teachers College*. **2000**, 13, 265-292. <https://academic.ntue.edu.tw/ezfiles/7/1007/img/41/13-10.pdf>
2. Evie Crotty, Alberto Magni. *The Psychology of Children's Fear*. Publisher: Posts & Telecommunications Press, **2021**. ISBN: 9787115558947
3. Jui-Hsuan Li, Yi-Chen Chang, Fang-Chi Hu, Whei-Mei Shih, Chen-Li Chen, and Mei-Chuan Shih. Reducing Pain Indexes for Preschool Children when Receiving Emergency Room Intravenous Injections. *The Journal of Health Sciences*. **2018**, 4(2), 20-34. [http://doi.org/10.6979/TJHS.201803_4\(2\).0002](http://doi.org/10.6979/TJHS.201803_4(2).0002)
4. Jeng-Chung Woo. A Study on the Application of Emotional Design on the Anxiety and Pain Alleviation in Pediatric Outpatients. *Journal of Design*. **2012**, 17(3), 69-93. <http://doi.org/10.6381/JD.201209.0069>
5. Pi-Chun Wu, Jason C. Chan. Reflecting on the Perspective Transformation of Competency-Based Education. *Journal of Educational Research and Development*. **2018**, 14(2), 35-64. <http://doi.org/10.3966/181665042018061402002>
6. Ming-Sea Tseng, Fen Liu, Shu-Chun Lin. Cultivating Professional Nursing Behaviors—Experiences of Nursing Students for Fundamental Nursing Practicum. *Chang Gung Nursing*. **2018**, 29(2), 173-189. [https://doi.org/10.6386/CGN.201806_29\(2\).0002](https://doi.org/10.6386/CGN.201806_29(2).0002)
7. Shiah-Lian Chen, Weillie Lee. The Impact of Learning Styles of Nursing Students and Clinical Instructors on Academic Achievement and Teaching Satisfaction. *Nursing Research*. **2000**, 8(3), 313-324. <http://doi.org/10.7081/NR.200006.0313>
8. Yi-Jung Lin, Yu-Ching Tu, Hsu-Ling Hsieh. A Study of the Nursing Competency of Nursing College Students: From the Perspective of Nurses and Clinical Nursing Teachers. *Journal of Nursing and Healthcare Research*. **2009**, 5(3), 220-230. <http://doi.org/10.6225/JNHR.5.3.220>
9. Chih-Feng Tseng. The Study of the Influences of Parents' Social-Economic Background on Parenting: The Analysis of Mediated Effect of Personal Value and Perception of Educational Purpose. *National Chiayi University Journal of the Educational Research*. **2013**, 31, 85-118. <http://doi.org/10.6474/NCUJER>
10. Pi-Chi Wu, Fan-Hao Chou, and Bih-O Lee. Challenges and Reflections of Clinical Nursing Education. *Chang Gung Nursing*. **2016**, 27(1), 26-33. <http://doi.org/10.3966/102673012016032701003>
11. Yu-Chin Wang, Pei-Hua Chiu, Jia-Xin Li. Reducing the Fear of Intravenous Injection in Preschool Children by Using Openable Fixed Strap. *Tzu Chi Nursing Journal*. **2021**, 20(3), 106-116. <https://www.airitilibrary.com/Publication/alDetailedMesh?DocID=16831624-202106-202106240016-202106240016-106-116>
12. Ministry of Health and Welfare. *Taiwan's Leading Causes of Death in 2017*. Ministry of Health and Welfare: Taipei, Taiwan, **2017**. <https://www.mohw.gov.tw/cp-3795-41794-1.html>2018/6/15
13. Directorate-General of Budget, Accounting and Statistics, Executive Yuan. *10 Common Childhood Illnesses and Their Treatments*. Directorate-General of Budget, Accounting and Statistics, Executive Yuan: Taipei, Taiwan, **2020**. <http://www1.stat.gov.tw/public/Data/069135930A8RC8SAF.pdf>
14. Hsiu-Fang Pan, Hsiang-Ying Hung, Tzu-Wei Tseng, Mei-Chuan Wu. A Project to Improve the Rate of Accuracy of Aerosol Therapy in Pediatric Wards. *Show Chwan Medical Journal*. **2020**, 19(2), 149-160. <http://doi.org/10.3966/156104972020121902003>
15. Hsin-Hung Lin. Common Abdominal Emergencies in Children. *Journal of Traditional Chinese Pediatric Medicine*. **2012**, 14(1), 16-22. <http://doi.org/10.30033/JTCDM.201206.0003>
16. Juiju Tseng. Children's constipation is no longer bothersome, find the right method to improve it. *CHANGHUA CHRISTIAN HOSPITAL*. **2016**, 33(11), 12-13. <http://doi.org/10.29598/XLZY>

17. Li-Min Wu, Yi-Fang Chen, Yu-Chien Chen, Bao-Ru Chen. Clinical Practice of the Therapeutic Play. *Journal of MacKay Nursing*. **2017**, 11(1), 7-17. [http://doi.org/10.29415/JMKN.201701_11\(1\).0001](http://doi.org/10.29415/JMKN.201701_11(1).0001)
18. Tsao, Y., Kuo, H. C., Lee, H. C., & Yiin, S. J. Developing a medical picture book for reducing venipuncture distress in preschool-aged children. *International journal of nursing practice*. **2017**, 23(5), e12569. <http://doi.org/10.1111/ijn.12569>
19. Brondani, J. P., & Pedro, E. N. R. The use of children's stories in nursing care for the child: an integrative review. *Revista brasileira de enfermagem*. **2019**, 72, 333-342. <https://doi.org/10.1590/0034-7167-2018-0456>
20. Schwarzer, R., Bäßler, J., Kwiatek, P., Schröder, K., & Zhang, J. X. The assessment of optimistic self-beliefs: Comparison of the german, spanish, and chinese versions of the general self-efficacy scale. *Applied Psychology*. **1997**, 46(1), 69-88. <http://doi.org/10.1111/j.1464-0597.1997.tb01096.x>
21. Fang Mei Huang, Su Fen Cheng, Ya Ling Lee, Chi Wen Chen. Effectiveness of Distraction Strategies in Preschool Children Undergoing Intravenous Line Placement at the Emergency Department in Regional Hospitals. *VGH Nursing*. **2017**, 34(4), 342-355. [http://doi.org/10.6142/VGHN.201712_34\(4\).0002](http://doi.org/10.6142/VGHN.201712_34(4).0002)
22. Chi-Wen Chang, Yueh-Tao Chiang, Hsing-Yi Yu. Clinical Application of Therapeutic Play in Children. *Yuan-Yuan Nursing*. **2017**, 11(1), 44-51. https://www.airitilibrary.com/Common/Click_DOI?DOI=10.6530%2fYYN%2f2017.5.4
23. Natalya A., L., Chiao-Feng, C., Sarah, C., Jeffrey M., S., Sinem, A., Meany-Walen, & Kristin, K. The impact of child-centered play therapy training on attitude, knowledge, and skills. *International Journal of Play Therapy*. **2012**, 21(3), 149-166. DOI:10.1037/a0028039. <https://doi.org/10.1037/a0028039>
24. Holliman, R. P., & Blanco, P. J. An Exploratory Mixed Methods Study of the Development of Self-Efficacy, Attitudes, Knowledge, and Skills for Master's Level School-Based Internship Students Attending Play Therapy Supervision. *The Practitioner Scholar: Journal of the International Trauma Training Institute*. **2019**, 1, 23-39. <https://thepractitionerscholar.com/article/view/18511/13137>
25. Bliss, H., & Gildner, J. "Using Echo360 to Support Active Learning". *South Carolina Conference on Innovations in Teaching and Learning*. **2021**. <https://digitalcommons.coastal.edu/sccitl/2021/fullconference/67/>
26. Nagi, F., Salih, R., Aizubaidi, M., Shah, H., Alam, T., Shah, Z., & Househ, M. Applications of Artificial Intelligence (AI) in Medical Education: A Scoping Review. This article is published online with Open Access by IOS Press and distributed under the terms of the Creative Commons Attribution Non-Commercial License 4.0 (CC BY-NC 4.0). **2023**. doi:10.3233/SHTI230581. <https://doi.org/10.5281/zenodo.7866503>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.