

Review

Not peer-reviewed version

Unveiling the Potential of Phytocannabinoids: Exploring Marijuana's Lesser-Known Constituents for Neurological Disorders

[Balapal S Basavarajappa](#)* and [Shivakumar Subbanna](#)

Posted Date: 27 September 2024

doi: 10.20944/preprints202409.2224.v1

Keywords: Endocannabinoids; Neurodegenerative disorders; Alcohol use disorders; substance use disorders; Psychiatric disorders; Epilepsy; Cannabinoid receptors; Alzheimer's disease; Parkinson's disease; Huntington's disease



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

Unveiling the Potential of Phytocannabinoids: Exploring Marijuana's Lesser-Known Constituents for Neurological Disorders

Balopal S. Basavarajappa ^{1,2,*} and Shivakumar Subbanna ¹

¹ Center for Dementia Research, Nathan Kline Institute for Psychiatric Research, Orangeburg, NY 10962, USA

² Department of Psychiatry, New York University Grossman School of Medicine, New York, NY 10016, USA

* Basavaraj.Balopal@nki.rfmh.org; Tel.: +1-845-398-3234

Abstract: Cannabis sativa is known for producing over 120 distinct phytocannabinoids, with Δ^9 -tetrahydrocannabinol (Δ^9 -THC) and cannabidiol (CBD) being the most prominent, primarily in their acidic forms. Beyond Δ^9 -THC and CBD, a wide array of lesser-known phytocannabinoids, along with terpenes, flavonoids, and alkaloids, demonstrate diverse pharmacological activities, interacting with the endocannabinoid system (eCB) and other biological pathways. These compounds, characterized by phenolic structures and hydroxyl groups, possess lipophilic properties, allowing them to cross the blood-brain barrier (BBB) effectively. Notably, their antioxidant, anti-inflammatory, and neuro-modulatory effects position them as promising agents in treating neurodegenerative disorders. While research has extensively examined the neuropsychiatric and neuroprotective effects of Δ^9 -THC, other minor phytocannabinoids remain underexplored. Given the well-established neuroprotective potential of CBD, there is growing interest in the therapeutic benefits of non-psychotropic minor phytocannabinoids (NMPs) in brain disorders. This review highlights the emerging research on these lesser-known compounds and their neuroprotective potential. It offers insights into their therapeutic applications across various major neurological conditions.

Keywords: endocannabinoids; neurodegenerative disorders; alcohol use disorders; substance use disorders; psychiatric disorders; epilepsy; cannabinoid receptors; Alzheimer's disease; Parkinson's disease; Huntington's disease

1. Introduction

Neurodegenerative disorders (NDs) pose a significant global challenge, contributing extensively to disability and mortality [1–3]. The World Health Organization (WHO) projects that by 2040, NDs will be the second leading cause of death worldwide [4]. These disorders involve the steady loss of neurons, resulting in impaired brain function, the breakdown of neuronal networks, and ultimately leading to disability. Despite substantial research efforts, current treatments primarily focus on managing symptoms, as neuronal degeneration often progresses silently for years before clinical signs such as cognitive, motor, or other neurological symptoms manifest. Cell death, synaptic dysfunction, and behavioral abnormalities underlie most neurodegenerative diseases, such as Alzheimer's disease (AD), Parkinson's disease (PD), and Huntington's disease (HD). In AD, amyloid plaques and tau protein aggregates drive neurodegeneration, while synucleinopathies in PD are marked by Lewy body accumulations [5]. HD, in contrast, is caused by a genetic mutation in the huntingtin gene (Htt) involving expanded CAG repeats [6–8]. Other conditions like epilepsy and stroke, though distinct in pathology [9], share overlapping features with NDs, including neuronal damage and behavioral symptoms. Environmental factors, such as substance abuse, can exacerbate these conditions, which often lead to chronic cognitive, emotional, and behavioral impairments. Current therapies for NDs primarily focus on symptom management and slowing disease progression. Medications like cholinesterase inhibitors and NMDA receptor antagonists offer limited cognitive improvement in AD [10,11]. At the same time, levodopa and dopamine agonists alleviate

PD motor symptoms but fail to halt disease progression [12]. In HD, tetrabenazine reduces chorea but introduces neuropsychiatric side effects [13–16]. Although advances in gene therapy, stem cell research, and precision medicine offer hope for future treatments, the urgent need remains for novel therapies that manage symptoms and address underlying disease mechanisms with fewer side effects.

The discovery of cannabinoid receptors sparked interest in the endocannabinoid system (eCB), with endogenous ligands like anandamide (AEA) and 2-arachidonoyl-glycerol (2-AG) [17,18] playing critical roles in CNS functions, including pain regulation, neurogenesis, and immune modulation. Though CB1 receptors are abundant in the brain and mediate the psychoactive effects of THC (for references, see [19]), CB2 receptors, primarily found in immune cells, exhibit minimal psychoactivity [20–22]. Both receptor types are now recognized for their broader involvement in neuroprotective and anti-inflammatory processes. In recent years, we and others have reviewed a detailed discussion on the function of the eCB system in health and disease [23–27]. Over 120 phytocannabinoids have been identified in *Cannabis sativa* [28], with Δ^9 -tetrahydrocannabinol (Δ^9 -THC) and cannabidiol (CBD) being the most well-known. While Δ^9 -THC is primarily associated with psychotropic effects, both Δ^9 -THC and Δ^8 -THC [29] have demonstrated therapeutic benefits, including appetite stimulation, analgesia, and antiemetic properties [30–33]. Non-psychotropic cannabinoids like CBD have gained attention for their anti-inflammatory, anxiolytic, and neuroprotective effects [34]. These compounds are now understood to target multiple receptors and biological pathways beyond the classical CB1 and CB2 receptors. Their phenolic structures endow them with potent antioxidant properties, while their lipophilicity facilitates blood-brain barrier (BBB) penetration [35–37], making them attractive candidates for treating neurodegenerative diseases.

This review aims to explore the neuroprotective potential of non-psychotropic minor phytocannabinoids (NMPs)—such as CBD, Δ^9 -tetrahydrocannabivarin (THCV), cannabidivarin (CBDV), and cannabigerol (CBG)—in neurodegenerative conditions. These phytocannabinoids are selected based on their abundance, ease of synthesis, and demonstrated therapeutic potential in other areas, such as cancer and inflammatory disorders. Their structural similarity to CBD, a well-studied neuroprotective agent, suggests they may also hold significant promise for future clinical applications.

Role of NMPs in Epilepsy

CBD

CBD function as a negative allosteric modulator on CB1 and CB2 receptors (**Figure 1**). It also exhibits affinity for various other receptors, such as the transient receptor potential vanilloid type 1 channel (TRPV1), peroxisome proliferator-activated receptor γ (PPAR γ), G protein-coupled receptor 55 (GPR55), 5-hydroxytryptamine family 1A (5-HT1A) receptor, γ -aminobutyric acid type A (GABAA) receptor, and transient receptor potential cation channel subfamily M member 8 (TRPM8) (for references, see [19]). However, challenges remain in establishing the neuroprotective function of CBD through any one of these targets. Studies have indicated that eCBs regulate seizure mechanisms in developing animals through CB1 receptor signaling [38]. In pre-clinical seizure models, an anticonvulsant effect was observed due to a pharmacological increase in AEA and 2-AG following neuronal hyperexcitability, which reduces glutamate excitotoxicity during seizures [38–40]. Many axon terminals in the CNS express CB1 receptors, which inhibit the release of both excitatory and inhibitory neurotransmitters [41,42].




Figure 1. Receptors mediating CBD's agonistic and antagonistic actions. CBD interacts with various receptors, exhibiting a diverse pharmacological profile. Unlike THC, CBD does not exhibit a high affinity at the CB1 and CB2 receptors but can function as a negative allosteric modulator of CB1 [43], indirectly influencing endocannabinoid signaling. Additionally, CBD acts as an agonist at the 5-HT_{1A} serotonin receptor [44], contributing to its anxiolytic and antidepressant effects. It also modulates TRPV1/2 [45], which are involved in inflammation and pain perception. CBD is an antagonist of GPR55, a receptor implicated in anxiety-related behaviors [46], learning and memory [47] and pain perception [48]. CBD's receptor specificity contributes to its broad therapeutic potential across various neurological disorders. CBD, Cannabidiol; THC, Δ^9 -tetrahydrocannabinol; TRPV, Transient receptor potential vanilloid. →, agonist, antagonist.

Additionally, the anticonvulsant action of CBD involves several mechanisms beyond neuronal death prevention, including reinstating functions of hippocampal interneuron in a temporal lobe epilepsy model [49–51]. Furthermore, CBD enhances key molecular components of downstream ERK1/2 [52], PI3K/AKT [53], and pCREB pathways [54] typically linked to endocannabinoid-driven facilitation of adult neurogenesis. Subchronic CBD treatments have been shown to upregulate critical effectors involved in protein synthesis and neuronal survival in the hippocampus, including brain-derived neurotrophic factor (BDNF), MAP-2, calbindin, synapsin 1, and the activation of PPAR γ . Further, CBD activates different survival and synaptic remodeling signaling cascades such as ERK1/2-CREB [55], GSK3 β and PSD95 [56] or PI3K/mTOR/p70S6K [52,57–59] (for general signaling mechanism see Figure 2).




Figure 2. Schematic representation of the signaling pathways modulated by CBD through various receptors in the brain. CBD acts as an agonist, activating several receptors such as 5-HT1A, NMDA/AMPA, Wnt-Frizzled (Fz) family receptors, and TRPV1/2—additionally, CBD functions as a negative allosteric modulator of CB1 and an inhibitor of FAAH, an enzyme involved in the degradation of anandamide (AEA) and an antagonist of GPR55. While the precise mechanisms of action in specific neurodegenerative diseases remain unclear, the collective synergistic effects of CBD have been shown to contribute to its neuroprotective role in several neurological disorder models. CBD's influence on the ERK1/2[52], PI3K/AKT [53], and pCREB [60] pathways, which are critical for regulating genes required for cell death, survival, neurogenesis, and synaptic plasticity including BDNF, MAP-2, synapsin 1, calbindin and PPAR γ , suggests its potential to elicit neuroprotective effects.

CBD decreases epileptiform local field potential burst amplitude in the CA1 and dentate gyrus regions in a CB1 receptor-independent manner [61]. This observation is significant because, in addition to CB1 and CB2 receptors, CBD also binds to 5-HT1A receptors in the human brain [62]. These Gi/o protein-coupled serotonergic receptors, when activated, induce inhibitory effects. In temporal lobe epilepsy, 5-HT1A receptor signaling appears crucial in activating seizure-induced cell proliferation and survival in the dentate gyrus [63]. Experimental and clinical findings suggest that serotonergic signaling impacts seizure susceptibility and epilepsy-associated psychiatric comorbidities [64]. A reduction in serotonergic receptor activity in the dentate gyrus may lead to an excitatory/inhibitory imbalance in the trisynaptic circuit, triggering the neuropathology of epilepsy. Another study observed a significant loss of 5-HT1A binding in the hippocampus, temporal cortex, amygdaloid complex, and frontal lobe in patients with temporal lobe epilepsy [65]. Additionally, the downregulation of postsynaptic 5-HT1A receptors exacerbated the severity of epilepsy-associated depression [66]. Decreased serotonin signaling in brain regions related to epilepsy and emotions,

such as the hippocampus, amygdaloid complex, and medial prefrontal lobe structures, can promote epilepsy and mental disorders, making CBD a potential therapeutic agent. CBD has also attenuated seizures in genetic models of brainstem and limbic seizures [67]. Thus, several well-designed trials have demonstrated the effectiveness of CBD in controlling epileptic seizures, with minimal significant adverse effects, although agitation, irritability, mood changes, and aggressive behavior have been noted [68]. Based on preclinical and clinical trials, the use of CBD (branded as Epidiolex) for Dravet syndrome, Lennox–Gastaut syndrome, or tuberous sclerosis complex, was authorized by the FDA in 2018 and by the European Medicines Agency in 2019.

THCV

THCV (Figure 3) is a relatively abundant, non-psychoactive phytocannabinoid found in the *Cannabis sativa* [69] plant. It primarily functions as a CB1 receptor antagonist, displacing radiolabeled CB1 agonists in mouse whole-brain membranes [70]. THCV also antagonizes CB1-specific GTP γ S binding in various mouse brain regions, including the whole brain, cerebellum, and piriform cortex [70,77]. Moreover, it inhibits Δ^9 -THC-induced anti-nociception and hypothermia in mice [78]. There is evidence from a study in rats that THCV is anti-psychotic in a phencyclidine model of psychosis to a similar degree as clozapine [73]. In tissues enriched with CB2 receptors, THCV acts as a partial antagonist in Gi protein-coupled activities, as evidenced by its effect on forskolin-induced cytosolic cAMP levels [79]. Additionally, THCV regulates calcium transport in epithelial cells by inhibiting TRPV channels [80].




Figure 3. Receptors mediating THCV's agonistic and antagonistic actions. THCV is a minor cannabinoid with a distinctive pharmacological profile, exhibiting both agonistic and antagonistic effects on cannabinoid and non-cannabinoid receptors. THCV has a unique interaction with CB1 receptors [21,71]. At low doses, it acts as a CB1 antagonist, which can counteract the psychoactive effects of THC. However, at higher doses, THCV can act as a CB1 agonist, producing effects similar to THC, such as appetite suppression and potential psychoactivity. THCV is generally regarded as a partial agonist of CB2, which may contribute to its anti-inflammatory and immunomodulatory properties [21,71]. THCV also modulates transient receptor potential (TRP) channels, particularly TRPV1 [72]. These channels play roles in pain perception, thermoregulation, and inflammation, suggesting THCV's potential in managing pain and inflammatory conditions. Similar to CBD, THCV has been shown to interact with the 5-HT_{1A} receptor [73], which is involved in spatial learning and memory, anxiety regulation and mood modulation [74,75]. This receptor interaction may underlie

some of THCV's potential anxiolytic and antipsychotic effects. THCV activates GPR55 [76] anxiety-related behaviors [46], learning and memory [47]. THCV's receptor specificity and its ability to shift between agonist and antagonist roles give it diverse therapeutic potential, particularly in areas like metabolic disorders and neuroprotection. THCV, Δ^9 -Tetrahydrocannabivarin; CBD, Cannabidiol; THC, Δ^9 -tetrahydrocannabinol; TRPV, Transient receptor potential vanilloid. \rightarrow , agonist; \dashv , antagonist.

THCV has also demonstrated protective effects in healthy volunteers who exhibited enhanced neural responses to various stimuli [64]. It was shown to be safe and holds likely for treating central nervous system (CNS) disorders. This makes THCV one of the most promising phytocannabinoids poised for drug approval by regulatory bodies. Indeed, THCV acts as a CB1 receptor antagonist without exerting agonist effects on [35 S]GTP γ S binding, significantly reducing seizure incidence in adult rat piriform cortex slices [81]. In another study, THCV reduced Purkinje cell firing by enhancing inhibitory neurotransmission at interneuron-Purkinje cell synapses in mouse cerebellar brain slices. This was achieved by decreasing CB1 receptor-mediated, endocannabinoid-induced inhibition of GABA release [82]. The proposed mechanism for THCV's anticonvulsant action is the increased GABA release through the blockade of CB1 receptors for endocannabinoids [21]. THCV has also shown efficacy in animal models of epilepsy and in NMR neuroimaging studies involving patients with epilepsy [83,84]. However, further studies with varying doses are needed to evaluate its efficacy in epilepsy treatment.

CBDV

CBDV (Figure 4) is found in unhybridized pure sativa and indica cannabis plants, known as landrace strains. These strains contain high amounts of CBD and low amounts of Δ^9 -THC. CBDV exhibits low binding affinity for the CB1 and CB2 receptors [85,86] and, therefore, appears to lack the psychotropic effects seen with THC. Consistent with this function, high concentrations of CBDV are required to stimulate CB1 receptor-coupled activation of (35 S) GTP γ S binding, inhibit cAMP synthesis, and recruit β -arrestin 2 (β arr2) [85,87,88]. Primarily, CBDV is a more potent and efficacious agonist at CB2 receptors [87,88]. It also activates TRPV1-4 channels [89,90] and stimulates ERK1/2 phosphorylation, inhibiting LPS-mediated signaling [76] through the GPR55 receptor. Additionally, CBDV acts as an inverse agonist at the Gs-coupled GPR6 receptor, leading to the stimulation of adenylyl cyclase and the recruitment of β -arrestin 2 [91]. CBDV has poor oral bioavailability; however, its liposoluble properties enable it to cross the blood-brain barrier [78] efficiently.




Figure 4. Receptors mediating CBDV's agonistic and antagonistic actions. CBDV, a non-psychoactive cannabinoid, exhibits both agonistic and antagonistic activity on several receptors, contributing to its potential therapeutic benefits in conditions like epilepsy, developmental disorders, and inflammation. Unlike THC, CBDV has a low affinity for CB1 and CB2 receptors [86]. It is thought to act more as a modulator of the endocannabinoid system rather than directly engaging as an agonist or antagonist at these receptors. This indirect modulation could influence the signaling pathways mediated by endocannabinoids, contributing to its anticonvulsant properties [92]. CBDV significantly affects TRPV1 and TRPV2, where it acts as an agonist [72]. These channels regulate pain, inflammation, and sensory perception, making CBDV potentially useful in managing neuropathic pain and inflammatory conditions. It is unknown whether CBDV activates other receptors such as α 2-adrenoreceptors, PPARs, and 5-HT_{1A}. CBDV's receptor specificity highlights its therapeutic potential, particularly in epilepsy, neuroinflammation, and pain management, making it a promising candidate for further medical research. CBD, Cannabidiol; CBDV, Cannabidivarin; THC, Δ^9 -tetrahydrocannabinol; TRPV, Transient receptor potential vanilloid; PPARs, Peroxisome proliferator-activated receptor. \rightarrow , agonist; \dashv , antagonist.

CBDV exhibited an antiseizure effect in animal models and a favorable safety profile in Human studies [93,94]. In *MECP2*-308 mouse models, CBDV improved sociability, brain weight, and overall health and partially improved motor function [95,96]. In rat studies, CBDV significantly reduced pentylenetetrazol (PTZ) seizure severity and mortality [93,97]. However, CBDV did not affect the severity of pilocarpine convulsions. Further, CBDV significantly suppressed PTZ seizures, decreased seizure severity, and rescued PTZ-induced gene expression [98]. In another study, long-term CBDV administration in RTT mice significantly improved brain weight without affecting the neurotrophin levels [95]. None of these *in vivo* studies used antagonist experiments further to elucidate the mechanism of CBDV's anticonvulsant effects. In an adult focal epilepsy cohort, CBDV exhibited safety in humans and significantly reduced seizure frequency [99]. In a pediatric RTT cohort, CBDV significantly improved seizure control in children with *MECP2*-related RTT [100]. Future studies are warranted to establish the specific mechanism CBDV offers for neuroprotective effects.

CBG

CBG (Figure 5) activates α_2 -adrenoceptors [101], PPAR γ [102], binds to cannabinoid CB1 and CB2 receptors [101], and blocks CB1 and 5-HT $_1A$ receptors [103] and has been shown to inhibit sodium channel currents in vitro; however, it was ineffective as an anticonvulsant in a PTZ-induced acute seizure model [104]. Furthermore, CBG did not prevent hyperthermia-induced seizures in a Scn1a $^{+/-}$ mouse model of Dravet syndrome, which involves mutations affecting Nav channels [105]. In stably transfected HEK cells and primary dorsal root ganglion (DRG) neurons expressing Nav channels, CBG acted as a low-affinity inhibitor of sodium channels [106]. Another study found that CBG produced modest inhibitory effects on peak currents elicited by this subset of sodium channels in human voltage-gated sodium channel subtypes expressed in mammalian cells [107]. These findings suggest that CBG may induce neuronal hyperexcitability. However, more in vivo studies are warranted to explore its neuroprotective mechanisms in controlling epileptic seizures.




Figure 5. Receptors mediating CBG's agonistic and antagonistic actions. CBG, often referred to as the "mother cannabinoid" due to its role as a precursor to other cannabinoids like THC and CBD, exhibits a diverse pharmacological profile, interacting with various receptors and modulating multiple physiological pathways. CBG is considered a partial agonist at both CB1 and CB2 receptors [101]. However, its affinity for these receptors is relatively low compared to THC. Its interaction with CB1 is thought to be non-psychoactive and may even counteract some of the psychoactive effects of THC, which acts as a full agonist. CBG's CB2 agonism has been linked to immunomodulatory and anti-inflammatory effects, which may aid in its therapeutic potential in conditions like neuroinflammation. CBG acts as an agonist at particularly TRPV1/2 [90], which is involved in pain, inflammation, and thermoregulation. CBG can activate alpha (2)-adrenoceptors, and block 5-HT $_1A$ receptors [103]. CBG activates PPAR γ [102], which are involved in lipid metabolism, inflammation, and cell differentiation. CBG's receptor interactions highlight its diverse pharmacological profile, making it a likely candidate for therapeutic applications in pain management, neuroprotection, mood disorders, and inflammation. Its ability to act as both an agonist and antagonist across different receptor systems contributes to its potential to treat a wide range of medical conditions. CBG, Cannabigerol; THC, Δ^9 -tetrahydrocannabinol; TRPV, Transient receptor potential vanilloid; PPAR γ , Peroxisome proliferator-activated receptor- γ . →, agonist; ⊥, antagonist.

Table 1. Neuroprotective functions of NMPs in epilepsy models.

Model	NMPs	Effect	Reference
Pilocarpine-epilepticus rat	CBD	Convulsant ↓ Neurodegeneration ↓	[49]
PTZ seizures	CBD and CBG	Nav current in cells ↓	[104]
Epilepsy-spontaneous LFPs in Cells	CBDV	Amplitude and duration of LFPs ↓ Mg ²⁺ free induced LFPs frequency ↑	[93]
Epilepsy in transfected cells (TRPV1, TRPV2, and TRPA1.	CBDV +CBD	Convulsant ↓ Phosphorylation of TRPV1 at the S800 site ↑	[108]
Electrophysiology (epileptiform bursting) (In vitro)	Δ ⁹ -THCV	Epileptiform burst ↓	[81]
PTZ seizures	CBDV	Seizure severity ↓ Latency to first signs of seizure ↑	[93,97,98]
PTZ seizures	Δ ⁹ -THCV	Median seizure severity, duration, progression, or latency was unaffected	[81]
6-hydroxytryptamine or LPS in rats and mice	Δ ⁹ -THCV	Neuronal loss, microglial activation, ↓ TH positive neurons and Motor activity↑	[109]
Rat model	CBD	Convulsant ↓ Seizure severity ↓	[61]
Rats (GEPR-3) strain	CBD	Seizure ↓	[67]

CBD, Cannabidiol; CBDV, Cannabidiol; CBG, Cannabigerol; THCV, Δ⁹-tetrahydrocannabivarin; LFS, local field potentials; PENK, proenkephalin; PTZ, pentylenetetrazole; LPS, lipopolysaccharide; TRPV, transient receptor potential vanilloid; TH, Tyrosine hydroxylase; GEPR-3, genetically epilepsy-prone rat; ↓, reduced; ↑, enhanced.

Role of NMPs in Parkinson's Disease

Parkinson's disease (PD) is the second most dominant neurodegenerative condition, affecting approximately 5% of individuals over the age of 85. Recently, the global burden of PD has doubled, making it one of the fastest-growing neurodegenerative diseases. Both genetic and environmental factors contribute to the risk of developing PD [110]. The PD is characterized by motor symptoms such as tremors, rigidity, bradykinesia, and a range of non-motor symptoms [111]. Motor symptoms are strongly associated with the damage and dysfunction of dopaminergic neurons in the nigrostriatal pathway, particularly during the intermediate stages of the disease [112]. This damage is often linked to the presence of α-synuclein, a protein involved in synaptic vesicle release, which can misfold and form aggregates known as Lewy bodies [113]. Developing α-synuclein aggregates can take more than 20 years, indicating that other neurotransmission systems may contribute to early non-motor symptoms, such as mild olfactory and cognitive impairments and depression [111,114]. eCBs, acting via CB1 and CB2 receptors, regulate synaptic and motor functions and can impact striatal rearrangement following dopamine depletion [115–117]. In PD subjects, cerebrospinal fluid exhibits heightened levels of AEA [117], and a reduction in CB1 receptors is observed in the substantia nigra

[118]. Conversely, increased CB1 receptor expression is found in the nigrostriatal, mesolimbic, and mesocortical dopaminergic projection areas [118]. CB1 receptors also appear to be involved in the action of 3,4-dihydroxy-L-phenylalanine (L-DOPA); CB1 agonists have been shown to prevent the motor fluctuations commonly observed during L-DOPA therapy [119]. Additionally, increased CB2 receptor expression has been found in glial cells in postmortem tissues of PD subjects [120], suggesting a role in the inflammatory aspects of the disease. Given the role of the eCB system in PD, pharmacological manipulation of the eCB system may represent a potential therapeutic approach for managing the disease.

CBD

A recent meta-analysis suggested that treatment with CBD significantly improves PD symptoms [121,122]. Nabilone has also been shown to be effective in alleviating anxious moods and night-time sleep problems [123], which are non-motor symptoms of PD. Additionally, CBD exhibits neuroprotective effects on the nigrostriatal pathway [124,125], and in vitro studies indicate its ability to activate tropomyosin receptor kinase A [126]. The anti-apoptotic effects of CBD are mediated by the ERK and Akt/mTOR pathways, with ERK activation modulated by TRPV1 and CB2 receptors [53]. Further studies have demonstrated that CBD exerts anti-apoptotic effects on dopaminergic neurons and reduces neuroinflammation by inhibiting the expression of NLRP3/caspase-1/IL-1 β , Bax, and caspase-3 while upregulating Bcl-2 [127]. Therefore, the neuroprotective effects of CBD in PD appear to be mediated by CB2, but not CB1, receptors [53,124]. CBD prevented memory impairments and decreased despair-like behavior that was induced by bilateral intraoral 6-OHDA lesions [128]. In addition, CBD prevented dopaminergic neuronal loss in the striatum, ventral tegmental area, substantia nigra pars compacta (SNpc) and a reduced hippocampal neurogenesis, reduced the mortality rate and decreased neuroinflammation in 6-OHDA-lesioned rats [128]. Repeated treatment with CBD favored the neuronal maturation of newborn neurons in the hippocampus in Parkinsonian rats [128]. Although these studies provide evidence of the therapeutic potential of CBD, further research is warranted before considering CBD for the treatment of PD.

THCV

THCV attenuated the motor impairments -induced by 6-hydroxydopamine, likely through changes in glutamatergic transmission [109]. Also, chronic administration of THCV rescued the loss of tyrosine hydroxylase-positive neurons caused by 6-hydroxydopamine in the substantia nigra, as antioxidant effects [109]. Additionally, THCV was protective against 6-hydroxydopamine (6-OHDA) or LPS-induced motor impairments similar to rimonabant in rats [109]. Chronic THCV impaired microglial activation and preserved nigrostriatal dopaminergic neurons after 6-OHDA administration and in the LPS model of PD. Additionally, THCV preserved TH-positive neurons, possibly through CB2 receptors [109]. Also, THCV antagonized the effects of the CB1 receptor agonist, CP55,940, indicating its effects as an antagonist of the CB1 receptor. In Pitx3ak mutant mice, THCV rescued abnormal involuntary movements and reduced the FosB protein and the histone pACh3, which had previously been found to be enhanced in the basal ganglia in L-DOPA-induced dyskinesia [129]. While future investigations are warranted to examine the clinical significance of THCV in humans, the findings, however, establish THCV as a promising agent for developing a nonpsychotropic cannabinoid-based therapy for patients with PD.

Table 2. Neuroprotective functions of NMPs in PD models.

Model	NMPs	Effect	Reference
PD Patients	CBD	PD symptoms ↓	[121]
PD Patients	CBD	Anxiety, tremor amplitude ↓	[122]
Unilateral lesions rat model	CBD	hydroxydopamine-induced DA depletion	[124]

Sprague–Dawley rats	CBD	Neuroprotection ↑	[125]
PC12 cells	CBD	Cell viability, differentiation, axonal (GAP-43), synaptophysin, and synapsin I ↑	[126]
SH-SY5Y cells	CBD	Cell viability↑ Apoptosis, Bax, and caspase 3. Moreover, nuclear PARP-1↓	[53]
Mice	CBD	Cognitive dysfunction, TNF- α , IL-1 β , IL-6, Bax and Caspase-3 and NLRP3/caspase-1/IL-1 β inflammasome pathway↓ Locomotion, 5-HT, DA, IL-10, TH, Bcl-2 ↑	[127]
Rats	CBD	SNpc, mortality rate, hippocampal neurogenesis, despair- behavior, Memory impairments, neuroinflammation↓ Neuronal maturation ↑	[130]
Mice	CBG	Motor tests , LAMP-1, TNF- α , IL-1 β , nitric oxide synthase and COX2↓	[131]
SH-SY5Y cells Mice (unilaterally lesioned)	CBG	Cytoprotection, GFAP, CD68 ↓ Motor activity ↑	[132]
SH-SY5Y cells Mice (unilaterally lesioned)	CBG	TH positive neurons, Motor activity ↑	[133]
Rat	Δ^9 -THCV	Motor activity, TH positive neurons ↑	[109]
Pitx3 ^{ak} mutant mice		AIMs, horizontal and vertical activities, FosB, pACh3, and dyskinesia ↓	[129]
C. elegans	CBDV	α -syn, DAergic neurons ↓	[134]

CBD, Cannabidiol; CBDV, Cannabidivarin ; DA, THC, Tetrahydrocannabinol; DA, Dopamine; CBG, Cannabigerol; TH, Tyrosine hydroxylase; GFAP, Glial fibrillary acidic protein; THCV , Δ -9-tetrahydrocannabivarin (THCV); THC, Δ -9-tetrahydrocannabinol; SH-SY5Y is a thrice-subcloned cell line derived from the SK-N-SH neuroblastoma cell line; C. elegans, Caenorhabditis elegans; FosB, FosB Proto-Oncogene, AP-1 Transcription Factor Subunit; pACh3 phosphoacetyl H3; CD68, Cluster of Differentiation 68; LAMP-1, Lysosome-associated membrane glycoprotein 1; TNF- α , Tumor necrosis factor alpha; IL-1 β , Interleukin 1 Beta; IL-10; Interleukin 10; IL-6, Interleukin 6; COX2, Cyclooxygenase 2; Bax, Bcl-2-associated X protein; NLRP3, Nod-like receptor protein 3; 5-HT, 5-hydroxytryptamine; Bcl-2, B-cell lymphoma-2, PARP-1, Poly (ADP-ribose) polymerase 1; SNpc, Substantia Nigra Pars Compacta; GAP-43, Growth-associated protein 43. Pitx3^{ak} mice, Microphthalmia and aphakia mice; PC12 cells, Rat adrenal tumor cells; ↓, reduced; ↑, enhanced.

CBDV

CBDV has been shown to block α -synuclein (α -syn) aggregation via DAF-16, a key transcription factor involved in regulating oxidative stress, in a transgenic *Caenorhabditis elegans* (C. elegans) model [134]. Additionally, CBDV protects dopaminergic neurons from injury and degeneration induced by 6-OHDA [134]. Further exploring the molecular targets through which CBDV ameliorates α -synuclein aggregation in vivo would be valuable for understanding its therapeutic potential in PD. CBDV blocked α -synuclein (α -syn) aggregation via DAF-16, a key transcription factor for regulating

oxidative stress in the transgenic *C. elegans* model [134]. CBDV prevented dopaminergic neurons from 6-OHDA-induced injury and degeneration [134]. The molecular target (s) for CBDV ameliorating α -synuclein aggregation in vivo would be deserving of exploration in the future to understand its therapeutic potential of PD.

CBG

CBG has recently been investigated for its neuroprotective properties in inflammatory models of PD in mice. In one study, CBG improved motor function and rescued the loss of tyrosine hydroxylase-containing nigrostriatal neurons in mice that were unilaterally lesioned with lipopolysaccharide (LPS) [131]. Additionally, CBG reduced the elevated levels of LAMP-1 immunolabeling (a marker for autophagy impairment) and proinflammatory mediators such as tumor necrosis factor- α , interleukin-1 β , inducible nitric oxide synthase, and cyclooxygenase-2, and glial reactivity caused by the LPS lesion [131]. In another study using a neurotoxin model of PD, CBG exhibited cytoprotective effects in cultured SH-SY5Y cells treated with 6-hydroxydopamine (6-OHDA) [132]. Additionally, in mice unilaterally lesioned with 6-OHDA, CBG rescued tyrosine hydroxylase (TH)-positive nigral neurons from 6-OHDA-induced damage. It also completely prevented astroglial (GFAP) and microglial (CD68) reactivity in the substantia nigra of lesioned mice, leading to recovery from the motor deficiencies caused by 6-OHDA [131,133]. Furthermore, CBG treatment restored dopamine levels and its metabolite DOPAC in the striatum of 6-OHDA-lesioned mice [132]. These neuroprotective effects were attributed to CBG's activity at the peroxisome proliferator-activated receptor- γ (PPAR- γ) rather than the CB2 [133].

Role of NMPs in Alzheimer's Disease

Alzheimer's disease (AD) is a chronic neurodegenerative condition that affects the central nervous system, leading to a decline in cognitive functions. Similar to other neurodegenerative diseases, AD involves multiple pathologies influenced by a combination of genetic and environmental factors. Several potential causes are associated with the onset of AD, including the deposition of beta-amyloid ($A\beta$) aggregates forming senile plaques, hyperphosphorylation of Tau protein resulting in neurofibrillary tangles (NFTs), oxidative stress, neuroinflammation due to microglial activation, metabolic dysfunction, and obesity (For references see [135]).

Despite numerous pharmacological strategies designed to slow down AD symptoms, these approaches have largely been ineffective. In AD patients, senile plaques have been shown to express CB1 and CB2 receptors along with microglial activation markers [136]. However, the number of CB1-positive neurons abundant in healthy individuals is significantly reduced in areas with activated microglia. Additionally, G protein coupling and CB1 receptor protein expression are markedly diminished in AD brains [136]. Furthermore, enhanced CB2 levels are found in AD patients and correlated with brain $A\beta$ 42 levels and senile plaque score [137]. The marked eCBs changes, as found in AD patients, were also observed in several AD mouse models [138–140]. These findings indicate that cannabinoid receptors play a vital role in the development of AD pathology and suggest that cannabinoids lacking psychiatric effects may have the potential to prevent the neurodegenerative process in AD.

CBD

In pre-clinical studies, CBD exhibited neuroprotection against many aspects of AD pathology. For example, the application of CBD prevented $A\beta$ -peptide toxicity in PC12 cells by inhibiting oxidative stress, caspase-3 activation [141], and Tau hyperphosphorylation [142]. Additionally, CBD downregulated p-GSK-3 β , an inhibitor of the Wnt pathway, causing enhanced Wnt/ β -catenin signaling and stimulation of PPAR γ and amyloid precursor protein (APP) ubiquitination. In the brain, Wnt/ β -catenin signaling is crucial for neuronal survival and neurogenesis. It regulates synaptic plasticity and blood-brain barrier integrity and function [143]. Furthermore, CBD attenuated oxidative stress and rescued mitochondrial function [142]. In mesenchymal stem cells, CBD exposure

reduced genes coding for the kinases responsible for Tau phosphorylation and the secretase enzymes involved in A β production [144]. Studies have shown that CBD rescues the formation of NFTs and prevents neuronal apoptosis via functioning as an endogenous cannabinoid receptor agonist [145].

In adult mice, intraventricular (i.v.) injection of fibrillar A β caused IL-6 gene expression and spatial memory deficits, which were rescued by CBD injection for two weeks [146]. In APP x PS1 mice, CBD administration rescued cognitive deficits measured by object recognition and social recognition memory without affecting anxiety behavior [147]. In a similar study, CBD was provided as pellets to APP x PS1 mice at the age of 2.5 months for 8 months and could prevent the development of social recognition memory deficits without affecting A β load, oxidative damage, or anxiety behavior [148]. In 5xFAD mice, chronic, low-dose CBD administration significantly improved memory and enhanced insoluble form of A β levels in the cortical regions [149]. Oral administration of CBD for an extended period reversed pathophysiology and rescued social recognition memory deficits in the APP x PS1 mouse model for AD [150]. CBD administration in the senescence-accelerated mouse prone 8 (SAMP8) model, improved cognitive function, increased hippocampal-activated microglia shift from M1 to M2, and restored gut microbial functions [151]. Additionally, CBD stimulates autophagy signal transduction through crosstalk between ERK1/2 and AKT kinases [52], which are key cell proliferation and survival regulators and have been implicated in AD pathology. In a randomized, double-blinded, placebo-controlled trial, CBD improved behavioral symptoms in AD [152]. Future studies are warranted to address research issues related to CBD's safety, efficacy, and variability. In this direction, nanoparticle-coated CBD exhibited better beneficial effects on rescuing learning and memory, increasing hippocampal CB1 and CB2 receptors, and reducing amyloid plaques in an AD rat model [153]. Additionally, administering CBD along with THC, a combination known to provide more significant therapeutic benefits than phytocannabinoids alone [154], was shown to improve behavior problems and rigidity in severely demented patients [155].

Table 3. Neuroprotective role of NMPs in AD models.

Model	NMPs	Effect	Reference
PC12 cells	CBD	Cell Survival \uparrow ROS, lipid peroxidation and Caspase 3 \downarrow	[141]
PC12 cells	CBD	Wnt/ β -catenin \uparrow Tau hyperphosphorylation, p-GSK-3 β \downarrow	[142]
MSC cells	CBD	GSK3 β , CDK5, DYRK1A, CAMK2A, MAPK1, MAPK12, and MAPK14, BACE1 \downarrow ,	[144]
N13 microglial cells Rat primary microglia Mice	CBD	Intracellular calcium \downarrow Nitrite generation, IL-6 gene expression \downarrow Spatial memory, Microglial migration \uparrow	[146]
APPxPS1 mice	CBD	social recognition and novel object recognition \uparrow	[147]
A β PPSwe/PS1 Δ E9	CBD	social recognition \uparrow	[148]
5x FAD mice	CBD and THC	Spatial memory, beta amyloid \uparrow	[149]
SAMP8 mice	CBD	Bacterioidetes, hippocampal activated microglia shift from M1 to M2 \uparrow and LPS \downarrow	[151]
Male wistar rats	CBD coated by nano-chitosan	learning and memory, CB1 and CB2 protein expression \uparrow Amyloid plaques \uparrow	[153]

Female patients	CBD	Direct contact, behavior ↑	[155]
MC65 cell	CBG	A β aggregation ↓	[156]
PC12 cells	CBG	A β aggregation, A β ₁₋₄₂ neurotoxicity ↓ Neuroprotection	[157]
Male human subjects	THCV	Memory impairment↓	[158]
In vitro assay	CBD, CBDV, CBG	AChE and BChE ↓	[159]

CBD, Cannabidiol; CBDV, Cannabidivarin ; THC, Tetrahydrocannabinol; DA, Dopamine; CBG, Cannabigerol; THCV, Δ^9 -tetrahydrocannabivarin; TH, Tyrosine hydroxylase; 5HT, 5-hydroxytryptamine; IL-10, Interleukin-10; IL-1 β , Interleukin-1 beta; Interleukin-6, Bcl2, Protein that regulates programmed cell death; COX-2, Cyclooxygenase-2; AIMS, abnormal involuntary movements; ROS, Reactive oxygen species, MSCs, Mesenchymal stem cells; LPS, Lipopolysaccharide; AChE, Acetylcholinesterase; BChE, butyrylcholinesterase; BDNF, brain-derived neurotrophic factor; PGC1 α , Peroxisome proliferator-activated receptor gamma coactivator 1-alpha; CDK5, Cyclin dependent kinases 5 gene; DYRK1A, Dual specificity tyrosine phosphorylation regulated kinase 1A; CAMK2A, Calcium/ calmodulin dependent protein kinase II α ; MAPK1, Mitogen activated protein kinase 1; MAPK12, Mitogen activated protein kinase 12; MAPK14, Mitogen activated protein kinase 14; BACE, β -Secretase 1; PC12 cells, Rat adrenal tumor cells; MSC, Multipotent stromal cells; N13 cells, primary mice microglial cells; APP, Amyloid precursor protein; PS1, Presenilin 1; FAD, 5 familial AD; SAMP8, senescence accelerated mouse prone 8; MC65, Human CNS nerve cell line; HT22, Mouse hippocampal Neuron cell; ↓, reduced; ↑, enhanced.

THCV

While THCV has not yet been tested in AD models, it has demonstrated a strong safety profile in healthy adults. In a placebo-controlled, double-blind, crossover pilot study, THCV was shown to reverse THC-induced memory impairment [158]. Future studies using THCV in AD animal models could provide potential neuroprotective mechanisms and therapeutic benefits.

CBDV

CBDV has demonstrated efficacy in an in vitro AD model by blocking oxytosis and energy loss in HT22 cells and reducing A β toxicity and trophic withdrawal. However, these experiments did not investigate the specific mechanisms underlying these effects [156]. Additionally, CBDV was found to inhibit the activities of acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) [159]. Abnormal hyperactivity of AChE and BChE contributes to cholinergic deficiency, associated with several neurological disorders, including the memory impairments and cognitive decline observed in AD [160]. Therefore, further research to elucidate the molecular mechanisms of CBDV in AD models could provide valuable insights into its therapeutic potential for this condition.

CBG

CBG has been shown to reduce the accumulation of A β deposits in vitro. Specifically, CBG cleared the preformed and aggregated A β from neurons and stimulated A β degradation in the MC65 cell line [156]. This widely recognized human neuron-like cell line lacks CB1 or CB2 receptors extensively used for proteotoxicity studies [156]. Additionally, CBG inhibited A β ₁₋₄₂ -induced neurotoxicity and morphological changes such as reduced neuritic projections and rounded cell morphology in PC12 cells [157]. Future studies with CBG in AD animal models could help elucidate this phytocannabinoid's neuroprotective mechanism.

NMPs Role in Huntington's Disease

Huntington's disease (HD) is a progressive neurodegenerative condition characterized by abnormal motor and cognitive functions. HD is triggered by mutations in the huntingtin gene (Htt) on chromosome 4, involving multiple cytosine-adenine-guanine (CAG) repeats at the exon 1 of the Htt gene [6–8]. Current pharmacological treatment for HD primarily utilizes atypical anti-psychotic drugs to manage hypermotor activity symptoms. However, the exact mechanisms underlying HD pathogenesis remain largely unclear, and there is currently no specific drug available to address the cognitive impairments associated with the disease [161]. Early in HD progression, research has revealed reduced CB1 mRNA and protein levels in medium spiny projection neurons of the caudate and putamen [162–164]. Moreover, CB1 transcription is suppressed by mutant huntingtin protein (mHtt) [165,166]. These changes in CB1 functionality significantly contribute to the cognitive, behavioral, and motor deficits observed in animal models of HD [167,168]. Consequently, there is increasing interest in pharmacological strategies to enhance CB1 signaling as a potential therapeutic approach for treating and managing HD.

CBD

Recent studies have evaluated the neuroprotective role of CBD in vitro using models of medium spiny projection neurons expressing mutant huntingtin protein (STHdhQ111/Q111) compared to wild-type (STHdhQ7/Q7). In these models, CBD enhanced CB1 expression and GABA release and promoted CB1-independent but 5HT1A-dependent phosphorylation of CREB (p-CREB). Additionally, CBD administration in rats with 3-nitropropionic acid (3NP)-induced HD reduced CB1 receptor expression and insulin-like growth factor 1 (IGF-1) while enhancing calpain expression in striatal neurons [60]. This treatment also restored levels of GABA, substance P, and neuron-specific enolase, which are involved in generating proinflammatory markers that contribute to neuronal atrophy. Furthermore, CBD reinstated antioxidant enzyme SOD-1 and proenkephalin levels in the striatum and substantia nigra, which play crucial roles in HD pathogenesis [60]. However, despite these insights into CBD's protective function in HD, its therapeutic utility remains controversial. While some studies indicate no improvement in animal models and human trials [169,170], others have reported beneficial effects in animal models [60,171].

Table 4. Neuroprotective functions of NMPs in different HD models.

Model	NMPs	Effect	Reference
STHdh(7/7) cells	ACEA, mAEA	ATP production, BDNF-2, PGC1 α CB1 mRNA levels \uparrow	[165]
Rats	CBD	mRNA SP, mRNA NSE, mRNA SOD-2 \uparrow	[60]
Rats	CBD	3NP-induced GABA, Nissl-stained neurons, CB1 and IGF-1 expression and SOD-1 expression \downarrow Calpain expression \uparrow	[171]
RBL-2H3 cells	CBG	Human TRPV1, rat TRPV2 \downarrow	[89]
HT29 cells	CBG	COX-2 enzyme, prostaglandins \downarrow	[172]
Mice	CBG	Reactive microgliosis, expression of COX-2, iNOS, TNF- α , <i>Cd44</i> , <i>Sgk1</i> \downarrow PPAR γ , catalase, SOD and GSH \uparrow	[173]
NSC-34	CBG	<i>HAP1</i> , <i>SLC32A1</i> , <i>ADCY5</i> , <i>AKT</i> , <i>ATF4</i> , <i>DLGAP1</i> , <i>DRD4</i> , <i>GNB4</i> , <i>PRKCA</i> \uparrow <i>ADCY9</i> , <i>CAMK2B</i> , <i>CLOCK</i> , <i>CREB1</i> , <i>DRD2</i> , <i>GNAL</i> , <i>PLD1</i> , <i>PPP3R1</i> , <i>PRKCB</i> , <i>SHANK1</i> , <i>SLC1A2</i> , <i>SLC18A1</i> , <i>SLC38A1</i> \downarrow	[174]

CBD, Cannabidiol; CBG, Cannabigerol; ACEA, Arachidonyl-2'-chloroethylamide hydrate; mAEA, Methanandamide; 3-NP, 3-Nitropropionic acid; SOD-1, Superoxide dismutase-1; SOD-2, Superoxide dismutase-2; SP, Substance P; NSE, Neuron-specific enolase; TRPV1, Transient receptor potential vanilloid 1; TRPV2, Transient receptor potential vanilloid 2; COX-2, Cyclooxygenase enzyme 2; iNOS, Inducible nitric oxide synthase; TNF- α , Tumor necrosis factor alpha; IL-6, Interleukin-6; PPAR γ , peroxisome proliferator-activated receptor- γ ; *cd44*, Non-kinase transmembrane glycoprotein gene; *SGK1* Serum/Glucocorticoid Regulated Kinase 1 gene; GSH, Glutathione; *SLC32A1*, solute carrier family 32 member 1 gene; *ADCY5*, Adenylate cyclase 5 gene, *AKT*, Serine /threonine kinase 1 gene; *ATF4*, Activating transcription factor4 gene; *DLGAP1*, DLG associated protein 1 gene; *DRD4*, Dopamine receptor D4 gene; G protein, Guanine nucleotide binding protein gene; *GNAI2*, Alpha inhibiting 2 gene; *GNB4*, Beta 4 gene; *HAP1*, Huntingtin-associated protein 1 gene; *PRKCA*, Protein kinase C, alpha gene; *ADCY9*, Adenylate cyclase 9 gene; *CAMK2B*, Calcium/calmodulin-dependent protein kinase II, beta; *CLOCK*, Circadian locomotor output cycles kaput; *CREB1*, AMP responsive element binding protein 1; *DRD2*, Dopamine receptor D2; *GNAL*, Guanine nucleotide binding protein, alpha stimulating, olfactory type; *PLD1*, Phospholipase; *D1PPP3R1*, Protein phosphatase 3, regulatory subunit B, alpha isoform; *PRKCB*, Protein kinase C, beta; *SHANK1*, SH3 and multiple ankyrin repeat domains 1; *SLC1A2*, Solute carrier family 1 (glial high affinity glutamate transporter), member 2; *SLC18A1*, solute carrier family 18 (vesicular monoamine), member 1; *SLC38A1*, solute carrier family 38, member 1; STHdh(7/7) cells, Cell line derived from the striatum of a mouse; RBL-2H3 cells, Basophilic leukemia cell line; HT29, Colon Cancer Line; NSC-34, motor neuron-like cells; ↓, reduced; ↑, enhanced.

THCV

Therapeutic studies of cannabinoid-based agents in HD animal models suggest that CB1 and endovanilloid receptor agonists [175,176] and AEA reuptake inhibitors [177] can prevent hyperkinesia in the early phases of HD. This is likely due to the gradual loss of CB1 receptors [178] in the advanced stages of the disease. Although further studies are warranted, the potential use of THCV in HD is limited since it functions as an antagonist of CB1 and CB2 receptors. An improved understanding of the eCB system in HD may help identify specific NMPs or combinations that provide therapeutic or neuroprotective benefits in patients with HD.

CBDV

While phytocannabinoids show promise as potential treatments for motor-related diseases, no studies have evaluated the neuroprotective effects of CBDV in models of HD. Future research is needed to explore neuroprotective mechanisms within the relevant brain regions in animal models treated with CBDV.

CBG

CBG is a biologically active constituent of the marijuana plant, present in much smaller quantities compared to other cannabinoids. It acts as a precursor to various phytocannabinoids. CBG is a non-psychoactive compound that exhibits a wide range of biological activities [179] and is potentially a therapeutic compound for treating diseases requiring multidirectional pharmacotherapy. CBG interacts with CB1, CB2, TRPV1, and PPAR receptors [88,180,181]. Additionally, CBG suppresses the activity of FAAH, an enzyme that metabolizes anandamide (AEA), affecting its levels and biological effects. However, compared to CBD, CBG is less effective as an FAAH inhibitor [89,182]. Furthermore, CBG reduces the activity of DAGL, the enzyme responsible for the biosynthesis of 2-AG, and inhibits the activities of COX-1 and COX-2, which metabolize polyunsaturated fatty acids (PUFAs), mainly arachidonic acid, into lipid mediators [172,183,184]. CBG also acts as a neutral 5-HT1A receptor antagonist, a CB1 receptor antagonist, and a potent α 2-adrenoceptor agonist [103].

CBG exhibits similar pharmacokinetic (PK) profiles in rats and mice, though it shows slower brain penetration in mice. Both species have higher concentrations of CBG following intraperitoneal (i.p.) injection compared to oral administration; however, in rats, this does not correspond to higher concentrations in brain tissue [185]. CBG has demonstrated neuroprotective effects in experimental

HD models through both cannabinoid receptor-dependent and independent mechanisms. For instance, in an *in vivo* model of HD using 3-NP, CBG significantly reduced the expression of upregulated inflammatory markers such as COX-2, iNOS, IL-6, and TNF- α [173] and prevented the 3-NP-induced neuronal loss. Additionally, CBG reversed the activities of antioxidant enzymes, including catalase, superoxide dismutase (SOD), and glutathione (GSH), compared to control groups. CBG also rescued the expression of down-regulated genes directly related to HD, such as *sgk1*, *Cd44*, and huntingtin-associated protein-1, while reducing mutant huntingtin (mHTT) protein aggregation and improving motor function [173]. Moreover, CBG was found to activate PPAR γ dose-dependently in cultured striatal cells containing both wild-type and mutant huntingtin [173]. A recent transcriptomic study in motor neuron-like cells (NSC-34) revealed that CBG reduced the expression of genes involved in glutamate release, enhanced the expression of genes related to GABA release, and upregulated the dopamine D4 receptor and its downstream effectors [174], suggesting CBG's influence on neurotransmission pathways. Future studies are needed to establish the link between these transcriptomic changes and behavioral and neuronal signaling to understand the role of CBG in neuroprotection better.

Neuroprotective Role of NMPs in Substance and Alcohol Use Disorders

Cannabinoid receptors are densely enriched in brain areas related to reward function and developing and maintaining addictive behaviors [186]. The eCB system has been implicated in the pathophysiology of addiction by modulating pathways that affect drug and alcohol-seeking behaviors, cravings, withdrawal [24,187,188], and memory and emotional processes [189]. Alcohol [190,191] and drug abuse [192,193] act as one of the environmental factors that promote many of the neurodegenerative disorders. Therefore, studies have explored whether phytocannabinoids provide neuroprotection against substance and AUD-related disorders. In the following sections, we discuss the current evidence on the neuroprotective functions of phytocannabinoids on substance and alcohol use disorders.

Substance Use Disorders (SUD)

In a randomized, double-blind, placebo-controlled study, CBD significantly reduced cigarette smoking, although it did not affect cravings [194]. Similarly, another study found that CBD decreased the salience and pleasantness of cigarette cues without impacting cravings, withdrawal symptoms [195], or impulsivity [196]. In an open-label crossover study involving daily users of nicotine-containing e-cigarettes, CBD was shown to reduce both the severity of nicotine withdrawal symptoms and state anxiety during e-cigarette abstinence [197]. Additionally, CBD has been found to mitigate nicotine withdrawal and hyperalgesia-inducing effects in both mice [198] and rats [199]. While the potential of other phytocannabinoids to protect against nicotine use disorder remains unknown, future research is warranted to explore this possibility.

There has been substantial scientific discussion regarding the potential of CBD as a treatment for cannabinoid use disorder (CUD) [200,201]. In animal models, CBD has been shown to reduce spontaneous withdrawal symptoms in THC-dependent rodents [202–204]. In a case study, CBD alleviated cannabis withdrawal symptoms and improved anxiety and sleep function [204–206]. Additionally, CBD significantly reduced cannabis use in participants of a non-randomized open-label study [207]. Moreover, CBD mitigated adverse cognitive and mental health effects [202,203] in long-term THC users, including reduced cognitive deficits, psychotic-like and depressive symptoms, and increased hippocampal volumes [208] and resting-state functional connectivity [209]. These studies suggest that the antipsychotic and anxiolytic properties of CBD may improve psychological and cognitive functions related to long-term cannabis use—the use of other NMPs to combat the effects of THC warrants further study.

CBD has been studied in opioid use disorders and has been shown to mitigate morphine reward and reduce the reinstatement of heroin-seeking behavior in rats [210,211] as well as alleviate opioid withdrawal symptoms in rodents [212,213]. It also decreased naloxone-precipitated jumping behavior in mice [214]. CBD prevented the reinstatement of oxycodone-induced conditioned place

preference (CPP) and rescued recognition memory deficits in adolescent male mice. These mice exhibited increased CB1 receptors and a reduced mu-opioid receptor (MOR) expression in the prefrontal cortex (PFC) [215]. CBG was found to reverse oxaliplatin-associated mechanical sensitivity [216] and attenuate the acute antinociceptive effects of morphine through interactions with α 2-adrenergic, CB1, or CB2 receptors [214]. These findings suggest that continued research on CBD, CBG, and other NMPs could enhance our understanding of their potential in treating pain and substance use disorders.

Table 5. Neuroprotective functions of NMPs in different SUD models.

Model	NMPs	Effect	Reference
Humans	CBD (+THC)	Anxiety ↓	[200]
Humans	CBD (+THC)	Satiety ↑	[201]
Humans	CBD	Emotion, reward processing, effects of THC ↓	[203]
Human	CBD	Withdrawal, anxiety and dissociative symptoms ↓	[205]
Human	CBD	Anxiety, cannabis use ↓ Sleep ↑	[206]
Humans	CBD	Cannabis use ↓	[207]
Humans	CBD	Psychological symptoms ↓ Cognition ↑	[208]
Human	CBD	Functional connectivity ↑	[217]
Wistar rats	CBD	Morphine reward behavior ↓	[210]
Rat	CBD	heroin-seeking behavior, CB1R expression ↓	[211]
Mice	CBD	Anxiety, <i>Cnr1</i> , <i>Pomc</i> ↓ Motor activity, TH expression, ↑	[212]
Mice	CBD	Gastrointestinal symptoms, Jumping behavior ↓	[213]
Mice	CBD	Mechanical sensitivity, Jumping behavior ↓	[214]
Rats	CBD	Locomotor Hyperactivity, MOR ↓ Recognition Memory, CB1R expression ↑	[215]
Mice	CBG	Mechanical sensitivity, <i>Aif1</i> , <i>Ccl2</i> , <i>Calca</i> , and <i>Tlr4</i> ↓	[216]

CBD, Cannabidiol; THC, Tetrahydrocannabinol; CBG, Cannabigerol; TH, tyrosine hydroxylase; Pomc, Proopiomelanocortin; Cnr1, Cannabinoid receptor; MOR Mu-opioid receptor; *Aif1*, Allograft Inflammatory Factor 1; *Ccl2*, C-C Motif Chemokine Ligand. 2; *Calca*, Calcitonin Related Polypeptide Alpha; Tlr4, Toll-Like Receptor 4; ↓, reduced; ↑, enhanced.

Alcohol Use Disorders (AUD)

CBD has shown promise in preventing alcohol abuse and mitigating alcohol-related effects, owing to its anxiolytic properties [218, 219]. Preclinical studies strongly suggest that the eCB system plays a significant role in the motivational properties of alcohol and its effects [24, 220-222]. Specifically, antagonism of the CB1 receptor has been shown to suppress rodent alcohol consumption [24, 220-223]. As CBD may act as a negative allosteric modulator of the CB1 receptor, this supports the notion that CBD could be a promising treatment for AUD.

CBD has demonstrated protective effects against alcohol-induced neurodegeneration. When administered alongside binge ethanol exposure, CBD rescued neurodegeneration in the

hippocampus and entorhinal cortex [224]. Transdermal administration of CBD also significantly reduced neurodegeneration in the entorhinal cortex [225]. Furthermore, CBD has been shown to prevent alcohol-induced cell death and improve neuronal viability in rat hippocampal cultures [218,219][226]. In alcohol dependence models, CBD inhibited impulsive choice behavior [227,228], a factor linked to both alcohol use and the risk of relapse [229,230]. CBD has also been found to reduce alcohol consumption [204], motivation, and relapse in a two-bottle choice paradigm [231]. In these studies, CBD reduced ethanol-induced hypothermia and handling-induced convulsions without affecting blood ethanol concentration [231]. Moreover, CBD does not alter breath alcohol levels in humans [232]. CBD also reversed alcohol-induced changes in gene expression, such as tyrosine hydroxylase in the ventral tegmental area, *Oprm1*, *CNR1*, and *GPR55* in the nucleus accumbens (NAcc), while significantly enhancing CB2 receptor expression in the Nacc [231].

In selectively bred Sardinian alcohol-preferring (sP) rats, CBD significantly reduced lever responses for alcohol and the amount of self-administered alcohol [233]. Although the exact neuroprotective mechanisms of CBD require further investigation, these findings strongly suggest that CBD may be beneficial for treating AUDs. In another study, CBD significantly reduced alcohol withdrawal symptoms such as anxiety behavior and altered gene expression in related brain regions [234]. In the alcohol binge-like drinking model, CBD inhibited alcohol consumption and preference, normalized abnormal corticolimbic calcitonin gene-related peptide (CGRP) expression, and reduced reward and aversion-related hyper-responsivity and glucocorticoid levels in rats [235]. CBD also significantly mitigated the blunted stress response and corticosterone levels caused by binge-like alcohol exposure, restored dopamine transmission, and facilitated excitatory postsynaptic strength and remodeling in rats [236]. Additionally, CBD inhibited the development of tolerance to alcohol's hypothermic and sedative effects and restored reduced CB2R gene transcription in the striatum caused by ethanol [237]. In another alcohol withdrawal study, CBD reduced anxiety-like behaviors in mice [238]. However, some studies show limitations: CBD failed to attenuate alcohol-induced locomotor sensitization [239], and oral CBD did not prevent alcohol seeking, self-administration, or intake in baboons [240]. Although the precise mechanisms remain unknown, these studies indicate that CBD and potentially other phytocannabinoids could offer neuroprotection against alcohol toxicity and AUD.

Fetal Alcohol Spectrum Disorder (FASD) embodies a range of neurobehavioral impairments caused by prenatal and postnatal exposure to ethanol. Recent studies have shown that CBD treatment during the peri-adolescence period can mitigate the increased levels of TNF α and IL-6 in the hippocampus, as well as the cognitive deficits associated with pre- and postnatal alcohol exposure [241]. Data from four case studies suggest that CBD has neuroprotective effects, reducing FASD symptoms such as restlessness, aggression, agitation, impulsivity, and high scores on the Nisonger Disruptive Behavior Scale [242]. Additionally, CBD administration has been shown to reverse alcohol-induced gene expression defects, anxiety, depressive-like behaviors, and cognitive impairments in animal models exposed to alcohol both prenatally and postnatally [243]. Although the precise mechanisms are not yet fully understood, early research suggests that CBD, or possibly other phytocannabinoids, could provide neuroprotective effects against the developmental impact of alcohol exposure.

Table 6. Neuroprotective functions of NMPs in different AUD models.

Model	NMPs	Effect	Reference
Rat	CBD	Anxiolytic effect \uparrow	[218]
Rats	CBD	Social interaction \uparrow	[219]
Rats	CBD	Neurodegeneration \downarrow	[244]
Rats	CBD	Neurodegeneration \downarrow	[225]
Hippocampal cultures	CBD	Neuroprotection \uparrow	[226]
Mice	CBD	Alcohol intake,	[227]

		TH, Oprm1, CB ₁ R and GPR55 gene expression ↓	
Humans	CBD	BrAC ↓	[232]
sP Rats	CBD	Lever responses to self-administered alcohol ↓	[233]
Mice	CBD (+THC)	Locomotor sensitization ↓	[239]
Baboons	CBD	Alcohol seeking, self-administration, drinking patterns ↓	[240]
Mice (SAW model)	CBD	Rearings, groomings, anxiogenic behavior, <i>Cnr2</i> , <i>Oprm1</i> expression ↑ <i>Th</i> , <i>Pomc</i> gene expressions ↓	[234]
Rats	CBD	CGRP, alcohol consumption and preference ↓	[235]
Rats	CBD	Corticosterone ↓ DA, postsynaptic strength ↑	[236]
Rats	CBD	Hypothermic and sedation CB ₁ R, DRD ₁ , DRD ₂ mRNA ↓ CB ₂ R gene transcription ↑	[237]
Mice	CBD	Anxiety behavior, S100β and Iba1 ↓	[238]
Mice	CBD	Cognitive deficits, TNFα IL-6 ↑	[241]
Human	CBD	Disruptive behavior score ↓	[242]
Mice	CBD	Emotional cognitive disturbance ↓	[243]

CBD, Cannabidiol; THC, Tetrahydrocannabinol; Oprm1, Opioid Receptor Mu 1; GPR55, G Protein-Coupled Receptor 55; CB₁R, Cannabinoid Receptor 1, TH, tyrosine hydroxylase; BrAC, Breath alcohol level; sP, Sardinian alcohol-preferring; *Cnr2*, Cannabinoid Receptor 2 gene; SAW, Spontaneous alcohol withdrawal; CGRP, Calcitonin gene-related peptide; DA, Dopamine; DRD₁, Dopamine receptor D1; DRD₂, Dopamine receptor D2; S100B, calcium-binding protein; Iba, Ionized calcium-binding adaptor molecule 1; TNF-α, Tumor necrosis factor-alpha; IL-6, Interleukin-6, SAW, Spontaneous alcohol withdrawal; ↓, reduced; ↑, enhanced.

Conclusions

In summary, the therapeutic potential of *Cannabis sativa* extends well beyond the widely studied CBD, encompassing a diverse range of lesser-known phytocannabinoids that show promise in addressing various neurological disorders. The neuroprotective functions of these NMPs, particularly their antioxidant, anti-inflammatory, and immune-modulating properties, offer new avenues for research and treatment. While the pharmacological mechanisms of many NMPs remain underexplored, emerging studies suggest their potential to develop novel therapies for brain disorders. As research continues to unfold, these findings could pave the way for innovative cannabinoid plant-based treatments that go beyond the scope of traditional approaches, offering new hope in neuroprotection and disease management.

Author Contributions: B.S.B. and S.S. participated in the collection of literature, writing, and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported in part by NIH/NIAAA grant AA029686 (BSB).

Institutional Review Board Statement: All experiments were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals (National Institutes of Health Publication, 8th edition, revised 2011), as confirmed by the Nathan Kline Institute Ethical Committee.

Acknowledgments: This study was supported by funds from NIH/NIAAA.

References

1. Basavarajappa BS, Shivakumar M, Joshi V, Subbanna S. Endocannabinoid system in neurodegenerative disorders. *J Neurochem* 142(5), 624-648 (2017).
2. Ferrari R, Kapogiannis D, Huey ED, Momeni P. FTD and ALS: a tale of two diseases. *Curr Alzheimer Res* 8(3), 273-294 (2011).
3. Gibson SB, Figueroa KP, Bromberg MB, Pulst SM, Cannon-Albright L. Familial clustering of ALS in a population-based resource. *Neurology* 82(1), 17-22 (2014).
4. Gammon K. Neurodegenerative disease: brain windfall. *Nature* 515(7526), 299-300 (2014).
5. Kovacs GG. Concepts and classification of neurodegenerative diseases. *Handb Clin Neurol* 145 301-307 (2017).
6. Gonzalez-Alegre P. Recent advances in molecular therapies for neurological disease: triplet repeat disorders. *Hum Mol Genet* 28(R1), R80-R87 (2019).
7. Nopoulos PC. Huntington disease: a single-gene degenerative disorder of the striatum. *Dialogues Clin Neurosci* 18(1), 91-98 (2016).
8. Paulson H. Repeat expansion diseases. *Handb Clin Neurol* 147 105-123 (2018).
9. Taylor JJ, Williams NR, George MS. Beyond neural cubism: promoting a multidimensional view of brain disorders by enhancing the integration of neurology and psychiatry in education. *Acad Med* 90(5), 581-586 (2015).
10. Hampel H, Vergallo A, Caraci F *et al.* Future avenues for Alzheimer's disease detection and therapy: liquid biopsy, intracellular signaling modulation, systems pharmacology drug discovery. *Neuropharmacology* 185 108081 (2021).
11. Trinh NH, Hoblyn J, Mohanty S, Yaffe K. Efficacy of cholinesterase inhibitors in the treatment of neuropsychiatric symptoms and functional impairment in Alzheimer disease: a meta-analysis. *JAMA* 289(2), 210-216 (2003).
12. Bhushan M, Akash S, Pettarusp W. Adverse effects of medications used to treat motor symptoms of Parkinson's disease: A narrative review. *Annals of Movement Disorders* 6(2), 45-57 (2023).
13. Hayden MR, Leavitt BR, Yasothan U, Kirkpatrick P. Tetrabenazine. *Nat Rev Drug Discov* 8(1), 17-18 (2009).
14. Kaur N, Kumar P, Jamwal S, Deshmukh R, Gauttam V. Tetrabenazine: Spotlight on Drug Review. *Ann Neurosci* 23(3), 176-185 (2016).
15. Sampaio C. Huntington disease - Update on ongoing therapeutic developments and a look toward the future. *Parkinsonism Relat Disord* 122 106049 (2024).
16. Wyant KJ, Ridder AJ, Dayalu P. Huntington's Disease-Update on Treatments. *Curr Neurol Neurosci Rep* 17(4), 33 (2017).
17. Mechoulam R, Fride E, Di Marzo V. Endocannabinoids. *Eur J Pharmacol* 359(1), 1-18 (1998).
18. Di Marzo V, Melck D, Bisogno T, De Petrocellis L. Endocannabinoids: endogenous cannabinoid receptor ligands with neuromodulatory action. *Trends Neurosci* 21(12), 521-528 (1998).
19. Peng J, Fan M, An C, Ni F, Huang W, Luo J. A narrative review of molecular mechanism and therapeutic effect of cannabidiol (CBD). *Basic Clin Pharmacol Toxicol* 130(4), 439-456 (2022).
20. Guindon J, Hohmann AG. Cannabinoid CB2 receptors: a therapeutic target for the treatment of inflammatory and neuropathic pain. *Br J Pharmacol* 153(2), 319-334 (2008).
21. Pertwee RG. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol* 153(2), 199-215 (2008).
22. Wilson RI, Nicoll RA. Endocannabinoid Signaling in the Brain. *Science* 296 678-682 (2002).
23. Aoki J, Isokawa M. Understanding Cellular, Molecular, and Functional Specificity, Heterogeneity, and Diversity of the Endocannabinoid System. *Cells* 13(12), (2024).
24. Basavarajappa BS. Endocannabinoid System and Alcohol Abuse Disorders. In: *RECENT ADVANCES IN CANNABINOID PHYSIOLOGY AND PATHOLOGY*, Bukiya AN (Ed. ^ (Eds). Nature Springer Switzerland p.00. 25p. (2019).
25. Basavarajappa BS, Subbanna S. Molecular Insights into Epigenetics and Cannabinoid Receptors. *Biomolecules* 12(11), (2022).
26. Dallabrida KG, De Oliveira Bender JM, Chade ES, Rodrigues N, Sampaio TB. Endocannabinoid System Changes throughout Life: Implications and Therapeutic Potential for Autism, ADHD, and Alzheimer's Disease. *Brain Sci* 14(6), (2024).
27. Wright NJD. A review of the direct targets of the cannabinoids cannabidiol, Delta9-tetrahydrocannabinol, N-arachidonylethanolamine and 2-arachidonoylglycerol. *AIMS Neurosci* 11(2), 144-165 (2024).
28. Morales P, Hurst DP, Reggio PH. Molecular Targets of the Phytocannabinoids: A Complex Picture. *Prog Chem Org Nat Prod* 103 103-131 (2017).
29. Mechoulam R, Gaoni Y. The absolute configuration of delta-1-tetrahydrocannabinol, the major active constituent of hashish. *Tetrahedron Lett* 12 1109-1111 (1967).

30. Boychuk DG, Goddard G, Mauro G, Orellana MF. The effectiveness of cannabinoids in the management of chronic nonmalignant neuropathic pain: a systematic review. *J Oral Facial Pain Headache* 29(1), 7-14 (2015).
31. Novack GD. Cannabinoids for treatment of glaucoma. *Curr Opin Ophthalmol* 27(2), 146-150 (2016).
32. Phillips RS, Friend AJ, Gibson F *et al.* Antiemetic medication for prevention and treatment of chemotherapy-induced nausea and vomiting in childhood. *Cochrane Database Syst Rev* 2(2), CD007786 (2016).
33. Pomorska DK, Do-Rego JC, Do-Rego JL, Zubrzycka M, Janecka A. Opioid and Cannabinoid System in Food Intake. *Curr Pharm Des* 22(10), 1361-1370 (2016).
34. Mechoulam R, Peters M, Murillo-Rodriguez E, Hanus LO. Cannabidiol--recent advances. *Chem Biodivers* 4(8), 1678-1692 (2007).
35. Borges RS, Batista J, Jr., Viana RB *et al.* Understanding the molecular aspects of tetrahydrocannabinol and cannabidiol as antioxidants. *Molecules* 18(10), 12663-12674 (2013).
36. Hampson AJ, Grimaldi M, Axelrod J, Wink D. Cannabidiol and (-)-Delta9-tetrahydrocannabinol are neuroprotective antioxidants. *Proc Natl Acad Sci U S A* 95(14), 8268-8273 (1998).
37. Yamaori S, Ebisawa J, Okushima Y, Yamamoto I, Watanabe K. Potent inhibition of human cytochrome P450 3A isoforms by cannabidiol: role of phenolic hydroxyl groups in the resorcinol moiety. *Life Sci* 88(15-16), 730-736 (2011).
38. Huizenga MN, Wicker E, Beck VC, Forcelli PA. Anticonvulsant effect of cannabinoid receptor agonists in models of seizures in developing rats. *Epilepsia* 58(9), 1593-1602 (2017).
39. Cristino L, Bisogno T, Di Marzo V. Cannabinoids and the expanded endocannabinoid system in neurological disorders. *Nat Rev Neurol* 16(1), 9-29 (2020).
40. Wallace MJ, Blair RE, Falenski KW, Martin BR, Delorenzo RJ. The endogenous cannabinoid system regulates seizure frequency and duration in a model of temporal lobe epilepsy. *J Pharmacol Exp Ther* 307(1), 129-137 (2003).
41. Alger BE. Retrograde signaling in the regulation of synaptic transmission: focus on endocannabinoids. *Progress in Neurobiology*. 68 247-286 (2002).
42. Schlicker E, Kathmann M. Modulation of transmitter release via presynaptic cannabinoid receptors. *Trends Pharmacol Sci* 22(11), 565-572 (2001).
43. Laprairie RB, Bagher AM, Kelly ME, Denovan-Wright EM. Cannabidiol is a negative allosteric modulator of the cannabinoid CB1 receptor. *Br J Pharmacol* 172(20), 4790-4805 (2015).
44. Russo EB, Burnett A, Hall B, Parker KK. Agonistic properties of cannabidiol at 5-HT1a receptors. *Neurochem Res* 30(8), 1037-1043 (2005).
45. Anand U, Jones B, Korchev Y *et al.* CBD Effects on TRPV1 Signaling Pathways in Cultured DRG Neurons. *J Pain Res* 13 2269-2278 (2020).
46. Shi QX, Yang LK, Shi WL *et al.* The novel cannabinoid receptor GPR55 mediates anxiolytic-like effects in the medial orbital cortex of mice with acute stress. *Mol Brain* 10(1), 38 (2017).
47. Marichal-Cancino BA, Fajardo-Valdez A, Ruiz-Contreras AE, Mendez-Diaz M, Prospero-Garcia O. Advances in the Physiology of GPR55 in the Central Nervous System. *Curr Neuropharmacol* 15(5), 771-778 (2017).
48. Armin S, Muenster S, Abood M, Benamar K. GPR55 in the brain and chronic neuropathic pain. *Behav Brain Res* 406 113248 (2021).
49. Do Val-Da Silva RA, Peixoto-Santos JE, Kandratavicius L *et al.* Protective Effects of Cannabidiol against Seizures and Neuronal Death in a Rat Model of Mesial Temporal Lobe Epilepsy. *Front Pharmacol* 8 131 (2017).
50. Khan AA, Shekh-Ahmad T, Khalil A, Walker MC, Ali AB. Cannabidiol exerts antiepileptic effects by restoring hippocampal interneuron functions in a temporal lobe epilepsy model. *Br J Pharmacol* 175(11), 2097-2115 (2018).
51. Legare CA, Raup-Konsavage WM, Vrana KE. Therapeutic Potential of Cannabis, Cannabidiol, and Cannabinoid-Based Pharmaceuticals. *Pharmacology* 107(3-4), 131-149 (2022).
52. Vrechi TaM, Leao A, Morais IBM *et al.* Cannabidiol induces autophagy via ERK1/2 activation in neural cells. *Sci Rep* 11(1), 5434 (2021).
53. Gugliandolo A, Pollastro F, Bramanti P, Mazzon E. Cannabidiol exerts protective effects in an in vitro model of Parkinson's disease activating AKT/mTOR pathway. *Fitoterapia* 143 104553 (2020).
54. Lujan MA, Valverde O. The Pro-neurogenic Effects of Cannabidiol and Its Potential Therapeutic Implications in Psychiatric Disorders. *Front Behav Neurosci* 14 109 (2020).
55. Lujan MA, Castro-Zavala A, Alegre-Zurano L, Valverde O. Repeated Cannabidiol treatment reduces cocaine intake and modulates neural proliferation and CB1R expression in the mouse hippocampus. *Neuropharmacology* 143 163-175 (2018).
56. Campos AC, Ortega Z, Palazuelos J *et al.* The anxiolytic effect of cannabidiol on chronically stressed mice depends on hippocampal neurogenesis: involvement of the endocannabinoid system. *Int J Neuropsychopharmacol* 16(6), 1407-1419 (2013).

57. Lanza Cariccio V, Scionti D, Raffa A *et al.* Treatment of Periodontal Ligament Stem Cells with MOR and CBD Promotes Cell Survival and Neuronal Differentiation via the PI3K/Akt/mTOR Pathway. *Int J Mol Sci* 19(8), (2018).
58. Giacoppo S, Pollastro F, Grassi G, Bramanti P, Mazzon E. Target regulation of PI3K/Akt/mTOR pathway by cannabidiol in treatment of experimental multiple sclerosis. *Fitoterapia* 116 77-84 (2017).
59. Renard J, Loureiro M, Rosen LG *et al.* Cannabidiol Counteracts Amphetamine-Induced Neuronal and Behavioral Sensitization of the Mesolimbic Dopamine Pathway through a Novel mTOR/p70S6 Kinase Signaling Pathway. *J Neurosci* 36(18), 5160-5169 (2016).
60. Sagredo O, Ramos JA, Decio A, Mechoulam R, Fernandez-Ruiz J. Cannabidiol reduced the striatal atrophy caused 3-nitropropionic acid in vivo by mechanisms independent of the activation of cannabinoid, vanilloid TRPV1 and adenosine A2A receptors. *Eur J Neurosci* 26(4), 843-851 (2007).
61. Jones NA, Hill AJ, Smith I *et al.* Cannabidiol displays antiepileptiform and antiseizure properties in vitro and in vivo. *J Pharmacol Exp Ther* 332(2), 569-577 (2010).
62. Martinez-Aguirre C, Carmona-Cruz F, Velasco AL *et al.* Cannabidiol Acts at 5-HT(1A) Receptors in the Human Brain: Relevance for Treating Temporal Lobe Epilepsy. *Front Behav Neurosci* 14 611278 (2020).
63. Radley JJ, Jacobs BL. Pilocarpine-induced status epilepticus increases cell proliferation in the dentate gyrus of adult rats via a 5-HT1A receptor-dependent mechanism. *Brain Res* 966(1), 1-12 (2003).
64. Schonhoff K, Von Ruden EL, Koska I, Seiffert I, Potschka H. Hippocampal and Septal 5-HT(1A) Receptor Expression in Two Rat Models of Temporal Lobe Epilepsy. *Neuroscience* 465 219-230 (2021).
65. Cai L, Xu Z, Luo H *et al.* The association between 5-HT1A binding and temporal lobe epilepsy: A meta-analysis of molecular imaging studies. *Epilepsy Behav* 145 109354 (2023).
66. Pineda EA, Hensler JG, Sankar R, Shin D, Burke TF, Mazarati AM. Plasticity of presynaptic and postsynaptic serotonin 1A receptors in an animal model of epilepsy-associated depression. *Neuropsychopharmacology* 36(6), 1305-1316 (2011).
67. Lazarini-Lopes W, Campos-Rodriguez C, Garcia-Cairasco N, N'gouemo P, Forcelli PA. Cannabidiol attenuates generalized tonic-clonic and suppresses limbic seizures in the genetically epilepsy-prone rats (GEPR-3) strain. *Pharmacol Rep* 75(1), 166-176 (2023).
68. Barnes JP, Dial H, Owens W *et al.* Adherence and discontinuation of prescription cannabidiol for the management of seizure disorders at an integrated care center. *Epilepsy Res* 200 107300 (2024).
69. Gill EW, Paton WD, Pertwee RG. Preliminary experiments on the chemistry and pharmacology of cannabis. *Nature* 228(5267), 134-136 (1970).
70. Thomas A, Stevenson LA, Wease KN *et al.* Evidence that the plant cannabinoid Delta9-tetrahydrocannabinol is a cannabinoid CB1 and CB2 receptor antagonist. *Br J Pharmacol* 146(7), 917-926 (2005).
71. Franco R, Rivas-Santisteban R, Reyes-Resina I *et al.* Pharmacological potential of varinic-, minor-, and acidic phytocannabinoids. *Pharmacol Res* 158 104801 (2020).
72. Muller C, Morales P, Reggio PH. Cannabinoid Ligands Targeting TRP Channels. *Front Mol Neurosci* 11 487 (2018).
73. Cascio MG, Zamberletti E, Marini P, Parolaro D, Pertwee RG. The phytocannabinoid, Delta(9)-tetrahydrocannabinol, can act through 5-HT(1)A receptors to produce antipsychotic effects. *Br J Pharmacol* 172(5), 1305-1318 (2015).
74. Garcia-Garcia AL, Newman-Tancredi A, Leonardo ED. 5-HT(1A) [corrected] receptors in mood and anxiety: recent insights into autoreceptor versus heteroreceptor function. *Psychopharmacology (Berl)* 231(4), 623-636 (2014).
75. Glikmann-Johnston Y, Saling MM, Reutens DC, Stout JC. Hippocampal 5-HT1A Receptor and Spatial Learning and Memory. *Front Pharmacol* 6 289 (2015).
76. Anavi-Goffer S, Baillie G, Irving AJ *et al.* Modulation of L-alpha-lysophosphatidylinositol/GPR55 mitogen-activated protein kinase (MAPK) signaling by cannabinoids. *J Biol Chem* 287(1), 91-104 (2012).
77. Dennis I, Whalley BJ, Stephens GJ. Effects of Delta9-tetrahydrocannabinol on [35S]GTPgammaS binding in mouse brain cerebellum and piriform cortex membranes. *Br J Pharmacol* 154(6), 1349-1358 (2008).
78. Pertwee RG, Thomas A, Stevenson LA *et al.* The psychoactive plant cannabinoid, Delta9-tetrahydrocannabinol, is antagonized by Delta8- and Delta9-tetrahydrocannabinol in mice in vivo. *Br J Pharmacol* 150(5), 586-594 (2007).
79. Bolognini D, Costa B, Maione S *et al.* The plant cannabinoid Delta9-tetrahydrocannabinol can decrease signs of inflammation and inflammatory pain in mice. *Br J Pharmacol* 160(3), 677-687 (2010).
80. Janssens A, Silvestri C, Martella A, Vanoevelen JM, Di Marzo V, Voets T. Delta(9)-tetrahydrocannabinol impairs epithelial calcium transport through inhibition of TRPV5 and TRPV6. *Pharmacol Res* 136 83-89 (2018).
81. Hill AJ, Weston SE, Jones NA *et al.* Delta(9)-Tetrahydrocannabinol suppresses in vitro epileptiform and in vivo seizure activity in adult rats. *Epilepsia* 51(8), 1522-1532 (2010).

82. Ma YL, Weston SE, Whalley BJ, Stephens GJ. The phytocannabinoid Delta(9)-tetrahydrocannabinol modulates inhibitory neurotransmission in the cerebellum. *Br J Pharmacol* 154(1), 204-215 (2008).
83. Allendorfer JB, Szaflarski JP. Neuroimaging studies towards understanding the central effects of pharmacological cannabis products on patients with epilepsy. *Epilepsy Behav* 70(Pt B), 349-354 (2017).
84. Gaston TE, Friedman D. Pharmacology of cannabinoids in the treatment of epilepsy. *Epilepsy Behav* 70(Pt B), 313-318 (2017).
85. Husni AS, Mccurdy CR, Radwan MM *et al.* Evaluation of Phytocannabinoids from High Potency Cannabis sativa using In Vitro Bioassays to Determine Structure-Activity Relationships for Cannabinoid Receptor 1 and Cannabinoid Receptor 2. *Med Chem Res* 23(9), 4295-4300 (2014).
86. Rosenthaler S, Pohn B, Kolmanz C *et al.* Differences in receptor binding affinity of several phytocannabinoids do not explain their effects on neural cell cultures. *Neurotoxicol Teratol* 46 49-56 (2014).
87. Navarro G, Varani K, Lillo A *et al.* Pharmacological data of cannabidiol- and cannabigerol-type phytocannabinoids acting on cannabinoid CB(1), CB(2) and CB(1)/CB(2) heteromer receptors. *Pharmacol Res* 159 104940 (2020).
88. Zagzoog A, Mohamed KA, Kim HJJ *et al.* In vitro and in vivo pharmacological activity of minor cannabinoids isolated from Cannabis sativa. *Sci Rep* 10(1), 20405 (2020).
89. De Petrocellis L, Ligresti A, Moriello AS *et al.* Effects of cannabinoids and cannabinoid-enriched Cannabis extracts on TRP channels and endocannabinoid metabolic enzymes. *Br J Pharmacol* 163(7), 1479-1494 (2011).
90. De Petrocellis L, Vellani V, Schiano-Moriello A *et al.* Plant-derived cannabinoids modulate the activity of transient receptor potential channels of ankyrin type-1 and melastatin type-8. *J Pharmacol Exp Ther* 325(3), 1007-1015 (2008).
91. Laun AS, Shrader SH, Song ZH. Novel inverse agonists for the orphan G protein-coupled receptor 6. *Heliyon* 4(11), e00933 (2018).
92. Devinsky O, Cilio MR, Cross H *et al.* Cannabidiol: pharmacology and potential therapeutic role in epilepsy and other neuropsychiatric disorders. *Epilepsia* 55(6), 791-802 (2014).
93. Hill AJ, Mercier MS, Hill TD *et al.* Cannabidivarin is anticonvulsant in mouse and rat. *Br J Pharmacol* 167(8), 1629-1642 (2012).
94. Huizenga MN, Sepulveda-Rodriguez A, Forcelli PA. Preclinical safety and efficacy of cannabidivarin for early life seizures. *Neuropharmacology* 148 189-198 (2019).
95. Vigli D, Cosentino L, Raggi C, Laviola G, Woolley-Roberts M, De Filippis B. Chronic treatment with the phytocannabinoid Cannabidivarin (CBDV) rescues behavioural alterations and brain atrophy in a mouse model of Rett syndrome. *Neuropharmacology* 140 121-129 (2018).
96. Zamberletti E, Gabaglio M, Piscitelli F *et al.* Cannabidivarin completely rescues cognitive deficits and delays neurological and motor defects in male Mecp2 mutant mice. *J Psychopharmacol* 33(7), 894-907 (2019).
97. Hill TD, Cascio MG, Romano B *et al.* Cannabidivarin-rich cannabis extracts are anticonvulsant in mouse and rat via a CB1 receptor-independent mechanism. *Br J Pharmacol* 170(3), 679-692 (2013).
98. Amada N, Yamasaki Y, Williams CM, Whalley BJ. Cannabidivarin (CBDV) suppresses pentylenetetrazole (PTZ)-induced increases in epilepsy-related gene expression. *PeerJ* 1 e214 (2013).
99. Brodie MJ, Czapinski P, Pazdera L *et al.* A Phase 2 Randomized Controlled Trial of the Efficacy and Safety of Cannabidivarin as Add-on Therapy in Participants with Inadequately Controlled Focal Seizures. *Cannabis Cannabinoid Res* 6(6), 528-536 (2021).
100. Hurley EN, Ellaway CJ, Johnson AM *et al.* Efficacy and safety of cannabidivarin treatment of epilepsy in girls with Rett syndrome: A phase 1 clinical trial. *Epilepsia* 63(7), 1736-1747 (2022).
101. Navarro G, Varani K, Reyes-Resina I *et al.* Cannabigerol Action at Cannabinoid CB(1) and CB(2) Receptors and at CB(1)-CB(2) Heteroreceptor Complexes. *Front Pharmacol* 9 632 (2018).
102. Nadal X, Del Rio C, Casano S *et al.* Tetrahydrocannabinolic acid is a potent PPARgamma agonist with neuroprotective activity. *Br J Pharmacol* 174(23), 4263-4276 (2017).
103. Cascio MG, Gauson LA, Stevenson LA, Ross RA, Pertwee RG. Evidence that the plant cannabinoid cannabigerol is a highly potent alpha2-adrenoceptor agonist and moderately potent 5HT1A receptor antagonist. *Br J Pharmacol* 159(1), 129-141 (2010).
104. Hill AJ, Jones NA, Smith I *et al.* Voltage-gated sodium (NaV) channel blockade by plant cannabinoids does not confer anticonvulsant effects per se. *Neurosci Lett* 566 269-274 (2014).
105. Anderson LL, Heblinski M, Absalom NL *et al.* Cannabigerolic acid, a major biosynthetic precursor molecule in cannabis, exhibits divergent effects on seizures in mouse models of epilepsy. *Br J Pharmacol* 178(24), 4826-4841 (2021).
106. Ghovanloo MR, Estacion M, Higerd-Rusli GP, Zhao P, Dib-Hajj S, Waxman SG. Inhibition of sodium conductance by cannabigerol contributes to a reduction of dorsal root ganglion neuron excitability. *Br J Pharmacol* 179(15), 4010-4030 (2022).
107. Milligan CJ, Anderson LL, Mcgregor IS, Arnold JC, Petrou S. Beyond CBD: Inhibitory effects of lesser studied phytocannabinoids on human voltage-gated sodium channels. *Front Physiol* 14 1081186 (2023).

108. Iannotti FA, Hill CL, Leo A *et al.* Nonpsychotropic plant cannabinoids, cannabidivarin (CBDV) and cannabidiol (CBD), activate and desensitize transient receptor potential vanilloid 1 (TRPV1) channels in vitro: potential for the treatment of neuronal hyperexcitability. *ACS Chem Neurosci* 5(11), 1131-1141 (2014).
109. Garcia C, Palomo-Garo C, Garcia-Arencibia M, Ramos J, Pertwee R, Fernandez-Ruiz J. Symptom-relieving and neuroprotective effects of the phytocannabinoid Delta(9)-THCV in animal models of Parkinson's disease. *Br J Pharmacol* 163(7), 1495-1506 (2011).
110. Nalls MA, Blauwendraat C, Vallerga CL *et al.* Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 18(12), 1091-1102 (2019).
111. Mack JM, Schamne MG, Sampaio TB *et al.* Melatonergic System in Parkinson's Disease: From Neuroprotection to the Management of Motor and Nonmotor Symptoms. *Oxid Med Cell Longev* 2016 3472032 (2016).
112. Braak H, Del Tredici K. Neuropathological Staging of Brain Pathology in Sporadic Parkinson's disease: Separating the Wheat from the Chaff. *J Parkinsons Dis* 7(s1), S71-S85 (2017).
113. Yasuda T, Nakata Y, Mochizuki H. alpha-Synuclein and neuronal cell death. *Mol Neurobiol* 47(2), 466-483 (2013).
114. Braak H, Ghebremedhin E, Rub U, Bratzke H, Del Tredici K. Stages in the development of Parkinson's disease-related pathology. *Cell Tissue Res* 318(1), 121-134 (2004).
115. Garcia C, Palomo-Garo C, Gomez-Galvez Y, Fernandez-Ruiz J. Cannabinoid-dopamine interactions in the physiology and pathophysiology of the basal ganglia. *Br J Pharmacol* 173(13), 2069-2079 (2016).
116. Oz M, Jaligam V, Galadari S, Petroianu G, Shuba YM, Shippenberg TS. The endogenous cannabinoid, anandamide, inhibits dopamine transporter function by a receptor-independent mechanism. *J Neurochem* 112(6), 1454-1464 (2010).
117. Pisani A, Fezza F, Galati S *et al.* High endogenous cannabinoid levels in the cerebrospinal fluid of untreated Parkinson's disease patients. *Ann Neurol* 57(5), 777-779 (2005).
118. Van Laere K, Casteels C, Lunsken S *et al.* Regional changes in type 1 cannabinoid receptor availability in Parkinson's disease in vivo. *Neurobiol Aging* 33(3), 620 e621-628 (2012).
119. Song L, Yang X, Ma Y, Wu N, Liu Z. The CB1 cannabinoid receptor agonist reduces L-DOPA-induced motor fluctuation and ERK1/2 phosphorylation in 6-OHDA-lesioned rats. *Drug Des Devel Ther* 8 2173-2179 (2014).
120. Gomez-Galvez Y, Palomo-Garo C, Fernandez-Ruiz J, Garcia C. Potential of the cannabinoid CB(2) receptor as a pharmacological target against inflammation in Parkinson's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 64 200-208 (2016).
121. Chagas MH, Zuardi AW, Tumas V *et al.* Effects of cannabidiol in the treatment of patients with Parkinson's disease: an exploratory double-blind trial. *J Psychopharmacol* 28(11), 1088-1098 (2014).
122. De Faria SM, De Moraes Fabricio D, Tumas V *et al.* Effects of acute cannabidiol administration on anxiety and tremors induced by a Simulated Public Speaking Test in patients with Parkinson's disease. *J Psychopharmacol* 34(2), 189-196 (2020).
123. Peball M, Krismer F, Knaus HG *et al.* Non-Motor Symptoms in Parkinson's Disease are Reduced by Nabilone. *Ann Neurol* 88(4), 712-722 (2020).
124. Garcia-Arencibia M, Gonzalez S, De Lago E, Ramos JA, Mechoulam R, Fernandez-Ruiz J. Evaluation of the neuroprotective effect of cannabinoids in a rat model of Parkinson's disease: importance of antioxidant and cannabinoid receptor-independent properties. *Brain Res* 1134(1), 162-170 (2007).
125. Lastres-Becker I, Molina-Holgado F, Ramos JA, Mechoulam R, Fernandez-Ruiz J. Cannabinoids provide neuroprotection against 6-hydroxydopamine toxicity in vivo and in vitro: relevance to Parkinson's disease. *Neurobiol Dis* 19(1-2), 96-107 (2005).
126. Santos NA, Martins NM, Sisti FM *et al.* The neuroprotection of cannabidiol against MPP(+)-induced toxicity in PC12 cells involves trkA receptors, upregulation of axonal and synaptic proteins, neuriteogenesis, and might be relevant to Parkinson's disease. *Toxicol In Vitro* 30(1 Pt B), 231-240 (2015).
127. Wang L, Wu X, Yang G *et al.* Cannabidiol Alleviates the Damage to Dopaminergic Neurons in 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine-Induced Parkinson's Disease Mice Via Regulating Neuronal Apoptosis and Neuroinflammation. *Neuroscience* 498 64-72 (2022).
128. Mattos BA, Bonato JM, Splendor MC, Del Bel E, Milani H, Oliveira RMW. Cannabidiol improves nonmotor symptoms, attenuates neuroinflammation and favors hippocampal newborn neuronal maturation in a rat model of Parkinsonism. *Acta Neuropsychiatr* doi:10.1017/neu.2024.15 1-30 (2024).
129. Espadas I, Keifman E, Palomo-Garo C *et al.* Beneficial effects of the phytocannabinoid Delta(9)-THCV in L-DOPA-induced dyskinesia in Parkinson's disease. *Neurobiol Dis* 141 104892 (2020).
130. De Mattos BA, Bonato JM, Splendor MC, Del Bel E, Milani H, De Oliveira RMW. Cannabidiol improves non-motor symptoms, attenuates neuroinflammation, and favours hippocampal newborn neuronal maturation in a rat model of Parkinsonism. *Acta Neuropsychiatr* doi:10.1017/neu.2024.15 1-13 (2024).

131. Burgaz S, Garcia C, Gomez-Canas M, Munoz E, Fernandez-Ruiz J. Development of An Oral Treatment with the PPAR-gamma-Acting Cannabinoid VCE-003.2 Against the Inflammation-Driven Neuronal Deterioration in Experimental Parkinson's Disease. *Molecules* 24(15), (2019).
132. Burgaz S, Garcia C, Gomez-Canas M *et al.* Neuroprotection with the cannabigerol quinone derivative VCE-003.2 and its analogs CBGA-Q and CBGA-Q-Salt in Parkinson's disease using 6-hydroxydopamine-lesioned mice. *Mol Cell Neurosci* 110 103583 (2021).
133. Burgaz S, Garcia C, Gomez-Canas M, Rolland A, Munoz E, Fernandez-Ruiz J. Neuroprotection with the Cannabidiol Quinone Derivative VCE-004.8 (EHP-101) against 6-Hydroxydopamine in Cell and Murine Models of Parkinson's Disease. *Molecules* 26(11), (2021).
134. Wang F, Jin T, Li H *et al.* Cannabidiol alleviates alpha-synuclein aggregation via DAF-16 in *Caenorhabditis elegans*. *FASEB J* 37(2), e22735 (2023).
135. Harrell ER, King JW, Stoeckel LE, Trevino M. National Institute on Aging's 50th anniversary: Advancing cognitive aging research and the cognitive health of older adults. *J Gerontol B Psychol Sci Soc Sci* doi:10.1093/geronb/gbae120 (2024).
136. Ramirez BG, Blazquez C, Gomez Del Pulgar T, Guzman M, De Ceballos ML. Prevention of Alzheimer's disease pathology by cannabinoids: neuroprotection mediated by blockade of microglial activation. *J Neurosci* 25(8), 1904-1913 (2005).
137. Solas M, Francis PT, Franco R, Ramirez MJ. CB2 receptor and amyloid pathology in frontal cortex of Alzheimer's disease patients. *Neurobiol Aging* 34(3), 805-808 (2013).
138. Li S, Huang Y, Yu L, Ji X, Wu J. Impact of the Cannabinoid System in Alzheimer's Disease. *Curr Neuropharmacol* 21(3), 715-726 (2023).
139. Khavandi M, Rao PPN, Beazely MA. Differential Effects of Endocannabinoids on Amyloid-Beta Aggregation and Toxicity. *Int J Mol Sci* 24(2), (2023).
140. Rapaka D, Adiukwu PC, Challa SR, Bitra VR. Interplay Between Astroglial Endocannabinoid System and the Cognitive Dysfunction in Alzheimer's Disease. *Physiol Res* 72(5), 575-586 (2023).
141. Iuvone T, Esposito G, Esposito R, Santamaria R, Di Rosa M, Izzo AA. Neuroprotective effect of cannabidiol, a non-psychoactive component from Cannabis sativa, on beta-amyloid-induced toxicity in PC12 cells. *J Neurochem* 89(1), 134-141 (2004).
142. Vallee A, Lecarpentier Y, Guillevin R, Vallee JN. Effects of cannabidiol interactions with Wnt/beta-catenin pathway and PPARgamma on oxidative stress and neuroinflammation in Alzheimer's disease. *Acta Biochim Biophys Sin (Shanghai)* 49(10), 853-866 (2017).
143. Zhang N, Parr CJC, Birch AM, Goldfinger MH, Sastre M. The amyloid precursor protein binds to beta-catenin and modulates its cellular distribution. *Neurosci Lett* 685 190-195 (2018).
144. Libro R, Diomedea F, Scionti D *et al.* Cannabidiol Modulates the Expression of Alzheimer's Disease-Related Genes in Mesenchymal Stem Cells. *Int J Mol Sci* 18(1), (2016).
145. Li H, Liu Y, Tian D *et al.* Overview of cannabidiol (CBD) and its analogues: Structures, biological activities, and neuroprotective mechanisms in epilepsy and Alzheimer's disease. *Eur J Med Chem* 192 112163 (2020).
146. Martin-Moreno AM, Reigada D, Ramirez BG *et al.* Cannabidiol and other cannabinoids reduce microglial activation in vitro and in vivo: relevance to Alzheimer's disease. *Mol Pharmacol* 79(6), 964-973 (2011).
147. Cheng D, Low JK, Logge W, Garner B, Karl T. Chronic cannabidiol treatment improves social and object recognition in double transgenic APP^{swe}/PS1 Δ E9 mice. *Psychopharmacology (Berl)* 231(15), 3009-3017 (2014).
148. Cheng D, Spiro AS, Jenner AM, Garner B, Karl T. Long-term cannabidiol treatment prevents the development of social recognition memory deficits in Alzheimer's disease transgenic mice. *J Alzheimers Dis* 42(4), 1383-1396 (2014).
149. Arnanz MA, Ruiz De Martin Esteban S, Martinez Relimpio AM *et al.* Effects of Chronic, Low-Dose Cannabinoids, Cannabidiol, Delta-9-Tetrahydrocannabinol and a Combination of Both, on Amyloid Pathology in the 5xFAD Mouse Model of Alzheimer's Disease. *Cannabis Cannabinoid Res* doi:10.1089/can.2023.0101 (2023).
150. Singh J, Neary JP. Neuroprotection Following Concussion: The Potential Role for Cannabidiol. *Can J Neurol Sci* 47(3), 289-300 (2020).
151. Ma BQ, Jia JX, Wang H *et al.* Cannabidiol improves the cognitive function of SAMP8 AD model mice involving the microbiota-gut-brain axis. *J Toxicol Environ Health A* 87(11), 471-479 (2024).
152. Velayudhan L, Dugonjic M, Pisani S *et al.* Cannabidiol for behavior symptoms in Alzheimer's disease (CANBiS-AD): a randomized, double-blind, placebo-controlled trial. *Int Psychogeriatr* doi:10.1017/S1041610224000516 1-3 (2024).
153. Amini M, Abdolmaleki Z. The Effect of Cannabidiol Coated by Nano-Chitosan on Learning and Memory, Hippocampal CB1 and CB2 Levels, and Amyloid Plaques in an Alzheimer's Disease Rat Model. *Neuropsychobiology* 81(3), 171-183 (2022).
154. Novotna A, Mares J, Ratcliffe S *et al.* A randomized, double-blind, placebo-controlled, parallel-group, enriched-design study of nabiximols* (Sativex((R))), as add-on therapy, in subjects with refractory spasticity caused by multiple sclerosis. *Eur J Neurol* 18(9), 1122-1131 (2011).

155. Broers B, Pata Z, Mina A, Wampfler J, De Saussure C, Pautex S. Prescription of a THC/CBD-Based Medication to Patients with Dementia: A Pilot Study in Geneva. *Med Cannabis Cannabinoids* 2(1), 56-59 (2019).
156. Schubert D, Kepchia D, Liang Z, Dargusch R, Goldberg J, Maher P. Efficacy of Cannabinoids in a Pre-Clinical Drug-Screening Platform for Alzheimer's Disease. *Mol Neurobiol* 56(11), 7719-7730 (2019).
157. Marsh DT, Sugiyama A, Imai Y, Kato R, Smid SD. The structurally diverse phytocannabinoids cannabichromene, cannabigerol and cannabinal significantly inhibit amyloid beta-evoked neurotoxicity and changes in cell morphology in PC12 cells. *Basic Clin Pharmacol Toxicol* 134(3), 293-309 (2024).
158. Englund A, Atakan Z, Kralj A, Tunstall N, Murray R, Morrison P. The effect of five day dosing with THC/V on THC-induced cognitive, psychological and physiological effects in healthy male human volunteers: A placebo-controlled, double-blind, crossover pilot trial. *J Psychopharmacol* 30(2), 140-151 (2016).
159. Puopolo T, Liu C, Ma H, Seeram NP. Inhibitory Effects of Cannabinoids on Acetylcholinesterase and Butyrylcholinesterase Enzyme Activities. *Med Cannabis Cannabinoids* 5(1), 85-94 (2022).
160. Melnikova I. Therapies for Alzheimer's disease. *Nat Rev Drug Discov* 6(5), 341-342 (2007).
161. Mccolgan P, Tabrizi SJ. Huntington's disease: a clinical review. *Eur J Neurol* 25(1), 24-34 (2018).
162. Denovan-Wright EM, Robertson HA. Cannabinoid receptor messenger RNA levels decrease in a subset of neurons of the lateral striatum, cortex and hippocampus of transgenic Huntington's disease mice. *Neuroscience* 98(4), 705-713 (2000).
163. Glass M, Dragunow M, Faull RL. The pattern of neurodegeneration in Huntington's disease: a comparative study of cannabinoid, dopamine, adenosine and GABA(A) receptor alterations in the human basal ganglia in Huntington's disease. *Neuroscience*. 97 505-519 (2000).
164. Van Laere K, Casteels C, Dhollander I *et al.* Widespread decrease of type 1 cannabinoid receptor availability in Huntington disease in vivo. *J Nucl Med* 51(9), 1413-1417 (2010).
165. Laprairie RB, Kelly ME, Denovan-Wright EM. Cannabinoids increase type 1 cannabinoid receptor expression in a cell culture model of striatal neurons: implications for Huntington's disease. *Neuropharmacology* 72 47-57 (2013).
166. Mccaw EA, Hu H, Gomez GT, Hebb AL, Kelly ME, Denovan-Wright EM. Structure, expression and regulation of the cannabinoid receptor gene (CB1) in Huntington's disease transgenic mice. *Eur J Biochem* 271(23-24), 4909-4920 (2004).
167. Blazquez C, Chiarlone A, Sagredo O *et al.* Loss of striatal type 1 cannabinoid receptors is a key pathogenic factor in Huntington's disease. *Brain* 134(Pt 1), 119-136 (2011).
168. Chiarlone A, Bellocchio L, Blazquez C *et al.* A restricted population of CB1 cannabinoid receptors with neuroprotective activity. *Proc Natl Acad Sci U S A* 111(22), 8257-8262 (2014).
169. Consroe P, Laguna J, Allender J *et al.* Controlled clinical trial of cannabidiol in Huntington's disease. *Pharmacol Biochem Behav* 40(3), 701-708 (1991).
170. Valdeolivas S, Satta V, Pertwee RG, Fernandez-Ruiz J, Sagredo O. Sativex-like combination of phytocannabinoids is neuroprotective in malonate-lesioned rats, an inflammatory model of Huntington's disease: role of CB1 and CB2 receptors. *ACS Chem Neurosci* 3(5), 400-406 (2012).
171. Sagredo O, Pazos MR, Satta V, Ramos JA, Pertwee RG, Fernandez-Ruiz J. Neuroprotective effects of phytocannabinoid-based medicines in experimental models of Huntington's disease. *J Neurosci Res* 89(9), 1509-1518 (2011).
172. Ruhaak LR, Felth J, Karlsson PC, Rafter JJ, Verpoorte R, Bohlin L. Evaluation of the cyclooxygenase inhibiting effects of six major cannabinoids isolated from Cannabis sativa. *Biol Pharm Bull* 34(5), 774-778 (2011).
173. Valdeolivas S, Navarrete C, Cantarero I, Bellido ML, Munoz E, Sagredo O. Neuroprotective properties of cannabigerol in Huntington's disease: studies in R6/2 mice and 3-nitropropionate-lesioned mice. *Neurotherapeutics* 12(1), 185-199 (2015).
174. Gugliandolo A, Silvestro S, Chiricosta L, Pollastro F, Bramanti P, Mazzon E. The Transcriptomic Analysis of NSC-34 Motor Neuron-Like Cells Reveals That Cannabigerol Influences Synaptic Pathways: A Comparative Study with Cannabidiol. *Life (Basel)* 10(10), (2020).
175. De Lago E, Urbani P, Ramos JA, Di Marzo V, Fernandez-Ruiz J. Arvanil, a hybrid endocannabinoid and vanilloid compound, behaves as an antihyperkinetic agent in a rat model of Huntington's disease. *Brain Res* 1050(1-2), 210-216 (2005).
176. Lastres-Becker I, Bizat N, Boyer F, Hantraye P, Brouillet E, Fernandez-Ruiz J. Effects of cannabinoids in the rat model of Huntington's disease generated by an intrastriatal injection of malonate. *Neuroreport* 14(6), 813-816 (2003).
177. De Lago E, Fernandez-Ruiz J, Ortega-Gutierrez S *et al.* UCM707, an inhibitor of the anandamide uptake, behaves as a symptom control agent in models of Huntington's disease and multiple sclerosis, but fails to delay/arrest the progression of different motor-related disorders. *Eur Neuropsychopharmacol* 16(1), 7-18 (2006).

178. Lastres-Becker I, Fezza F, Cebeira M *et al.* Changes in endocannabinoid transmission in the basal ganglia in a rat model of Huntington's disease. *Neuroreport* 12(10), 2125-2129 (2001).
179. Borrelli F, Fasolino I, Romano B *et al.* Beneficial effect of the non-psychotropic plant cannabinoid cannabigerol on experimental inflammatory bowel disease. *Biochem Pharmacol* 85(9), 1306-1316 (2013).
180. Nachnani R, Raup-Konsavage WM, Vrana KE. The Pharmacological Case for Cannabigerol. *J Pharmacol Exp Ther* 376(2), 204-212 (2021).
181. Starkus J, Jansen C, Shimoda LMN, Stokes AJ, Small-Howard AL, Turner H. Diverse TRPV1 responses to cannabinoids. *Channels (Austin)* 13(1), 172-191 (2019).
182. De Almeida DL, Devi LA. Diversity of molecular targets and signaling pathways for CBD. *Pharmacol Res Perspect* 8(6), e00682 (2020).
183. Di Marzo V, Piscitelli F. The Endocannabinoid System and its Modulation by Phytocannabinoids. *Neurotherapeutics* 12(4), 692-698 (2015).
184. Ramer R, Heinemann K, Merkord J *et al.* COX-2 and PPAR-gamma confer cannabidiol-induced apoptosis of human lung cancer cells. *Mol Cancer Ther* 12(1), 69-82 (2013).
185. Deiana S, Watanabe A, Yamasaki Y *et al.* Plasma and brain pharmacokinetic profile of cannabidiol (CBD), cannabidivarin (CBDV), Delta(9)-tetrahydrocannabivarin (THCV) and cannabigerol (CBG) in rats and mice following oral and intraperitoneal administration and CBD action on obsessive-compulsive behaviour. *Psychopharmacology (Berl)* 219(3), 859-873 (2012).
186. Glass M, Dragunow M, Faull RL. Cannabinoid receptors in the human brain: a detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience* 77(2), 299-318 (1997).
187. Basavarajappa BS, Joshi V, Shivakumar M, Subbanna S. Distinct Functions of Endogenous Cannabinoid System in Alcohol Abuse Disorders. *Br J Pharmacol* doi:10.1111/bph.14780 (2019).
188. Parsons LH, Hurd YL. Endocannabinoid signalling in reward and addiction. *Nat Rev Neurosci* 16(10), 579-594 (2015).
189. Stern CAJ, De Carvalho CR, Bertoglio LJ, Takahashi RN. Effects of Cannabinoid Drugs on Aversive or Rewarding Drug-Associated Memory Extinction and Reconsolidation. *Neuroscience* 370 62-80 (2018).
190. Sabia S, Fayosse A, Dumurgier J *et al.* Alcohol consumption and risk of dementia: 23 year follow-up of Whitehall II cohort study. *BMJ* 362 k2927 (2018).
191. Stampfer MJ, Kang JH, Chen J, Cherry R, Grodstein F. Effects of moderate alcohol consumption on cognitive function in women. *N Engl J Med* 352(3), 245-253 (2005).
192. Liu SJ, Fang ZY, Yang Y, Deng HM, Wang JZ. Alzheimer-like phosphorylation of tau and neurofilament induced by cocaine in vivo. *Acta Pharmacol Sin* 24(6), 512-518 (2003).
193. Shukla M, Vincent B. The multi-faceted impact of methamphetamine on Alzheimer's disease: From a triggering role to a possible therapeutic use. *Ageing Res Rev* 60 101062 (2020).
194. Morgan CJ, Das RK, Joye A, Curran HV, Kamboj SK. Cannabidiol reduces cigarette consumption in tobacco smokers: preliminary findings. *Addict Behav* 38(9), 2433-2436 (2013).
195. Hindocha C, Freeman TP, Grabski M *et al.* Cannabidiol reverses attentional bias to cigarette cues in a human experimental model of tobacco withdrawal. *Addiction* 113(9), 1696-1705 (2018).
196. Hindocha C, Freeman TP, Grabski M *et al.* The effects of cannabidiol on impulsivity and memory during abstinence in cigarette dependent smokers. *Sci Rep* 8(1), 7568 (2018).
197. Gournay LR, Petry J, Bilsky S *et al.* Cannabidiol Reduces Nicotine Withdrawal Severity and State Anxiety During an Acute E-cigarette Abstinence Period: A Novel, Open-Label Study. *Cannabis Cannabinoid Res* doi:10.1089/can.2022.0317 (2023).
198. Cheeks SN, Buzzi B, Valdez A, Mogul AS, Damaj MI, Fowler CD. Cannabidiol as a potential cessation therapeutic: Effects on intravenous nicotine self-administration and withdrawal symptoms in mice. *Neuropharmacology* 246 109833 (2024).
199. Smith LC, Tieu L, Suhandynata RT *et al.* Cannabidiol reduces withdrawal symptoms in nicotine-dependent rats. *Psychopharmacology (Berl)* 238(8), 2201-2211 (2021).
200. Black N, Stockings E, Campbell G *et al.* Cannabinoids for the treatment of mental disorders and symptoms of mental disorders: a systematic review and meta-analysis. *Lancet Psychiatry* 6(12), 995-1010 (2019).
201. Hindocha C, Freeman TP, Schafer G *et al.* Acute effects of cannabinoids on addiction endophenotypes are moderated by genes encoding the CB1 receptor and FAAH enzyme. *Addict Biol* 25(3), e12762 (2020).
202. Broyd SJ, Van Hell HH, Beale C, Yucel M, Solowij N. Acute and Chronic Effects of Cannabinoids on Human Cognition-A Systematic Review. *Biol Psychiatry* 79(7), 557-567 (2016).
203. Freeman AM, Petrilli K, Lees R *et al.* How does cannabidiol (CBD) influence the acute effects of delta-9-tetrahydrocannabinol (THC) in humans? A systematic review. *Neurosci Biobehav Rev* 107 696-712 (2019).
204. Pokorski I, Clement N, Phung N, Weltman M, Fu S, J. C. Cannabidiol in the management of inpatient cannabis withdrawal: clinical case series. *Future Neurol.* 12(3), 133-140. (2017).
205. Crippa JA, Hallak JE, Machado-De-Sousa JP *et al.* Cannabidiol for the treatment of cannabis withdrawal syndrome: a case report. *J Clin Pharm Ther* 38(2), 162-164 (2013).

206. Shannon S, Opila-Lehman J. Cannabidiol Oil for Decreasing Addictive Use of Marijuana: A Case Report. *Integr Med (Encinitas)* 14(6), 31-35 (2015).
207. Cleirec G, Desmier E, Lacatus C *et al.* Efficiency of Inhaled Cannabidiol in Cannabis Use Disorder: The Pilot Study Cannavap. *Front Psychiatry* 13 899221 (2022).
208. Solowij N, Broyd SJ, Beale C *et al.* Therapeutic Effects of Prolonged Cannabidiol Treatment on Psychological Symptoms and Cognitive Function in Regular Cannabis Users: A Pragmatic Open-Label Clinical Trial. *Cannabis Cannabinoid Res* 3(1), 21-34 (2018).
209. Lorenzetti V, Mctavish E, Broyd S *et al.* Daily Cannabidiol Administration for 10 Weeks Modulates Hippocampal and Amygdalar Resting-State Functional Connectivity in Cannabis Users: A Functional Magnetic Resonance Imaging Open-Label Clinical Trial. *Cannabis Cannabinoid Res* doi:10.1089/can.2022.0336 (2023).
210. De Carvalho CR, Takahashi RN. Cannabidiol disrupts the reconsolidation of contextual drug-associated memories in Wistar rats. *Addict Biol* 22(3), 742-751 (2017).
211. Ren Y, Whittard J, Higuera-Matas A, Morris CV, Hurd YL. Cannabidiol, a nonpsychotropic component of cannabis, inhibits cue-induced heroin seeking and normalizes discrete mesolimbic neuronal disturbances. *J Neurosci* 29(47), 14764-14769 (2009).
212. Navarrete F, Gasparyan A, Manzanares J. CBD-mediated regulation of heroin withdrawal-induced behavioural and molecular changes in mice. *Addict Biol* 27(2), e13150 (2022).
213. Scicluna RL, Wilson BB, Thelaus SH, Arnold JC, Mcgregor IS, Bowen MT. Cannabidiol Reduced the Severity of Gastrointestinal Symptoms of Opioid Withdrawal in Male and Female Mice. *Cannabis Cannabinoid Res* 9(2), 547-560 (2024).
214. Hayduk SA, Hughes AC, Winter RL, Milton MD, Ward SJ. Single and Combined Effects of Cannabigerol (CBG) and Cannabidiol (CBD) in Mouse Models of Oxaliplatin-Associated Mechanical Sensitivity, Opioid Antinociception, and Naloxone-Precipitated Opioid Withdrawal. *Biomedicines* 12(6), (2024).
215. Socha J, Grochecki P, Marszalek-Grabska M *et al.* Cannabidiol Protects against the Reinstatement of Oxycodone-Induced Conditioned Place Preference in Adolescent Male but Not Female Rats: The Role of MOR and CB1R. *Int J Mol Sci* 25(12), (2024).
216. Li H, Ward SJ. Paclitaxel-Associated Mechanical Sensitivity and Neuroinflammation Are Sex-, Time-, and Site-Specific and Prevented through Cannabigerol Administration in C57Bl/6 Mice. *Int J Mol Sci* 25(8), (2024).
217. Lorenzetti V, Mctavish E, Broyd S *et al.* Daily Cannabidiol Administration for 10 Weeks Modulates Hippocampal and Amygdalar Resting-State Functional Connectivity in Cannabis Users: A Functional Magnetic Resonance Imaging Open-Label Clinical Trial. *Cannabis Cannabinoid Res* 9(4), e1108-e1121 (2024).
218. Guimaraes FS, Chiaretti TM, Graeff FG, Zuardi AW. Antianxiety effect of cannabidiol in the elevated plus-maze. *Psychopharmacology (Berl)* 100(4), 558-559 (1990).
219. Almeida V, Levin R, Peres FF *et al.* Cannabidiol exhibits anxiolytic but not antipsychotic property evaluated in the social interaction test. *Prog Neuropsychopharmacol Biol Psychiatry* 41 30-35 (2013).
220. Arnone M, Maruani J, Chaperon F *et al.* Selective inhibition of sucrose and ethanol intake by SR 141716, an antagonist of central cannabinoid (CB1) receptors. *Psychopharmacology*. 132 104-106 (1997).
221. Basavarajappa BS. Endocannabinoid system in the development of tolerance to alcohol. . *Klinik and Forschung (Journal of Clinical Research)*. 11 16-19 (2005).
222. Colombo G, Agabio R, Fa M *et al.* Reduction of voluntary ethanol intake in ethanol-preferring sP rats by the cannabinoid antagonist SR-141716. *Alcohol and alcoholism* 33 126-130 (1998).
223. Hungund BL, Szakall I, Adam A, Basavarajappa BS, Vadasz C. Cannabinoid CB1 receptor knockout mice exhibit markedly reduced voluntary alcohol consumption and lack alcohol-induced dopamine release in the nucleus accumbens. *J Neurochem* 84(4), 698-704 (2003).
224. Hamelink C, Hampson A, Wink DA, Eiden LE, Eskay RL. Comparison of cannabidiol, antioxidants, and diuretics in reversing binge ethanol-induced neurotoxicity. *J Pharmacol Exp Ther* 314(2), 780-788 (2005).
225. Liput DJ, Hammell DC, Stinchcomb AL, Nixon K. Transdermal delivery of cannabidiol attenuates binge alcohol-induced neurodegeneration in a rodent model of an alcohol use disorder. *Pharmacol Biochem Behav* 111 120-127 (2013).
226. Brenneman DE, Petkanas D, Kinney WA. Pharmacological Comparisons Between Cannabidiol and KLS-13019. *J Mol Neurosci* 66(1), 121-134 (2018).
227. Oberlin BG, Grahame NJ. High-alcohol preferring mice are more impulsive than low-alcohol preferring mice as measured in the delay discounting task. *Alcohol Clin Exp Res* 33(7), 1294-1303 (2009).
228. Wilhelm CJ, Mitchell SH. Strain differences in delay discounting using inbred rats. *Genes Brain Behav* 8(4), 426-434 (2009).
229. Amlung M, Vedelago L, Acker J, Balodis I, Mackillop J. Steep delay discounting and addictive behavior: a meta-analysis of continuous associations. *Addiction* 112(1), 51-62 (2017).
230. Mackillop J, Amlung MT, Few LR, Ray LA, Sweet LH, Munafo MR. Delayed reward discounting and addictive behavior: a meta-analysis. *Psychopharmacology (Berl)* 216(3), 305-321 (2011).

231. Viudez-Martinez A, Garcia-Gutierrez MS, Navarron CM *et al.* Cannabidiol reduces ethanol consumption, motivation and relapse in mice. *Addict Biol* 23(1), 154-164 (2018).
232. Karoly HC, Drennan ML, Prince MA, Zulic L, Dooley G. Consuming oral cannabidiol prior to a standard alcohol dose has minimal effect on breath alcohol level and subjective effects of alcohol. *Psychopharmacology (Berl)* 240(5), 1119-1129 (2023).
233. Maccioni P, Bratzu J, Carai MaM, Colombo G, Gessa GL. Reducing Effect of Cannabidiol on Alcohol Self-Administration in Sardinian Alcohol-Preferring Rats. *Cannabis Cannabinoid Res* 7(2), 161-169 (2022).
234. Gasparyan A, Navarrete F, Navarro D, Manzanares J. Cannabidiol regulates behavioral and brain alterations induced by spontaneous alcohol withdrawal. *Neuropharmacology* 233 109549 (2023).
235. Tringali G, Lavanco G, Castelli V *et al.* Cannabidiol tempers alcohol intake and neuroendocrine and behavioural correlates in alcohol binge drinking adolescent rats. Focus on calcitonin gene-related peptide's brain levels. *Phytother Res* 37(11), 4870-4884 (2023).
236. Brancato A, Castelli V, Lavanco G *et al.* Social stress under binge-like alcohol withdrawal in adolescence: evidence of cannabidiol effect on maladaptive plasticity in rats. *Psychol Med* 53(12), 5538-5550 (2023).
237. Szulc M, Kujawski R, Pacholak A *et al.* Cannabidiol as a Modulator of the Development of Alcohol Tolerance in Rats. *Nutrients* 15(7), (2023).
238. Melkumyan M, Annaswamy VM, Evans AM *et al.* Effects of cannabidiol, with and without Δ^9 -tetrahydrocannabinol, on anxiety-like behavior following alcohol withdrawal in mice. *Front Neurosci* 18 1375440 (2024).
239. Filev R, Engelke DS, Da Silveira DX, Mello LE, Santos-Junior JG. THC inhibits the expression of ethanol-induced locomotor sensitization in mice. *Alcohol* 65 31-35 (2017).
240. Moore CF, Zamarripa CA, Weerts EM. Oral Cannabidiol does not alter Alcohol Seeking and Self-Administration in Baboons. *Drug Alcohol Depend* 245 109829 (2023).
241. Garcia-Baos A, Puig-Reyne X, Garcia-Algar O, Valverde O. Cannabidiol attenuates cognitive deficits and neuroinflammation induced by early alcohol exposure in a mice model. *Biomed Pharmacother* 141 111813 (2021).
242. Koren G, Cohen R, Sachs O. Use of Cannabis in Fetal Alcohol Spectrum Disorder. *Cannabis Cannabinoid Res* 6(1), 74-76 (2021).
243. Gasparyan A, Navarro D, Navarrete F *et al.* Cannabidiol repairs behavioral and brain disturbances in a model of fetal alcohol spectrum disorder. *Pharmacol Res* 188 106655 (2023).
244. Arnone M, Maruani J, Chaperon F *et al.* Selective inhibition of sucrose and ethanol intake by SR 141716, an antagonist of central cannabinoid (CB1) receptors. *Psychopharmacology (Berl)* 132(1), 104-106 (1997).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.