

Article

Not peer-reviewed version

Pathological Study of Demyelination with Cellular Reactions in the Cerebellum of Dogs Infected with Canine Distemper Virus

[José Manuel Verdes](#)*, Camila Larrañaga, [Guillermo Godiño](#), Belén Varela, Victoria Yozzi, [Victoria Iribarnegaray](#), Luis Delucchi, Kanji Yamasaki

Posted Date: 9 October 2024

doi: 10.20944/preprints202410.0627.v1

Keywords: canine distemper; demyelination; glial cells; pathology; Purkinje cells



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Pathological Study of Demyelination with Cellular Reactions in the Cerebellum of Dogs Infected with Canine Distemper Virus

José Manuel Verdes ^{1,*}, Camila Larrañaga ¹, Guillermo Godíño ¹, Belén Varela ¹, Victoria Yozzi ¹, Victoria Iribarnegaray ², Luis Delucchi ³ and Kanji Yamasaki ^{1,4}

¹ Department of Pathobiology. Pathology Unit. Faculty of Veterinary, Universidad de la República (Udelar). Route 8 Km 18, 13000, Montevideo, Uruguay; patologia@fvet.edu.uy

² Department of Pathobiology. Microbiology Unit. Faculty of Veterinary, Udelar

³ Department of Clinics & Veterinary Hospital. Small Animals Medicine Unit, Clinical Neurology, Faculty of Veterinary, Udelar

⁴ Japan International Cooperation Agency (JICA). Br. Artigas 417 Of.601, 11300, Montevideo, Uruguay

* Correspondence: jose.verdes@fvet.edu.uy; Tel.: +(598) 2 1903 2519.

Abstract: The purpose of this study was to examine the relationship between demyelination and cellular reactions in the cerebellum of Canine Distemper Virus (CDV)-infected dogs. We subdivided the disease staging by adding the degree of demyelination determined by Luxol Fast Blue staining to the previously reported disease staging from the acute stage to the chronic stage, and investigated the relationship between demyelination in cerebellum and the number and histological changes of astroglia, microglia and Purkinje cells in each stage. Reactions of astrocytes and microglia were observed at an early stage when demyelination was not evident. Changes progressed with demyelination. Demyelination initially began in the medulla adjoining the fourth ventricle and gradually spread to the entire cerebellum, including lobes. CDV immune-positive granules were seen from the early stage, and inclusion bodies also appeared at the same time. CDV immune-positive reaction and inclusion bodies were observed in astrocytes, microglia, neurons, ependymal cells, and even leptomeningeal mononuclear cells. On the other hand, infiltration of CDV immunoreactive particles from the pia mater to the gray matter and further into the white matter through the granular layer was observed from an early stage. Purkinje cells decreased from the intermediate stage, and a decrease in cells in the granular layer was also observed. There was no clear association between age and each stage, and the stages did not progress with age.

Keywords: canine distemper; demyelination; glial cells; pathology; Purkinje cells

1. Introduction

Canine distemper is an infectious disease caused by a Morbillivirus, belonging to Paramyxoviridae family, which infect a broad range of terrestrial and aquatic carnivores [1]. It has been reported that demyelination is a characteristic change in the brain of dogs infected with Canine Distemper Virus (CDV), and it has been discussed that glial cells such as astrocytes, oligodendroglia, and microglia are related with demyelination [1–5]. Many studies have focused on changes in astrocytes during the early stages of CDV infection [6–8], and Kabakci et al. [9] reported that CDV infected both astrocytes and oligodendrocytes, and the gradual loss of oligodendrocytes was most likely responsible for the progressive demyelination. In addition, demyelination during CDV-demyelinating lesions was said to be appear representing a biphasic process with a primary virus-induced oligodendroglial dystrophy followed by a secondary wave of immune-mediated myelin destruction [10,11]. Two possible routes for demyelination have been reported that one route is that CDV infects astrocytes, which then infects oligodendroglia, and the other route is that CDV infects microglia through astrocytes and ultimately affects oligodendroglia [3]. It has been well known that

CDV primarily affects the cerebellum [5,12,13]. On the other hand, studies using gene expression analysis of astrocytes after infection have also been reported recently, as astrocytes are the main cellular target of CDV and already undergo reactive changes in brain lesions before demyelination [2].

Histopathologically, cerebellar lesions have been classified based on Hematoxylin-Eosin (HE) stained specimens, with the addition of immunohistochemistry, transcriptome, and various other methods [3,8,10,14–16]. On the other hand, it has been reported that CDV infection experiments on ferret preferentially target neurons in general, and particularly Purkinje cells and granule cells [17]. The purpose of this research is to clarify the relationship between demyelination and glial cells, using the cerebellum of natural CDV-infected dogs, in which the presence of virus was confirmed by immunohistochemistry against CDV, and positive Polymerase Chain Reaction (PCR) tests. Furthermore, we subdivided the stage classification by adding the degree of demyelination using Luxol Fast Blue (LFB) staining to the conventional classification from acute- to chronic- stage, and examined the relationship between demyelination and each glial cell in each stage. Finally, we investigated the Purkinje cells changes in each of these stages.

2. Materials and Methods

2.1. Dogs

Twenty-eight dogs naturally infected by CDV were used in this study. The ages ranged from 25 days to 10 years. There were 15 male and 13 female dogs, respectively. CDV immunopositive granules were confirmed in several regions of the brain of the dead dogs through immunohistochemical testing, and infection in twenty-eight dogs was confirmed through PCR analysis. In addition, considering the age of CDV-infected dogs, five dogs ranging from 45 days old to 6 years old were selected as a control group. In the control group, infectious diseases were excluded as the cause of death, and immunohistochemical or PCR tests were negative.

In our previous study in natural CDV infected dogs [18], we classified the stages of the histopathological lesions following as acute, subacute, or chronic based on previous studies in natural and experimental infection in dogs [3,11,14–16,19,20]. Namely, acute lesions were characteristic by focal vacuolation, gliosis, inclusion bodies and CDV-immunopositive cells without demyelination, subacute lesions were demyelination, gliosis, necrosis, inclusion bodies and CDV-immunopositive cells, and perivascular mononuclear infiltration (cuffing) of two to three layers of thickness, and chronic lesions were similar to the subacute stage, but perivascular infiltration was more prominent, of at least three layers of thickness. In the present study, we subdivided subacute changes based on the degree of demyelination determined by LFB staining in order to clarify the relationship between demyelination and other changes. Acute was a group in which no demyelination was observed, subacute 1 group was a group in which demyelination is less than 30% of the entire specimen, subacute 2 group was a group in which demyelination was in 30 to 70%, and subacute 3 was a group in which demyelination was 70% or more. Namely, we classified the disease into acute, subacute 1, subacute 2, subacute 3, and chronic groups based on HE staining and LFB staining. As a result, the 28 dogs of CDV infection were subdivided into 4 acute cases, 4 subacute 1 cases, 9 subacute 2 cases, 5 subacute 3 cases, and 6 chronic cases.

2.2. Histopathology and Immunohistochemistry

Routine autopsy was performed soon after death and selected tissues including cerebellum were fixed in 10% buffered neutral formalin solution and routinely processed for histologic examination. Tissue sections were stained with hematoxylin and eosin (HE), as well as, with LFB stains according to Verdes et al. [21].

For immunohistochemistry, a mouse anti-CDV monoclonal antibody (Biorad, MCA 1893, California, USA) was used, following by a conjugated secondary antibody in a HRP polymer detection system (Mouse-on-canine HRP-polymer, Biocare Medical, California, USA), positive antigen-antibody reactions were observed by incubation with 3,3-diaminobenzidine-

tetrahydrochloride (DAB) as described previously [22]. Glia cells were immunostained using specific primary antibodies against intermediate filaments of astrocytes (GFAP) and microglia (Iba1). According to Verdes et al. [23], a primary antibody against the calcium binding protein Calbindin D 28k (CbD28k), was used as a marker to identify and count Purkinje cells. Details of type and origin of primary antibodies, used dilution, and sources, are summarized in Table 1.

Table 1. Summary of used primary antibodies and detection systems.

Primary Antibody	Source	Dilution	Species / type	Detection system
MCA 1893 Canine Distemper Virus	Biorad	1:250	Mouse / monoclonal	Mouse on canine HRP-Polymer
Glial Fibrillary Protein (GFAP)	Biocare Medical	RTU*	Mouse / monoclonal	MACH 4 Universal HRP-Polymer
Anti-Iba1 (ab5076)	Abcam	1:1000	Goat / polyclonal	Rabbit Anti-Goat IgG H&L (HRP) ab6741
Calbindin (orb5912)	Biorbyt	1:200	Rabbit / polyclonal	MACH 4 Universal HRP-Polymer

* Ready to use.

In addition, four pathologists (JMV, CL, BV, KY) classified the stages of the histopathological lesions following as acute, subacute, or chronic based on the previous studies [3,10,11,14–16,19,24]. Namely, acute lesions were characteristic by focal vacuolation, gliosis, inclusion bodies and CDV-positive cells, subacute lesions were demyelination, gliosis, necrosis, perivascular mononuclear infiltration (cuffing) of two to three layers of thickness, inclusion bodies and CDV-immunopositive cells, and chronic lesions were similar to the subacute stage, but perivascular infiltration was more prominent, of at least three layers of thickness.

Changes in the numbers of astrocytes, microglia, and Purkinje cells were examined based on the results of the above immunohistochemical staining. The area to be examined was the white matter adjacent to the fourth ventricle for astroglia and microglia, where the changes first appeared and where the changes were characteristic. Purkinje cells were evaluated in cerebellar cortex selecting the neighbour cerebellar folia to the same anatomical region studied to evaluate glial cells.

2.3. Image Capture and Analysis

All histological slides were scanned with a slide scanner (Motic Easy Scan One®, China) for subsequent analysis, the whole slide images were viewed and captured using the Motic DSAssistant® software (Motic VM V1 Viewer 2.0®, version 2019-08-02, China).

Astrocytes, microglia and Purkinje cells were counted in canine cerebellar sections, based on the results of the specific immunohistochemistry against GFAP (astrocytes marker), Iba-1 (microglia marker), and CbD28k (Purkinje cell marker).

The counting was done manually in images captured in 400x fields for astrocytes and microglia. The examined region included the fourth ventricle region and the central white matter, capturing two images per animal, one for each area. For the counting of Purkinje cells, a whole image of a cerebellar folia per animal was used, in fields viewed at 40x.

2.4. Statistical Analysis

Statistical analysis was performed using GraphPad Prism® (version 10.1.1 -323- for Windows 64-bit, GraphPad Software, Boston, Massachusetts USA, www.graphpad.com). One-way ANOVA tests followed by multiple comparisons using Tukey’s test were used to assess the difference between groups. Data are presented as means ± standard deviation. The level of statistical significance used in all studies was $p \leq 0.05$.

3. Results

3.1. Relationship between Each Stage and Age or Sex

There was no relationship between the stage and the dog’s age, and there was no tendency for acute to be young and chronic to be old. For example, an animal was died at 45 days of age in the chronic stage. There was also no relationship between each stage and sex. Many of the dogs were mixed breed, and there was no relationship between each stage and a specific breed. In the control group, the numbers of astrocytes, microglia, and Purkinje cells were not related to age, breed, sex, or body weight.

3.2. Histopathological and Immunohistochemical Findings

A summary of histological findings is shown in Table 2.

Table 2. Histological findings in cerebellum in each stage, including a summary of number of studied dogs and average age in each group.

Groups	Number of animals	Average month age (range)	Histopathological findings
Control	5	(45 days old – 6 years old)	No changes
Acute	4	(25 days old – 3 years old)	glial cell reaction, inclusion bodies, CDV immunoreactive particles, meningitis
Subacute 1	4	(30 days old – 3 years old)	demyelination, glial cell reaction, inclusion bodies, CDV immunoreactive particles, meningitis
Subacute 2	9	(2 months old – 10 years old)	demyelination, glial cell reaction, inclusion bodies, CDV immunoreactive particles, meningitis, mild decreased granular layer cells and Purkinje cells
Subacute 3	5	(3 months old – 3 years old)	demyelination, glial cell reaction, inclusion bodies, CDV immunoreactive particles, meningitis, moderate decreased granular layer cells and Purkinje cells
Chronic	6	(3 months old – 3 years old)	demyelination, gliosis, glial cell reaction, inclusion bodies, CDV immunoreactive particles, meningitis, severe decreased granular layer cells and Purkinje cells

A summary of immunohistochemical study is shown in Tables 3, and the graphical representation of total number of GFAP-immunoreactive astrocytes in each stage, with statistically significant differences observed between the control vs chronic group $p=0.0002$ (Figure 1a), and representative immunohistochemistry against GFAP in a cerebellum of normal dog (Figure 1b) and cerebellum of a dog with chronic stage (Figure 1c) are shown. In Figure 2, the graphical representation of total number of Iba-1-immunoreactive microglia in each stage, without statistically significant differences between the groups (Figure 2a), and representative immunohistochemistry against Iba-1 in a cerebellum of a dog of subacute group 2 (Figure 1b and 1c) are shown. In Figure 3a, the graphical representation of total number CbD28k-immunoreactive Purkinje cells in each stage, with statistically significant differences observed between the control vs subacute group 2 ($p=0.0144$), control vs subacute 3 ($p=0.0024$) and between the control vs chronic groups ($p=0.0163$). Representative immunohistochemistry against CbD28k-immunoreactive Purkinje cells in a cerebellum of normal dog (Figure 3b) and cerebellum of CDV infected dog from subacute 3 stage (Figure 3c) are shown.

Table 3. Numbers of each cell type in cerebellum in each stage. Values (mean ± standard deviation) marked with different letters within the same column are significantly different (P < 0.05).

Groups	Number of animals	Average month age (range)	Numbers		
			astrocytes	microglias	Purkinje cells
Control	5	(45 days old – 6 years old)	35 ± 11 ^a	64 ± 19	49 ± 6 ^a
Acute	4	(25 days old – 3 years old)	41 ± 12	64 ± 13	42 ± 17
Subacute 1	4	(30 days old – 3 years old)	42 ± 13	56 ± 16	25 ± 6
Subacute 2	9	(2 months old – 10 years old)	36 ± 13	75 ± 26	24 ± 14 ^b
Subacute 3	5	(3 months old – 3 years old)	32 ± 12	63 ± 26	15 ± 12 ^c
Chronic	6	(3 months old – 3 years old)	63 ± 22 ^b	44 ± 17	23 ± 13 ^d




Figure 1. (a) Comparative number of astrocytes with positive immunostaining against GFAP in the fourth ventricle and central white matter region between the control, acute, subacute 1, subacute 2, subacute 3 and chronic animal groups. Statistically significant differences were observed between the control vs chronic group $p=0.0002$. **(b)** Representative GFAP-immunohistochemistry of cerebellar white matter of a normal dog (control group). Immunolabeling for GFAP is observed in normal astrocytes. Scale = 50 μ m. See details in the upper right inset of the figure. **(c)** Representative GFAP-immunohistochemistry of cerebellar white matter of an CDV infected dog (chronic group). Immunostaining for GFAP is observed in reactive astrocytes, showing an increase in cytoplasmic volume in gemistocytic astrocytes. Scale = 50 μ m. See details in the upper right inset of the figure.




Figure 2. (a) Comparative number of microglia with positive immunostaining against Iba1 in the region of the fourth ventricle and the central white matter between the control, acute, subacute 1, subacute 2, subacute 3 and chronic animal groups. No significant differences were found. **(b)** Representative Iba-1-immunohistochemistry of cerebellar white matter of a CDV infected dog from subacute group 2. Immunostaining against iba-1 is observed in the periventricular region, the microglia have an amoeboid shape. Scale = 60µm. See details in the upper right inset of the figure. **(c)** Representative Iba-1-immunohistochemistry of cerebellar white matter of a CDV infected dog (chronic group). Same animal and region of **(b)**. Intense immunostaining against Iba-1 is observed in the cerebellar white matter. Scale = 60 µm. See details in the upper right inset of the figure.




Figure 3. (a) Comparative number of Purkinje cells with positive immunostaining against CbD28k in the cerebellar cortex between the control, acute, subacute 1, subacute 2, subacute 3 and chronic animal groups. Statistically significant differences were observed between the control vs subacute group 2 ($p=0.0144$), control vs subacute 3 ($p=0.0024$) and between the control vs chronic groups ($p=0.0163$). **(b)** Representative Calbindin (CbD28k)-immunohistochemistry of cerebellar cortex of a normal dog (control group). Positive immunostaining against CbD28k is observed in all the Purkinje cells (both in the soma, axonal process and dendritic arborization). Scale = 100 μ m. See details in the upper right inset of the figure. **(c)** Representative CbD28k-immunohistochemistry of cerebellar cortex of a CDV infected dog from subacute group 3. Positive immunostaining against CbD28k is observed in all

Purkinje cells, showing a reduction in the number of immunoreactive Purkinje cells. Scale = 100 μ m. See details in the upper right inset of the figure.

LFB staining at each stage is shown in Figures 4a to 4f, demyelinating changes in white matter are shown in Figure 5, and CDV immunoreactive particle deposition is shown in Figures 8 a to 8 d.




Figure 4. (a) Demyelination is not detected. No. 397. Control group. LFB. Scale=60 μ m. See the asterisk. (b) Demyelination is not apparent. No. 010. Acute group. LFB. Scale=60 μ m. See the asterisk. (c) Slight demyelination is observed. No. 366. Subacute 1 group. LFB. Scale=60 μ m. See the asterisk. (d) Demyelination is apparent. No. 463. Subacute 2 group. LFB. Scale=100 μ m. See the asterisk. (e) Demyelination is progressed. No. 23. Subacute 3 group. LFB. Scale=100 μ m. See the asterisk. (f) Demyelination is severe. No. 11. Chronic group. LFB. Scale=100 μ m. See the asterisk.




Figure 5. Demyelination is apparent around blood vessel (asterisk) showing cuffing (black arrow). Dog No. 11. chronic group. LFB. Scale=90µm.

Histologically, the main lesions were demyelination, glial cell reaction, reduced number of Purkinje cells and meningitis.

In acute group, all animals subjected to PCR testing were positive. No demyelination and vacuoles were observed by HE and LFB stainings. CDV particles were detected in astrocytes, microglia, neurons, pia mater and choroid plexus. In addition, CDV immunoreactive particles were observed in the white matter adjacent to the fourth ventricle in all cases, and are also observed in endothelial cells of blood vessels, mononuclear cells inside and outside blood vessels in the white matter away from the fourth ventricle, white matter of cerebellar lobes, and granular layer. Nuclear and cytoplasmic inclusions were observed mainly in astrocytes, microglia, and pia mater. Although there was no significant increase in the number of astrocytes and microglia compared to those in the control group, they were enlarged compared to cells in the control group. CDV immunoreactive granules were observed in these enlarged cells.

In subacute 1 group, changes were similar to those in the acute stage, but demyelination was apparent by LFB staining. Changes were occurring in the white matter adjacent to the fourth ventricle. In one case, CDV immunoreactive granules infiltrated from the pia mater into the gray matter, and were also observed in Purkinje cells, and in cells of the granular layer. Although there was no significant increase in the number of astrocytes and microglia compared to the acute group, many enlarged cells were observed.

In subacute 2 group, demyelination occurred frequently in areas other than the white matter adjacent to the fourth ventricle, and demyelination in the white matter of the cerebellar folia was observed in more than half of the cases. Demyelination and associated changes in astrocytes and microglia progressed around blood vessels, such that areas of demyelination were scattered throughout the white matter. Although there was no significant increase in astrocytes and microglia compared to the acute and subacute 1 groups, gemistocytic astrocytes, multinuclear astrocytes, fibrillary gliosis, and rod cells were observed. Foamy cells also appeared in areas where changes were progressing. LFB staining revealed swelling, collapse, and disappearance of the myelin sheath around the myelinated axons. CDV immunoreactive particles and inclusion bodies were increased compared to the subacute 1 group.

In the subacute 3 group, the changes in the subacute 2 group progressed, and the area of demyelination increased and spread throughout the white matter. Although no significant increase in the number of astrocytes was observed, many gemistocytic astrocytes and multinuclear astrocytes were observed (Figure 6). In addition, CDV particles, increase in inclusion bodies, and CDV particles

and inclusion bodies were increased. Although lymphocytes, plasma cells, monocytes, and macrophages were aggregated around blood vessels, the number of these cells was 1 to 2 layers.

In the chronic group, changes such as severe demyelination, a significant increase in the number and enlargement of astrocytes and microglia, an increase in CDV immunoreactive particles, and an increase in inclusion bodies were progressing. The perivascular cuffing showed three or more layers.

Meningitis was observed in 1 subacute 1, 3 subacute 2, 3 subacute 3, and 3 chronic cases. Mononuclear cells with CDV immunoreactive particles were detected in and around blood vessels.

As described, astrocytes and microglia did not increase in number compared to the control group, but were enlarged in the acute group. Furthermore, gemistocytic astrocytes, multinuclear astrocytes, fibrillary gliosis, and rod cells were observed in the subacute groups. Demyelination was observed in the white matter of the cerebellar folia depending on the stages, and CDV infiltration from the pia mater into the gray matter was also observed. According to these changes, a decrease in Purkinje cells and necrotic changes in the granular layer were detected (Figure 7).




Figure 6. Marked increase in the number of astrocytes and microglia is detected in the white matter showing demyelination. Astrocytes are enlarged, recognized as gemistocytes (asterisk), and multinucleated cells are also seen. Nuclear inclusion bodies are observed (black arrow). Dog No. 436. subacute 2 group. HE. Scale=30µm.




Figure 7. The white matter stains on some cerebellar folia have disappeared (asterisk). The number of cells in the granular layer decreases with decreased number of Purkinje cells. No. 15. chronic group. HE. Scale=100µm.




Figure 8. (a) Canine distemper virus (CDV)-immunopositive particles are detected in the subventricular parenchyma(asterisk). Immunohistochemical staining for CDV. No. 53. Acute group. Scale=60µm. (b) Canine distemper virus (CDV)-positive particles are observed mainly in astrocytic podocytes around blood vessel arrowhead). Vascular endothelial cells and intravascular mononuclear cell also show CDV-immunopositivity. Immunohistochemical staining for CDV. No. 53. Acute group. Scale=30µm. (c) Many canine distemper virus (CDV)-immunopositive particles are observed in glial cells around blood vessels. No. 249. Subacute 3 group. Scale=100µm. (d) Many canine distemper virus (CDV)-immunopositive particles are observed in the three layers (molecular, Purkinje cells and granular) of gray matter and white matter (see asterisk in molecular layer). Immunohistochemical staining for CDV. No. 6. Subacute 3 group. Scale=300µm.

4. Discussion

Our results showed that in the cerebellum, demyelination, reactions of astrocytes, oligodendroglia, and microglia, the appearance of inclusion bodies, a reduction in the number of Purkinje cells, and degeneration of the granular layer were observed. Demyelination was evident with LFB staining, and was observed as a decrease in staining with HE staining in the subacute 1 group. Demyelination was consistent with vacuolization in the subacute 2 group and later. Astrocytes and microglia were hypertrophic even in the acute stage, when no demyelination was observed, and progressed with the stage. It has been reported that in demyelinating distemper lesions, the majority (95%) of infected cells have been identified as astrocytes, representing the main target for CDV [25,26]. In this study, CDV immunoreactive granules were observed not only in astrocytes but also in microglia at the early stage of infection. Furthermore, it has been said that astrocytes and microglia responded during the early stages of infection, and there are also reports that their numbers increased [2,3,20]. In this study, there was no significant increase in the number of astrocytes and microglia between the control group and each infection group, but these cells changed morphologically from an early stage, and CDV immunoreactive granules were also observed in these cells. It is clear that normal function is impaired. The fact that morphological changes, rather than increases in the number of astrocytes and microglia, were a characteristic feature is considered to be important information obtained in this study. Various functions of astrocytes have been reported, among which the key function of astrocytes is said that the removal of neurotransmitters, such as glutamate, from

the synaptic cleft by specific transporters and subsequent degradation by glutamine synthetase (glutamate - glutamine cycle), Thus preventing excitotoxic cell death of neurons and myelin producing oligodendrocytes [16,27]. In this study, astrocyte responses were observed before demyelination, so it is apparent that astrocytes are involved in myelin changes from the early stage of infection.

Cytoplasmic and intranuclear inclusion bodies could be frequently found in acute and subacute lesions [28]. In the present study, CDV granules and inclusion bodies were observed from the acute stage, before the onset of demyelination, and progressed with the stage. It has been reported that viral antigen may disappear from the inflammatory demyelinating lesions due to the antiviral immune responses in chronic CDV encephalomyelitis [29,30]. However, the responses of astrocytes and microglial cells, the increase in the number of CDV immunoreactive particles within these cells, and the appearance of inclusion bodies did not decrease until the chronic stage in this study. Therefore, it was clear that changes of astrocytes and microglial cells appeared at an early stage and progressed with the stage. Two possible routes for demyelination have been reported that one route is that CDV infects astrocytes, which then infects oligodendroglia, and the other route is that CDV infects microglia through astrocytes and ultimately affects oligodendroglia [3]. Our results do not support either of these possibilities.

Changes were initially observed in the periventricular region and gradually became perivascular in the white matter. The periventricular region is said to be the first area that the virus infects. The main route of neural entry is via infected mononuclear cells that cross the blood-brain barrier, resulting in local viral release and subsequent infection of resident epithelial and endothelial cells. It has also been reported that the virus can spread through the brain once it enters the brain. It infects the cerebrospinal fluid (CSF), infects the ependymal lining cells of the ventricles, and ultimately infects glial cells and neurons [19,31–34]. On the other hand, demyelination, astrocyte and microglial reactions, and myelin degeneration were also observed in the white matter of cerebellar folia in the subacute 2 group. Furthermore, in the subacute 1 group, infiltration of CDV immunoreactive granules from the leptomeninges into the gray matter was observed. In these cases, an astrocytic response was also observed in the gray matter, and cell loss in the granular layer was also observed. In cerebellar folia, CDV immunoreactive particles were found in astrocytes, Golgi cells, and Purkinje cells. CDV immunoreactive granules have been reported to extend from pial cells to the subpial gray matter [5,13,28,34,35]. We confirmed CDV infection from the leptomeninges to the gray and white matter described in our previous report [18].

In this study, we observed changes to Purkinje cells. The number of Purkinje cells decreased more significantly than in the subacute 2 group and gradually progressed. This fact was thought to be due not only to the influence of CDV infection in the white matter to the gray matter, but also to CDV infection from the leptomeninges to the gray and white matter. Effects on Purkinje cells have been reported in viral infections in various species of animals [7,36]. Focal vacuolation and spongy appearance near the Purkinje cell layer in the cerebellum have also been observed in dogs with CDV infection [4], and Rudd et al. [17] also stated that Purkinje cells and granule cells are specifically targeted by CDV infection. In addition, changes in Purkinje cells and granular layers have been reported in CDV infection in other animal species [37], and the changes in Purkinje cells and granule cells have been observed in the CDV infection experimental study in ferrets [38]. Based on the above reports and the present results, changes in Purkinje cells and the granular layer can be said to be one of the characteristics of CDV infection.

Age and each stage of disease have been discussed based on the dog's immune status. Young or immunocompromised dogs are said to show acute lesions, while mature dogs usually develop the chronic encephalomyelitis [1,11,20,39]. This study did not measure antibodies in the blood. Furthermore, there may be an issue with the strain of the CDV virus. In this study, the samples in the acute group ranged from 25 days after birth to 3 years old, and the onset at age 10 was identified in the subacute 2 group, instead of in the chronic group. These facts suggest that the stage of the lesions are related to the immune status due to various factors, and it seems unlikely that age alone is a factor associated with each stage.

5. Conclusions

Astrocyte and microglial responses were observed at an early stage when demyelination was not evident. CDV immunoreactive granules and inclusion bodies were also observed in various cells from the acute stage. A decrease in Purkinje cells and degenerative changes in the granular layer were also observed with demyelination. These changes were progressing along with the stage. There was no clear relationship between age and stage.

Author Contributions: Conceptualization, J.M.V.; C.L.; G.G.; K.Y.; methodology, J.M.V.; C.L.; G.G.; B.V.; V.I.; V.Y.; L.D.; K.Y.; software, C.L.; G.G.; validation, J.M.V.; C.L.; G.G.; B.V.; K.Y.; formal analysis, C.L.; G.G.; J.M.V.; K.Y.; investigation, C.L.; G.G.; J.M.V.; K.Y.; resources, J.M.V.; L.D.; data curation, J.M.V.; C.L.; G.G.; B.V.; V.I.; V.Y.; L.D.; K.Y.; writing—original draft preparation, C.L.; J.M.V.; K.Y.; writing—review and editing, C.L.; J.M.V.; K.Y.; visualization, J.M.V.; C.L.; G.G.; B.V.; K.Y.; supervision, K.Y.; J.M.V.; project administration, J.M.V.; L.D.; funding acquisition, J.M.V.; L.D.

Funding: This research was funded by National Agency of Research and Innovation (ANII, Fondo María Viñas 2019), Grant FMV-1-2019-1-155934), C.L. was MSc fellowship recipient of National Agency of Research and Innovation (ANII). V.I. is PhD fellowship recipient of Udelar's Posgraduate Academic Commission (Comisión Académica de Posgrados, CAP-Udelar). J.M.V. is Research Career Member of the National Research System (SNI-ANII), Uruguay, and Program for the Development of Basic Sciences (PEDECIBA), Uruguay.

Institutional Review Board Statement: The animal study protocol was approved by the Institutional Review Board of Veterinary Faculty's Animal Use Ethics Committee (Comisión de Ética en el Uso de Animales de la Facultad de Veterinaria, CEUA-FVET, Udelar, Montevideo, Uruguay), protocol code: CEUA FVET-Udelar N° 320/16. Date of approval: October 10th., 2016.

Acknowledgments: The authors would like to thank all the owner's that agreed to donate their dogs to participate in this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Carvalho, O.V.; Botelho, C.V.; Ferreira, C.G.T.; Scherer, P.O.; Soares-Martins, J.A.P.; Almeida, M.R.; Júnior, A.S. Immunopathogenic and neurological mechanisms of canine distemper virus. *Adv Virol* 2012, 163860.
2. Klemens, J.; Ciurkiewicz, M.; Chludzinski, E.; Iseringhausen, M.; Klotz, D.; Pfankuche, V.M.; Ulrich, R.; Herder, V.; Puff, C.; Baumgärtner, W.; Beineke, A. Neurotoxic potential of reactive astrocytes in canine distemper demyelinating leukoencephalitis. *Sci Rep* 2019, 9, 11689.
3. Lempp, C.; Spitzbarth, I.; Puff, C.; Cana, A.; Kegler, K.; Techangamsuwan, S.; Baumgärtner, W.; Seehusen, F. New Aspects of the Pathogenesis of canine distemper leukoencephalitis. *Viruses* 2014, 6, 2571-2601.
4. Pan, Y.; Liu, X.Y.; Meng, L.P.; Zhu, G.R.; Xia, Y.K.; Chen, J.S.; Yoshikawa, T. Pathogenesis of demyelinating encephalopathy in dogs with spontaneous acute canine distemper. *J Integr Agric* 2013, 12, 334-343.
5. Vandeveld, M.; Zurbriggen, A. Demyelination in canine distemper virus infection: A review. *Acta Neuropathol* 2005, 109, 56-68.
6. De Nardo, T.F.S.; Bertolo, P.H.L.; Bernardes, P.A.; Munari, D.P.; Machado, G.F.; Jardim, L.S.; Moreira, P.R.R.; Rosolem, M.C.; Vasconcelos, R.O. Contribution of astrocytes and macrophage migration inhibitory factor to immune-mediated canine encephalitis caused by the distemper virus. *Vet Immunol Immunopathol* 2020, 221, 110010.
7. Zurbriggen, A.; Schmid, I.; Graber, H.U.; Vandeveld, M. Oligodendroglial pathology in canine distemper. *Acta Neuropathol* 1998, 95, 71-77.
8. Silva, M.C.; Figuera, R.A.; Mazzanti, A.; . Neuropatologia da cinomose canina: 70 casos (2005-2008). *Pesq Vet Bras* 2009, 29, 643-652.
9. Kabakci, N.; Yarim, M.; Karahan, S.; Guvenc, T.; Yagci, B.B.; Gurcan, I.S. Immunohistochemical investigation of cerebellum in dogs infected with canine distemper virus. *Acta Veterinaria Hungarica* 2004, 52, 327-337.
10. Ulrich, R.; Puff, C.; Wewetzer, K.; Kalkuhl, A.; Deschl, U.; Baumgärtner, W. Transcriptional changes in canine distemper virus-induced demyelinating leukoencephalitis favor a biphasic mode of demyelination. *PLOS One* 2014, 9, e95917.
11. Vandeveld, M.; Fankhauser, R.; Kristensen, F.; Kristensen, B. Immunoglobulins in demyelinating lesions in canine distemper encephalitis. An immunohistological study. *Acta Neuropathol* 1981, 54, 31-41.
12. Bathen-Noethen, A.; Stein, V.M.; Puff, C.; Baumgärtner, W.; Tipold, A. Magnetic resonance imaging findings in acute canine distemper virus infection. *J Small Anim Pract* 2008, 49, 460-467.

13. Beineke, A.; Puff, C.; Seehusen, F.; Baumgärtner, W. Pathogenesis and immunopathology of systemic and nervous canine distemper. *Vet Immunol Immunopathol* 2009, 127, 1-18.
14. Alldinger, S.; Fonfara, S.; Kremmer, E.; Baumgärtner, W. Up-regulation of the hyaluronate receptor CD44 in canine distemper demyelinated plaques. *Acta Neuropathol* 2000, 99, 138-146.
15. Gröters, S.; Alldinger, S.; Baumgärtner, W. Up-regulation of mRNA for matrix metalloproteinases-9 and -14 in advanced lesions of demyelinating canine distemper leukoencephalitis. *Acta Neuropathol* 2005, 110, 369-382.
16. Spitzbarth, I.; Lempp, C.; Kegler, K.; Ulrich, R.; Kalkuhl, A.; Deschl, U.; Baumgärtner, W.; Seehusen, F. Immunohistochemical and transcriptome analyses indicate complex breakdown of axonal transport mechanisms in canine distemper leukoencephalitis. *Brain Behav* 2016, 3, e00472.
17. Rudd, P.A.; Bastien-Hamel, L.E.; von Messling, V. Acute canine distemper encephalitis is associated with rapid neuronal loss and local immune activation. *J Gen Virol* 2010, 91, 980-989.
18. Feijóo, G.; Yamasaki, K.; Delucchi, L.; Verdes, J.M. Central nervous system lesions caused by canine distemper virus in 4 vaccinated dogs. *J Vet Diagn Invest* 2021, 33, 640-647.
19. Summers, B.A.; Greisen, H.A.; Appel, M.J. Early events in canine distemper demyelinating encephalomyelitis. *Acta Neuropathol* 1979, 46, 1-10.
20. Thomas, W.B.; Sorjonen, D.C.; Steiss, J.E. A retrospective evaluation of 38 cases of canine distemper encephalomyelitis. *J Am Anim Hosp Assoc* 1993, 29, 129-133.
21. Verdes, J.M.; Larrañaga, C.; Varela, B.; Iribarnegaray, V.; Yozzi, V.; Feijóo, G.; Yamasaki, K. Histopathological analysis of brains from dogs infected with canine distemper virus. In: Ma D. y Pfaller, CK. Eds. *Measles and Related Morbilliviruses: Methods and Protocols*. Springer Nature. New York, USA, 2024, Volume 2808, Chapter 14, pp. 177-195.
22. Fairley, R.A.; Knesl, O.; Pesavento, P.A.; Elias, B.C. Post-vaccinal distemper encephalitis in two Border Collie cross littermates. *N Z Vet J* 2015, 63, 117-20.
23. Verdes, J.M.; de Sant'Ana, F.J.F.; Sabalsagaray, M.J.; Okada, K.; Calliari, A.; Moraña, J.A.; de Barros, C.S.L. Calbindin D28k distribution in neurons and reactive gliosis in cerebellar cortex of natural rabies virus-infected cattle. *J Vet Diagn Invest* 2016, 28, 361-8.
24. Seehusen, F.; Orlando, E.A.; Wewetzer, K.; Baumgärtner, W. Vimentin-positive astrocytes in canine distemper: a target for canine distemper virus especially in chronic demyelinating lesions? *Acta Neuropathol* 2007, 114, 597-608.
25. Mutinelli, F.; Vandevelde, M.; Griot, C.; Richard, A. Astrocytic infection in canine distemper virus-induced demyelination. *Acta Neuropathol* 1989, 77, 333-335.
26. Summers, B.A.; Appel, M.J. Demyelination in canine distemper encephalomyelitis: an ultrastructural analysis. *J Neurocytol* 1987, 16, 871-881.
27. Correale, J.; Farez, M.F. The Role of Astrocytes in multiple sclerosis progression. *Front Neurol* 2015, 6, 180.
28. Baumgärtner, W.; Orvell, C.; Reinacher, M. Naturally occurring canine distemper virus encephalitis: distribution and expression of viral polypeptides in nervous tissues. *Acta Neuropathol* 1989, 78, 504-512.
29. Koutinas, A.F.; et al. Relation of clinical signs to pathological changes in 19 cases of canine distemper encephalomyelitis. *J Comp Pathol* 2002, 126, 47-56.
30. Muller, C.F.; Fatzer, R.S.; Beck, K.; Vandevelde, M.; Zurbriggen, A. Studies on canine distemper virus persistence in the central nervous system. *Acta Neuropathol* 1995, 89, 438-445.
31. Axthelm, M.K.; Krakowka, S. Canine distemper virus: the early blood-brain barrier lesion. *Acta Neuropathol* 1987, 75, 27-33.
32. Frisk, A.L.; König, M.; Moritz, A.; Baumgärtner, W. Detection of canine distemper virus nucleoprotein RNA by reverse transcription-PCR using serum, whole blood, and cerebrospinal fluid from dogs with distemper. *J Clin Microbiol* 1999, 37, 3634-3643.
33. Higgins, R.J.; Krakowka, S.G.; Metzler, A.E.; Koestner, A. Primary demyelination in experimental canine distemper virus induced encephalomyelitis in gnotobiotic dogs. Sequential immunologic and morphologic findings. *Acta Neuropathol* 1982, 58, 1-8.
34. Rudd, P.A.; Cattaneo, R.; von Messling, V. Canine distemper virus uses both the anterograde and the hematogenous pathway for neuroinvasion. *J Virol* 2006, 80, 9361-9370.
35. Summers, B.A.; Greisen, H.A.; Appel, M.J. Canine distemper encephalomyelitis: variation with virus strain. *J Comp Pathol* 1984, 94, 65-75.
36. Wünschmann A, Lopez-Astacio, R.; Armien, A.G.; Reed, L.; Parrish, C.R. Parvovirus-induced encephalitis in a juvenile raccoon. *J Vet Diagn Invest* 2021, 33, 140-143.
37. Potgieter, L.N.; Patton CS. Multifocal cerebellar cortical necrosis caused by canine distemper virus infection in a raccoon. *J Am Vet Med Assoc* 1984, 185, 1397-1399.
38. Dudd, P.A.; Bastien-Hamel, L.E.; von Messling, V. Acute canine distemper encephalitis is associated with rapid neuronal loss and local immune activation. *J Gen Virol* 2010, 91, 980-989.
39. Vandevelde, M.; Zurbriggen, A. The neurobiology of canine distemper virus infection. *Vet Microbiol* 1995, 44, 271-280.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.