

Review

Not peer-reviewed version

Role of Environmental Factors in Legume-Rhizobium Symbiosis. A review

[Liudmyla Yermko](#), [Katarzyna Czopek](#)^{*}, [Mariola Staniak](#), [Mykola Marenych](#), [Volodymyr Hanhur](#)

Posted Date: 14 December 2024

doi: 10.20944/preprints202412.1142.v1

Keywords: legumes, *Rhizobium*, biological nitrogen fixation BNF, abiotic stress, soil nutrients, sustainable agriculture, protein crops



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Review

Role of Environmental Factors in Legume-Rhizobium Symbiosis. A review

Liudmyla Yeremko ¹, Katarzyna Czopek ^{2,*}, Mariola Staniak ², Mykola Marenych ³ and Volodymyr Hanhur ¹

¹ Department of Crop Production, Poltava State Agrarian University, Skovoroda St., 1/3, 36000 Poltava, Ukraine

² Department of Crops and Yield Quality, Institute of Soil Science and Plant Cultivation—State Research Institute, 8 Czarotoryskich St., 24-100 Pulawy, Poland

³ Department of Breeding, Seed Production and Genetics, Poltava State Agrarian University, Skovoroda St., 1/3, 36000 Poltava, Ukraine

* Correspondence: kczopek@iung.pulawy.pl

Abstract: Legumes play a pivotal role in addressing global challenges of food and nutrition security by offering a sustainable source of protein and bioactive compounds. The capacity of legumes to establish symbiotic relationships with *Rhizobium* bacteria enables biological nitrogen fixation (BNF), reducing the dependence on chemical fertilizers while enhancing soil health. However, the efficiency of this symbiosis is significantly influenced by environmental factors, such as soil acidity, salinity, temperature, moisture content, light intensity, and nutrient availability. These factors affect key processes, including rhizobia survival, nodule formation, and nitrogenase activity, ultimately determining the growth and productivity of legumes. This review synthesizes current knowledge on legume-rhizobium interactions under varying abiotic conditions. It highlights the impact of salinity and acidity in limiting nodule development, soil temperature in regulating microbial community dynamics, and moisture availability in modulating metabolic and hormonal responses during drought and waterlogging. Moreover, the role of essential nutrients, including nitrogen, phosphorus, potassium, and trace elements such as iron, molybdenum, and boron, in optimizing symbiosis is critically analyzed.

Keywords: legumes, *Rhizobium*, biological nitrogen fixation BNF, abiotic stress, soil nutrients, sustainable agriculture, protein crops

1. Introduction

In solving global food and nutrition security problems, the cultivation of legumes as the main source of highly nutritious protein resources, is gaining strategic importance. It is well known that protein, as an important nutrient, that plays a key role in the physiological and biochemical reactions in the organism necessary to ensure the body growth and development [1,2]. In the structure of protein resources used for nutrition, proteins of animal origin play a significant role. However, according to the results of scientific research, the presence of a large amount of meat in the human diet can lead to obesity and disorders of the functioning of the cardiovascular system related to it. On the other hand, the so-called hidden hunger caused by insufficient micronutrients in the daily diet leads to many health problems, especially for children. They can be manifested in stunted growth, poor weight gain, cognitive impairment, and mental disorders [3,4]. One of the components, that can ensure a healthy diet for the population of both developed and developing countries can be highly nutritious plant products made from legume seeds [5].

The main constituent components of the legume seeds are protein, complex carbohydrates, dietary fibre, vitamins, minerals and biologically active compounds. Legume seeds are characterized by the absence of cholesterol and, with the exception of soybeans, chickpeas, white lupine and peanuts, contain a low amount of fat [6]. Biologically active compounds of legumes (including phytoestrogens, saponins, phenolic compounds, oligosaccharides, alkaloids) are characterized by antithrombotic, antioxidant, opioid-like activity, have cytomodulatory or immunomodulatory effects

and contribute to the improvement of mineral bioavailability [7–9]. Pharmaceuticals based on their raw materials are widely used in the prevention of occurrence and medical therapy of cardiovascular, cancer diseases, diabetes and obesity [10,11]. At the same time, legumes are a valuable source of highly nutritious feed resources used in livestock farming [12,13].

Cultivation of legumes improves the biological, physical and chemical characteristics of the soil [14,15], contributes to the reduction of greenhouse gases in the atmosphere [16], and limits the occurrence of diseases, pests, diseases and nematodes by disrupting the biological cycle of their development [17]. Legumes have a unique biological property, which consists in establishing symbiotic relationships with nitrogen fixing bacteria of the genus *Rhizobium* (*Rhizobium*). Kebebe [18], notes that in the process of legume-*Rhizobium* symbiosis, legumes can fix about 100 to 300 kg ha⁻¹ of atmospheric nitrogen (N₂) annually, thus providing about 139-175 million tons of nitrogen to the soil. This, in turn, reduces the cost of applying about 80-90 tons of mineral nitrogen fertilizers annually [19]. Nitrogen (N) is an element that occurs mainly in the atmosphere, lithosphere and biosphere in various degrees of oxidation [20,21]. Only about 2% of this element is found in living organisms and their biocoenoses, with the remainder occurring in inorganic form [22]. Nitrogen in the gaseous form (it makes up about 78% of the atmospheric composition) is very persistent due to the triple covalent bond between N atoms, but in this form it is unabsorbable by most higher organisms and unreactive [23]. In bioavailable form, N is found in soil in the form of organic and inorganic compounds such as nitrates –NO₃, nitrites –NO₂ and ammonium ions –NH₄ [24,25]. Biological nitrogen fixation (BNF) is one of the most important processes on Earth. It involves the conversion of N₂, which is not assimilated by plants, into ammonia, the form available to plants [26,27]. This process is a highly energetic transformation, requiring reducing forces, i.e. electrons and protons (H⁺), as well as large amounts of energy (ATP) [28]. The symbiotic relationship between *Rhizobium* and legumes plays a key role in sustainable agriculture by facilitating access to N₂, improving soil fertility and reducing the need for chemical fertilizers.

After the mutualistic relationship between the two symbiosis partners is established, the roots of plants form structures known as root nodules, in which MgATP-dependent N₂ reduction by the nitrogenase complex occurs (Figure 1).



Figure 1. Nodules on soybean roots [Photo: K. Czopek].

The produced ammonia is used by plants as a source of N in the synthesis of organic compounds. To support the bacterial endosymbionts' metabolism, photosynthetic products are supplied from the plant to nodules [29]. The establishment of the legume-*Rhizobium* symbiosis as an N₂-fixing system is interconnected with the physiological status of the plant host, and is determined by the influence of environmental factors, such as soil type, temperature, moisture, pH, salinity, micro and macronutrient content [19,30,31]. These factors can affect various aspects of symbiosis, including *Rhizobium* survival in the soil, infection process, legume development, nodule function and indirectly the growth of the host plant [32]. Due to symbiosis, legumes cover their nutritional needs in relation to N and therefore have lower requirements for mineral fertilization, which is limited to the necessary minimum, generally a small pre-sowing (starter) dose. This allows deficiencies of the nutrient to be made up in the initial period of plant growth and development, when the BNF process has not yet begun [33]. Individual legume species can only coexist with a specific bacterial species, for example beans (*Phaseolus vulgaris* L.) have the ability to symbiosis with *Rhizobium leguminosarum*

bv. *phaseoli*, and soybean (*Glycine max* (L.) Merr.) with *Bradyrhizobium japonicum* (Table 1). Bacteria coexisting with legumes do not always occur naturally in the soils in which these species are grown, so it is sometimes necessary to inoculate legume seeds prior to sowing with appropriately selected bacterial strains to enable rapid colonization of the rhizosphere and ensure effective nodulation, thereby improving the BNF process and maximizing yield [34–36]. However, the majority of soils used in agriculture contain *Rhizobium* populations that compete in the plant host nodulation process with strains introduced by inoculation.

The aim of this review is to summarize the current state of knowledge regarding the symbiosis between legumes and *Rhizobium* bacteria and to identify key environmental factors affecting the efficiency of biological nitrogen fixation (BNF). Particular focus is placed on the impact of factors such as soil salinity, acidity, temperature, moisture, light intensity, and the availability of macro- and micronutrients. This review seeks to highlight the challenges and opportunities associated with optimizing the symbiosis, which can enhance legume productivity and support sustainable agricultural practices.

Table 1. Examples of symbiotic interactions between bacteria and legume [37].

Type	Species	Host-legume
<i>Rhizobium</i>	<i>R. leguminosarum</i> bv. <i>viciae</i>	vetch, peas, lentils, chickling vetch
	<i>R. leguminosarum</i> bv. <i>phaseoli</i>	beans
	<i>R. leguminosarum</i> bv. <i>trifolii</i>	clover
	<i>R. leguminosarum</i> bv. <i>etli</i>	beans, common bean
<i>Bradyrhizobium</i>	<i>B. japonicum</i>	soybean
	<i>B. elkani</i>	common bean
<i>Sinorhizobium</i>	<i>S. meliloti</i>	melilot, alfalfa, fenugreek
	<i>S. fredii</i>	soybean, common bean
<i>Mezorhizobium</i>	<i>M. loti</i>	lotus, lupin
	<i>M. huakuii</i>	Chinese milk vetch

2. Soil Salinity

One of the factors that determine the physiological status of plants and the formation and functioning of legume with *Rhizobium* symbiosis is the presence of salts in the soil. The negative effect of soil salinity on plant growth and development is manifested in a decrease in the intensity of growth processes and BNF, due to a decrease in the photosynthetic activity [19]. At the same time, soil salinity can lead to a decrease in the population of soil microorganisms, distortion of their structure, as well as cause inhibition of root and root hair development and disappearance of the mucous layer on them. This, in turn, leads to a deterioration of bacterial survival and a decrease in the possibility of forming an infectious thread during nodulation, a decrease in the intensity of nodule respiration, and a decrease in the production of leghemoglobin [32].

In the course of scientific research in the variants of inoculation of soybean seeds with *B. japonicum* in the presence of 170 mm NaCl, a slight twisting or deformation of root hairs was observed, while in the case of increasing the NaCl concentration to 210 mm, nodules on soybean roots were not formed. At the same time, studies have shown different effects of salt on the development and vital activity of bacteria of the *Rhizobium* and *Bradyrhizobium* strains. Thus, in the presence of 100 mm NaCl, a slowdown in the growth of *B. japonicum* was observed, while *R. meliloti* proved to be quite resistant to the effects of increased to 300-700 mm NaCl concentration [19].

The *Rhizobium* osmotolerance is mainly related to the presence of accumulated or unaccumulated osmoprotectants (betaine, amino acids and sugars) [38,39]. They are low-molecular-weight organic solutions and are characterized by the effect of counteracting dehydration caused by low water activity in the environment [40]. Some authors noted, that in some cases, the interaction of plant roots with bacteria characterized by high salt tolerance (which is manifested in the ability to grow in the range of 300-700 mM NaCl), leads to the formation of inefficient nodules with a low rate of nitrogen fixation [32].

3. Soil pH

A factor that significantly limits the cultivation of legumes in the world is soil acidity. In general, soils with an indifferent or weakly alkaline pH are favorable for legumes, while an acidic environment prevents most legume species from initiating the symbiosis process. Low soil pH is caused by an excess of acidic cations, such as H^+ , Al^{3+} , Mn^{2+} compared to alkaline cations, such as Ca^{2+} , Mg^{2+} , K^+ and Na^+ . Its negative impact is reflected in the damage to the root system of plants and the corresponding weakening of the absorption and assimilation of water and nutrients dissolved in it, resulting in yield losses in legumes of up to 50% [41].

In acidic soils, the release of carbon compounds by the plant root system, which are substrates for soil microorganisms living in the rhizosphere, is significantly reduced. Due to a significant limitation of rhizobia survival, as well as a negative impact on nodule formation in acidic soils, BNF is significantly reduced [42]. The negative effect of soil acidity is manifested already in the early stages of the infection process, namely during the exchange of signals between the host plant and the microsymbiont, which negatively affects the attachment of rhizobia to root hairs and, accordingly, leads to a decrease in bacterial colonization of roots [19]. Ferguson et al. [41] attributed this effect to a decrease in flavonoid secretion into the rhizosphere by the root system of legumes, which leads to a decrease in the induction of the Nod gene by Rhizobium, inhibition of nod factor secretion and excretion of Nod metabolite. In turn, the decrease in nod factor signaling leads to deformation of root hairs and their twisting. At the same time, cell division and nodule primordial tissue formation are inhibited [43].

4. Soil Temperature

An important environmental factor that determines the diversity of bacterial communities, as well as the formation of plant-bacteria relationships, is soil temperature [44]. Numerous studies have shown that soil temperatures that are higher or lower than the optimum, have a negative impact on the relationship between both partners of the symbiosis, in general. The optimal range of soil temperature for the functioning of nodule bacteria is between 28 and 31°C. However, some strains are able to survive an increase in soil temperature up to 35-40°C [45]. The range of temperature optimum for legume-Rhizobium symbiosis is determined by the biological characteristics of both partners [46]. The results of Michiels et al. [47] showed, that the optimal soil temperature for BNF in soybean and peanut is 30°C, and the critical temperature is in the range of 35-40°C. The most favorable temperature regime for the root zone for BNF of common beans ranges from 25 to 30°C. Increasing the soil temperature to 30-33°C inhibits legume-Rhizobium symbiosis, which is manifested in a smaller size and weight of nodules, as well as in a decrease in nitrogenase activity [19]. Ramires and Damo [48] reported, that an increase in temperature in the root zone is critical for BNF, due to the deterioration of the physiological state of rhizobium, reducing their ability to infect root hairs, form infectious threads, initiate nodule formation and growth, reduce the content of leghemoglobin in nodules, and reduce the activity of the nitrogenase complex.

On the other hand, low temperatures may limit the survival of Rhizobium, as well as the secretion of flavonoids by legumes, hindering the exchange of specific signaling molecules between them, which ultimately leads to a reduction in the number of nodules and adversely affects legume growth [49]. According to various studies, the effect of cold stress is manifested in damage to bacterial cell membranes [50] and stabilization of secondary RNA/DNA structures, which in turn leads to a decrease in the efficiency of the main stages of the implementation of genetic information (translation, transcription and replication) [51]. At the same time, signalization between both partners of the legume-Rhizobium symbiosis is disrupted, the period of nodule initiation and formation is prolonged [52], and their size is reduced [53].

5. Soil Moisture Content

The moisture availability of plants is a key environmental factor that determines the conditions of vital activity not only microbial communities, but also plants. It plays a key role in biological processes occurring in plants at all stages of development [54]. Álvarez-Aragóon et al. [55] report, that the aboveground and underground parts of plants respond to drought conditions in different ways. Thus, if the aboveground part undergoes significant modifications aimed at reducing moisture loss and optimizing metabolism, changes in the root system contribute to better moisture supply to plants [56]. Dehydration of plant cells and tissues caused by water deficit leads to a decrease in

photosynthetic activity, efficiency of moisture use by plants, absorption and transport of nutrients from roots to shoots, inhibition of synthesis of organic compounds as well as changes in the hormonal balance of plants [57]. Morphological manifestation of these changes is a decrease in root proliferation, leaf blade size and internode length, which consequently leads to a decrease in plant productivity [58].

Drought affects the initiation of symbiosis, because water deficit inhibits the exchange of signaling molecules involved in communication between the legume and Rhizobium, resulting in poor nodulation and reduced BNF [59–61]. Furlan et al. [62] notes, that insufficient moisture supply of plants negatively affects the density of populations of nitrogen-fixing microorganisms and leads to disruption of the process of infection of root hairs, due to inhibition of the formation of infectious threads [63,64]. The main physiological responses to water stress in legume-Rhizobium systems are a reduction in the intensity of carbon metabolism in nodules, reduced nitrogenase activity due to oxygen limitation, regulation of feedback by accumulation of nitrogen fixation products, and modification of Rhizobium cells [19,65–68].

Numerous studies confirm the reduction of legume nodulation under drought stress [69,70]. Michałek [71] showed a reduction in the number, weight, and activity of nodules formed on the roots of soybean plants at the flowering stage under the influence of periodic drought, which had a significant impact on plant productivity. Soil water content of 50% and 25% FWC reduced, compared to the control, the number of nodules by 22 and 42%, root fresh weight by 46 and 69%, and root dry weight by 23 and 41%, respectively. Also, Sadeghipour and Abbasi [72] showed a decrease in symbiotic bacteria activity and a decrease in the number of pods per plant, seeds per pod, seed weight, and seed yield under drought stress conditions.

The nodulation process is adversely affected by both deficiency and excess of water. Increased soil moisture stimulates anaerobic processes, that lead to the appearance of phytotoxins, compounds produced by anaerobic microorganisms in the soil. These chemical compounds can negatively affect plant roots, restricting their growth, decreasing nutrient transpiration, damaging roots and impairing the ability to form root nodules. These compounds are also not indifferent to Rhizobium populations [73]. Particularly sensitive to fluctuations in soil moisture and oxygen content are nitrogenases, whose synthesis is inhibited in an environment with a high oxygen content. Higher soil moisture creates a low-oxygen environment for N₂-fixing bacteria, which may favor BNF productivity [74].

6. Light

Light is the main source of energy, that drives the process of photosynthesis, during which plants convert light energy into energy of chemical bonds contained in the sugar and other organic compounds [75].

Organic compounds synthesized during photosynthesis act as a primary source of carbon for plants, and its excess is released into the rhizosphere in the form of root exudates [76]. Sugars, organic acids, amino acids, and secondary metabolites released into the rhizosphere play an important role in providing microorganisms with carbon and energy, thus determining the number of rhizosphere communities and establishing mutualistic relationships between legume plants and microorganisms [77]. At the same time, photosynthetic products act as a source of energy for legume-Rhizobium symbiosis. According to Lepetit and Brouquisse [29], more than 25% of photosynthetic products are used by plants to support the functioning of BNF.

7. Nutrient's Content in the Soil

An important factor determining the efficiency of legume-Rhizobium symbiosis is the presence of mineral nutrients in the soil [44].

Nitrogen (N) is one of the essential macronutrients that determines the growth and development of the vegetative part of plants. This element is required for numerous processes of synthesis of organic compounds, including proteins (de Bruijn, 2015 [78], chlorophyll, nucleic acids and hormones [42,79]. Plants can absorb it from mineralized soil organic matter [80], industrial nitrogen fertilizers [81], and in the BNF process. Abd-Alla et al. [82] note, that if there is a sufficient amount of inorganic (mineral) nitrogen (NO₃⁻) in the soil, plants absorb nitrate from the soil as non-symbiotic higher plants. Scientists attribute this effect to a reduction in energy consumption by plants for the N₂ to NH₃ conversion by the nitrogenase complex in nodules. It is known, that for its transformation, the plant

uses organic compounds synthesized during photosynthesis. Their amount is equivalent to 25% of shoot dry matter at harvest. Therefore, in order to reduce these costs, if other alternative sources of N are available, legumes will use them to ensure their vital activity. Under conditions of mineral N deficiency in the soil, plants maintain BNF in nodules at the lowest level necessary to ensure growth processes [82].

Mineral N has an inhibitory effect on the interaction of legume-Rhizobium symbiosis partners [83]. This is indicated by a decrease in the colonization of root hairs by N-fixing microorganisms [66], weakening of root hairs twisting [84], reduction of the number of infectious threads and associated inhibition of nodule development and disruption of the BNF process [85,86]. In addition, the presence of mineral N causes early nodule senescence and reverse inhibition of nitrogenase activity [87]. In legumes, the mineral N content of the soil is particularly important at the beginning of vegetation, before root nodules appear and bacterial symbiosis develops. From the budding stage onwards, when the assimilation of N₂ from the air by Rhizobium begins, legumes are self-sufficient in this nutrient. Therefore, N deficiency may be related to disruptions in nodules establishment, their function or N transport to vegetative parts. A study by Salvagiotti et al. [80] showed, that about 80 kg N ha⁻¹ is needed to produce 1 t ha⁻¹ of soybean seed. This requirement is met 50-80% by the plant through the BNF pathway [88]. Therefore, if higher yields with satisfactory seed quality are desired, soil N and BNF may not be sufficient to meet the needs of the plant during the seed-filling period to realize the yield potential of soybean.

Phosphorus (P) plays a key role in the energy supply of physiological processes of plant life, in particular the BNF process [89]. This element is essential for nodules growth and proper symbiosis with Rhizobium bacteria. Root nodules typically contain two to three times more P than other plant organs [90,91]. When plants are sufficiently supplied with P, photosynthetic products are used to support both root development and nodule function [92,93]. Under P deficiency, on the contrary, most of the synthesized organic compounds are directed to the roots rather than to the nodules. Insufficient supply of P to plants leads to inhibition of the nodulation process [94,95], nitrogenase activity, reduction of nodule leghemoglobin content [96], decrease of active N₂-fixation in nodules [26] and relevant deterioration of BNF. Numerous studies confirm that P influences earlier formation of active root nodules, increases their size and number, and improves the amount of bioavailable N per unit weight [97,98], in addition to increasing root and nodules weight [99,100].

Potassium (K) is an important element that determines the adaptive capacity of plants to abiotic stresses. K acts as an activator of more than 60 enzyme systems that catalyze numerous metabolic reactions in plants. It maintains the osmotic status and turgor of cells, regulates their cation-anion balance and cytoplasmic pH, controls membrane polarization, cell expansion, and stomatal movements, thus regulating the supply of CO₂ and moisture to the plant [101]. In legumes, K, as the most abundant intracellular cation, plays an important role in the formation of a powerful root system and its moisture absorption, root hair growth, which in turn improves nodulation and BNF [44]. At the same time, its presence provides support for the turgor pressure of bacterial cells, pH regulation, gene expression, and activation of cellular enzymes [102].

Sulfur (S) is an element that has an important function in nitrogen management and is involved in the biosynthesis and function of molybdenum-containing enzyme structures (nitrogenase enzyme complex). As a component of the amino acids Cys and Met, metal cofactors, coenzymes and secondary metabolites, takes part in the processes of synthesis of proteins and chlorophyll, nutrient absorption and nitrogen use efficiency by plants during their growth and development, as well as plant resistance to various stress factors [103]. Its deficiency negatively affects the formation of legume-Rhizobium symbiosis, which is associated with inhibition of nodulation and BNF, due to a decrease in the intensity of biosynthesis and the activity of nitrogenase, that contains an increased amount of S in the form of FeS clusters [104]. Some authors attribute this to a decrease in the availability of Cys and Met amino acids and a general change in nodule metabolism. The reduction in S supply to plants leads to a limited supply of energy and carbon skeleton to the legume-Rhizobium symbiosis, due to a decrease in the content of leghemoglobin and glucose in nodules, ATP in mitochondria and bacteroids, and ferredoxin in bacteroids [105–107]. Ferredoxin and nitrogenase are crucial for BNF process, as they contain the iron-sulphur and molybdenum-iron-sulphur metal clusters [108]. S can affect BNF by modulating the growth and function of root nodules or by

influencing the growth of the host plant. S deficiency reduces BNF efficiency in legumes by decreasing ferredoxin and leghemoglobin concentrations and ATP supply [109].

Calcium (Ca) influences the proper development of the root system and the aboveground parts of the plant. Through adequate root growth, it increases the likelihood of nodules forming and their number on the roots. In the symbiotic association legume and Rhizobium, Ca plays a crucial role in signaling between plants and bacteria and recognition of rhizobial nodal factors in the early stages of symbiotic interaction [44]. Miwa et al. [110] associate this with an increase in the level of flavonoids in plant root exudates, which in turn enhances the induction of nodal genes through signal perception by Rhizobium. Ca enhances the colonization of the legume root system by Rhizobium due to an increased release of bacterial exopolysaccharides and their gel formation as a means of attaching Rhizobium to root hairs [42]. At the same time, when plants are sufficiently supplied with this element, an increase in the activity of Nod genes is observed [111].

Magnesium (Mg) plays a significant role in the growth processes of both the legumes and the Rhizobium. At the beginning of the vegetation period, this element is essential for proper growth of the root system, which later determines adequate water and nutrient uptake. In the plant, the importance of Mg for energy metabolism, photosynthesis, protein synthesis, carbon allocation, and stress resistance through the Mg transporter is well known [112]. Mg also has a beneficial effect on the metabolism and distribution of N in legumes, which translates into planning levels and seed protein content. Mg affects nodule formation by regulating C-N transport in nodules and by changing the plasmadermal permeability of internal cortical cells in nodules [112–114].

Iron (Fe) is a key component of the nitrogenase complex, which is a MoFe protein, as well as a cofactor of other proteins that are important for N₂ fixation, such as leghemoglobin, cytochrome or ferredoxin [115]. In plants, Fe deficiency leads to a decrease in chlorophyll synthesis and inhibition of photosynthesis, as well as causes chlorosis and serious growth disorders of plants [116]. Excess iron is harmful and can negatively affect the uptake of manganese by plants.

Molybdenum (Mo), which is a component of enzymes, plays an important role in key metabolic processes of plant, in particular, photosynthesis and protein synthesis [117]. Mo, as a constituent part of the FeMoCo nitrogenase complex, plays a key role in the process of nitrogen reduction [118]. Therefore, the amount of this element in the nodules is about 10 times higher than in other parts of the plant [119]. Mo stimulates the formation of new root nodules and N₂ fixation and influences the number of flowers and pods on the plant, which has a direct effect on seeds yield. Deficiency of the Mo in the soil leads to disorganization of biochemical and physiological reactions, which weakens growth of plants, reduces the seeds yield and deteriorates its quality indicators [120]. A decrease in molecular nitrogen fixation caused by Mo deficiency leads to N deficiency in legumes [121]. Mo is also involved in the metabolism of P, S and Fe. A water deficit results in a reduction of BNF and, when Mo content in the soil is low, it can reduce the amount of N transformed up to several kg ha⁻¹ [122].

Boron (B) is an essential component of cell walls, thus increasing stem resistance to cracking and secondary pathogen infestation. B is involved in reproduction, phenolic metabolism and N₂ fixation. B plays a key role at every stage of nodules development and morphogenesis. Redondo-Nieto et al. [123] point out, that it is important for maintaining the structure of Rhizobium cell membranes and for the exchange of signals between plants and bacteria. At the same time, its role in the processes of Rhizobium infection of legume roots, nodule invasion and bacteroid development, is well known. Deficiency of B reduces the formation and growth of root nodules, resulting in N deficiency, weaker plant growth and reduced plant flowering time.

Nickel (Ni) is an activator of the urease enzyme, which facilitates the hydrolysis of urea to produce NH₃ [124]. As an integral part of the hydrogenase, nickel take part in the process of deactivation of hydrogen gas, which is a by-product of BNF. Utilization of gaseous hydrogen increases the energy efficiency of N₂ fixation by producing more ATP and simultaneously reducing energy use. At the same time, this process mitigates the inhibitory effect of gaseous hydrogen and oxygen on nitrogenase activity [125].

Cobalt (Co) plays an important role in the development, morphological changes, and vital functions of Rhizobium, as well as in the differentiation of bacteroids [125]. This element is a constituent of cobalamin, which is a cofactor of Rhizobium enzymes, that act as catalysts for the conversion of N₂ to NH₃ [126]. Research results indicate a significant role of Co fertilization in

increasing the efficiency of legume-Rhizobium symbiosis, as well as in increasing the level of N content in legume tissues [125]. At the same time, some authors report a decrease in the concentration of reactive oxygen species in root nodules, which improves the growth conditions of Rhizobium [127].

Zinc (Zn) is an element involved in carrying out efficient and effective photosynthesis and is involved in the reduction of free radicals, which are formed during the action of various stress factors. It acts as a catalytic cofactor and structural element in a large number of proteins [128]. Its deficiency leads to significant changes in metabolism, which is expressed in the inhibition of growth processes and defects in the formation of the aboveground part of plants, in particular, chlorosis between the fibres, the appearance of necrotic areas on the leaves, reduction of internode length and leaf size, bending of the leaf blades [129] and reduce of nitrogenase activity [130]. Zn also has a yield-forming effect by stimulating the biosynthesis of plant hormones, including auxins, whose deficiency causes plant growth inhibition. Weisany et al. [131] noted, that Zn fertilization of plants growing on saline soils plants increases the intensity of photosynthesis and the efficiency of soil moisture utilization by plants.

Manganese (Mn) plays a crucial role in a variety of physiological and biochemical processes in plant life. It is an integral part of the more than 35 key enzymes in photosynthesis, respiration, nitrogen metabolism, and the processes of proteins, lipids, carbohydrates and chlorophyll synthesis [125,132]. In many enzymes, Mn is interchangeable with Ca, Co, Cu, Mg or Zn. Its importance in the process of assimilation of other mineral nutrients (Fe, Ca, Mg) by plants is widely known. In addition, it is involved in the conversion of nitrate (NO_3^-) to ammonia (NH_3) [125]. As a cofactor of the antioxidant enzyme superoxide dismutase, it has a protective effect on root nodules under the influence of oxidative stress caused by reactive oxygen species [133]. To ensure normal functioning, plants need only a small amount of this trace element, but its deficiency can lead to delayed root growth and reduced nutrient uptake, disruption of hormonal signaling, reduced immune response to adverse abiotic factors, and pathogen damage [134].

Copper (Cu) as a cofactor of enzymes plays a key role in photosynthesis, respiration, supply and distribution of photoassimilates between plant organs, ethylene signaling, regulation of free radical levels [135,136] and energy transfer to bacteroids [125]. Cu is a component of enzymes associated with free radical metabolism, respiration, Rhizobium vital functions in root nodules [125].

Numerous studies show that Cu and Mn reduce the activity of Rhizobium interfere with the formation of new nodules and inhibit bacterial activity in already formed nodules [137–139]. Cu has also been shown to have bactericidal effects [140]. These results relate to the detrimental effect of Cu and Mn on Rhizobium bacteria with which legume seeds have been treated in the form of sulphate or chloride [137–141]. These micronutrients already inhibit bacterial activity when applied to inoculated seeds [36]. The effect of these micronutrients in chelated form on the activity of symbiotic bacteria has not yet been investigated.

8. Conclusions

The review highlights the multifaceted role of legumes in sustainable agriculture and their ability to mitigate environmental challenges while providing high-quality protein sources. The symbiosis between legumes and Rhizobium bacteria is crucial for biological nitrogen fixation (BNF), contributing significantly to soil fertility and reducing the dependency on synthetic nitrogen fertilizers. However, the efficiency of this process is highly influenced by environmental factors, including soil salinity, acidity, temperature, moisture, light availability, and nutrient content. Addressing these challenges requires a comprehensive understanding of the mechanisms underlying legume-rhizobium interactions and their response to abiotic stresses. Innovations in seed inoculation, soil management practices, and nutrient supplementation can improve BNF efficiency and crop yields. The economic and environmental benefits of legumes underscore their importance in sustainable agricultural systems, where they enhance soil health, support biodiversity, and reduce greenhouse gas emissions.

Future research should focus on breeding stress-tolerant legume varieties and developing advanced inoculants to maximize BNF under diverse environmental conditions. This holistic approach will enable the full utilization of the potential of legumes, contributing to global food security and environmental sustainability.

Author Contributions: “Conceptualization, L.Y.; V.H.; M.M.; methodology, L.Y.; V.H.; resources, L.Y.; M.S.; K.C.; writing—original draft preparation, L.Y.; M.S.; K.C.; writing, L.Y.; M.S.; K.C.; visualization, M.S.; supervision, L.Y.; M.S.; K.C. funding acquisition, M.S. All authors have read and agreed to the published version of the manuscript.”

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Singh, P.; Krishnaswamy, K. Sustainable zero-waste processing system for soybeans and soy by-product valorization. *Trends Food Sci. Technol.* **2022**, *128*, 331–344. <https://doi.org/10.1016/j.tifs.2022.08.015>
2. Çakor, Ö.; Uçarlıo, C.; Tarhan, Ç.; Pekmez, M.; Tyrgut-Kara, N. Nutritional and health benefits of legumes and their distinctive genomic properties. *Food Sci. Technol.* **2019**, *39*, 1–12. <https://doi.org/10.1590/fst.42117>
3. Ekholuenetale, M.; Tudeme, G.; Onikan, A.; Ekholuenetale, C.E. Socioeconomic inequalities in hidden hunger, undernutrition, and overweight among under-five children in 35 sub-Saharan Africa countries. *J. Egypt Public Health Assoc.* **2020**, *95*, 9. <https://doi.org/10.1186/s42506-019-0034-5>
4. Ibeanu, V.N.; Edeh, C.G.; Ani, P.N. Evidence-based strategy for prevention of hidden hunger among adolescents in a suburb of Nigeria. *BMC Public Health.* **2020**, *20*, 1683. <https://doi.org/10.1186/s12889-020-09729-8>
5. Marventano, S.; Pulido, M. I.; Sánchez-González, C.; Godos, J.; Speciani, A.; Galvano, F.; Grosso, G. Legume consumption and CVD risk: A systematic review and meta-analysis. *Public Health Nutrition.* **2017**, *20*, 245–254. <https://doi.org/10.1017/S1368980016002299>
6. Jach, M.E.; Sajnaga, E.; Ziaja, M. Utilization of legume-nodule bacterial symbiosis in phytoremediation of heavy metal-contaminated soils. *Biology.* **2022**, *11*, 676. <https://doi.org/10.3390/biology11050676>
7. Mejia, E.G.; Dia, V.P. The role of nutraceutical proteins and peptides in apoptosis, angiogenesis, and metastasis of cancer cells. *Cancer and Metastasis Reviews.* **2010**, *29*, 511–528. <http://dx.doi.org/10.1007/s10555-010-9241-4>
8. Möller, N.P.; Scholz-Ahrens K.E.; Roos, N.; Schrezenmeir, J. Bioactive peptides and proteins from foods: indication for health effects. *Eur. J. Nutr.* **2008**, *47*, 171–182. <http://dx.doi.org/10.1007/s00394-008-0710-2>
9. Zambrowicz, A.; Timmer, M.; Polanowski, A.; Lubec, G.; Trziszka, T. Manufacturing of peptides exhibiting biological activity. *J. Amino Acids.* **2013**, *44*, 315–320. <http://dx.doi.org/10.1007/s00726-012-1379-7>
10. Duranti, M. Grain legume proteins and nutraceutical properties. *Fitoterapia.* **2006**, *77*, 67–82. <https://doi.org/10.1016/j.fitote.2005.11.008>
11. Arnoldi, A.; Boschini, G.; Zannoni, C.; Lammi, C. The health benefits of sweet lupin seed flours and isolated proteins. *J. Funct. Foods.* **2015**, *18*, 550–563. <http://dx.doi.org/10.1016/j.jff.2015.08.012>
12. Sońta, M.; Rekiel, A. Legumes – use for nutritional and feeding purposes. *J. Elementol.* **2020**, *25*, 835–849. <http://dx.doi.org/10.5601/jelem.2020.25.1.1953>
13. Voisin, A.S.; Guéguen, J.; Huyghe, C.; Jeuffroy, M.H.; Magrini, M.B.; Meynard, J.M.; Mougél, C.; Pellerin, S.; Pelzer, E. Legumes for feed, food, biomaterials and bioenergy in Europe: A review. *Agron. Sustain. Dev.* **2014**, *34*, 361–380. <http://dx.doi.org/10.1007/s13593-013-0189-y>
14. Meena, R.S.; Meena, V.S.; Meena, S.K.; Verma, J.P. The needs of healthy soils for a healthy world. *Journal of Cleaner Production.* **2015**, *102*, 560–561. <https://doi.org/10.1016/j.jclepro.2015.04.045>
15. Deakin, W.J.; Broughton, W.J. Symbiotic use of pathogenic strategies: rhizobial protein secretion systems. *Nat. Rev. Microbiol.* **2009**, *7*, 312–320. <https://doi.org/10.1038/nrmicro2091>
16. Stagnari, F.; Maggio, A.; Galiani, A.; Pisante, M. Multiple benefits of legumes for agriculture sustainability: an overview. *Chem. Biol. Technol. Agric.* **2017**, *4*, 2. <https://doi.org/10.1186/s40538-016-0085-1>
17. Kocira, A.; Staniak, M.; Tomaszewska, M.; Kornas, R.; Cymerman, J.; Panasiewicz, K.; Lipińska, H. Legume cover crops as one of the elements of strategic weed management and soil quality improvement. A review. *Agriculture.* **2020**, *10*, 394. <https://doi.org/10.3390/agriculture10090394>
18. Kebede, E. Contribution, utilization, and improvement of legumes-driven biological nitrogen fixation in agricultural systems. *Front. Sustain. Food Syst.* **2021**, *5*, 767998. <https://doi.org/10.3389/fsufs.2021.767998>
19. Zahran, H.H. *Rhizobium*-legume symbiosis and nitrogen fixation under severe conditions and in an arid climate. *Microbiol. Mol. Biol. Rev.* **1999**, *63*, 969–989. <https://doi.org/10.1128/mmr.63.4.968-989.1999>
20. Martínez-Espinosa, R.M.; Cole, J.A.; Richardson, D.J.; Wartmough, N.J. Enzymology and ecology of the nitrogen cycle. *Biochem. Soc. Trans.* **2011**, *39*, 175–178.
21. Podleśna, A. The process of N₂ fixation by leguminous plants as a source of nitrogen for cultivated crops. *Studia i Raporty IUNG-PIB*, **2018**, *56(10)*, 71–85. doi: 10.26114/sir.iung.2018.56.06 [in Polish]
22. Wielbo, J.; Skorupska, A. Ewolucja układu symbiotycznego *Rhizobium* – rośliny motylkowate. *Postępy Mikrobiologii* **2003**, *42(3)*, 263–283.

23. Oldroyd, G.E.; Dixon, R. Biotechnological solutions to the nitrogen problem. *Curr. Opin. Biotechnol.* **2014**, *26*, 19–24.
24. Krug, E.C.; Winstanley, D. The need for comprehensive and consistent treatment of the nitrogen cycle in nitrogen cycling mass balance studies: I. Terrestrial nitrogen cycle. *Sci. Total Environ.* **2002**, *293*, 1–29. DOI: 10.1016/S0048-9697(01)01133-0
25. Oleńska, E. Asocjacja symbiotyczna roślin bobowatych z ryzobiami. In *Różnorodność biologiczna – od komórki do ekosystemu. Funkcjonowanie roślin i grzybów. Środowisko-Eksperyment-Edukacja*. Bajguz, A., Ciereszko, I.; Publisher: Polskie Towarzystwo Botaniczne, 2015; p. 187–200 [in Polish].
26. Hirsch, A.M. Development biology of legume nodulation. *New Phytol.* **1992**, *122*, 211–237. <https://doi.org/10.1111/j.1469-8137.1992.tb04227.x>
27. Jensen, E.S.; Peoples, M.B.; Boddey, R.M.; Gresshoff, P.M.; Hauggaard-Nielsen, H.; Alves, B.J.R.; Morrison, M.J. Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. *Agron. Sustain. Dev.* **2012**, *32*, 329–364. DOI 10.1007/s13593-011-0056-7
28. Martyniuk, S. The importance of biological fixation of atmospheric nitrogen in ecological agriculture. *Int. J. Res. Appl. Sci. Eng. Technol.*, **2008**, *53(4)*, 9–14.
29. Lepetit, M.; Brouquisse, R. Control of the rhizobium–legume symbiosis by the plant nitrogen demand is tightly integrated at the whole plant level and requires interorgan systemic signaling. *Front. Plant Sci.* **2023**, *14*, 1114840.
30. Abd-Alla, M.H.; Bagy, M.K.; El-enany, A.W.E.S.; Bashandy, S.R. Activation of *Rhizobium tibeticum* with flavonoids enhances nodulation, nitrogen fixation, and growth of fenugreek (*Trigonella foenum-graecum* L.) grown in cobalt-polluted soil. *Arch. Environ. Contam. Toxicol.* **2014**, *66*, 303–315. DOI: 10.1007/s00244-013-9980-7
31. Abd-Alla, M.H.; Nafady, N.A.; Bashandy, S.R.; Hassan, A.A. Mitigation of effect of salt stress on the nodulation, nitrogen fixation and growth of chickpea (*Cicer arietinum* L.) by triple microbial inoculation. *Rhizosphere* **2019**, *10*, 100148. DOI: 10.1016/j.rhisph.2019.100148
32. Abd-Alla, M.H.; Abdel Wahab, A.M. Survival of *Rhizobium leguminosarum* biovar *viciae* subjected to heat, drought and salinity in soil. *Biol. Plant.* **1995**, *37*, 131–137. <https://doi.org/10.1007/BF02913008>
33. Kocoń, A. Indicators of symbiotic N₂ fixation depending on the varied nitrogen nutrition of faba beans. *Fragm. Agron.* **1999**, *2(16)*, 50–61. [in Polish]
34. Catroux, G.; Hartmann, A.; Revellin, C. Trends in rhizobial inoculant production and use. *Plant Soil*, **2001**, *230*, 21–30. DOI: 10.1023/A:1004777115628
35. Deaker, R.; Roughley, R. J.; Kennedy, I. R. Legume seed inoculation technology – A review. *Soil Biol. Biochem.* **2004**, *36(8)*, 1275–1288. doi: <https://doi.org/10.1016/j.soilbio.2004.04.009>
36. Brzezińska, A.; Mrozek-Niecko, A. Effect of selected micronutrient seed fertilizers on the viability of *Bradyrhizobium japonicum*. *Prog. Plant Prot.* **2021**, *61(1)*, 17–23.
37. Sujkowska, M. The course of the infection process in the symbiotic system of leguminous plants - *Rhizobium*. *Wiadomości Botaniczne* **2009**, *53(1/2)*, 35–53. [in Polish]
38. Bernard, T.; Pocard, J. A.; Perroud, B.; Le Rudulier, D. Variations in the response of salt-stressed *Rhizobium* strains to betaines. *Arch. Microbiol.* **1986**, *143*, 359–364. <https://doi.org/10.1007/BF00412803>
39. Gouffi, K.; Pica, N.; Pichereau, V.; Blanco, B. Disaccharides as a new class of nonaccumulated osmoprotectants for *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.* **1999**, *65*, 1491–1500. <https://doi.org/10.1128/AEM.65.4.1491-1500.1999>
40. Pereira, S.I. A.; Lima A.I.G.; Figueira, E.M.A.P. Heavy metal toxicity in *Rhizobium leguminosarum* biovar *viciae* isolated from soils subjected to different sources of heavy metal contamination: effect on protein expression. *Appl. Soil Ecol.* **2006**, *33*, 286–293. <https://doi.org/10.1016/j.apsoil.2005.10.002>
41. Ferguson, B.J.; Lin, M.-H.; Gresshoff, P.M. Regulation of legume nodulation by acidic growth conditions. *Plant Signal. Behav.* **2013**, *8*, e23426. <https://doi.org/10.4161/psb.23426>
42. Dabessa, A.; Abebe, Z.; Bekele, S. Limitations and strategies to enhance biological nitrogen fixation in sub-humid tropics of Western Ethiopia. *J. Agric. Biotech. Sustain. Dev.* **2018**, *10*, 122–131. <https://doi.org/10.5897/JABSD2018.0318>
43. Hawkins, J.P.; Oresnik, I.J. The rhizobium-legume symbiosis: co-opting successful stress management. *Front. Plant Sci.* **2022**, *12*, 796045. <https://doi.org/10.3389/fpls.2021.796045>
44. Nakei, M.D.; Venkataramana, P.B.; Ndakidemi, P.A. Soybean-nodulating rhizobia: ecology, characterization, diversity, and growth promoting functions. *Front. sustain. food syst.* **2022**, *6*, 824444. <https://doi.org/10.3389/fsufs.2022.824444>
45. Lebrazi, S.; Fikri Benbrahim, K. Environmental stress conditions affecting the N₂ fixing rhizobium-legume symbiosis and adaptation mechanisms. *Afr. J. Microbiol. Res.* **2014**, *8*, 4053–4061. <http://dx.doi.org/10.5897/AJMR2014.7213>
46. Arayangkcon, T.; Schomberg, H.H.; Weaver, R.W. Nodulation and N₂ fixation of guar at high root temperature. *Plant Soil.* **1990**, *126*, 209–213. <https://doi.org/10.1007/BF00012824>

47. Michiels, J.; Verreth, C.; Vanderleyden, J. Effects of temperature stress on bean nodulating *Rhizobium* strains. *Appl. Environ. Microbiol.* **1994**, *60*, 601994. <https://doi.org/10.1128/aem.60.4.1206-1212.1994>
48. Ramírez, M.D.A.; Damo, J.L.C. Symbiosis under abiotic stress and its challenges. In *Symbiosis in nature*. IntechOpen. **2023**. <http://dx.doi.org/10.5772/intechopen.109972>
49. Duzan, H.M.; Zhou, X.; Souleimanov, A.; Smith, D.L. Perception of *Bradyrhizobium japonicum* Nod factor by soybean [*Glycine max* (L.) Merr.] root hairs under abiotic stress conditions. *J. Exp. Bot.* **2004**, *55*, 408, 2641–2646. doi:10.1093/jxb/erh265
50. Liu, Y.-S.; Geng, J.-C.; Sha, X.-Y.; Zhao, Y.-X.; Hu, T.-M.; Yang, P.-Z. Effect of rhizobium symbiosis on low-temperature tolerance and antioxidant response in alfalfa (*Medicago sativa* L.). *Frontiers in Plant Science*. **2019**, *10*, 538. <https://doi.org/10.3389/fpls.2019.00538>
51. Phadtare, S.; Inouye, M. The cold shock response. *EcoSal Plus*. **2008**, *3*, 33–45. <https://doi.org/10.1128/ecosalplus.5.4.2>
52. Pan, B.; Smith, O.L. Genistein and daidzein concentrations and contents in seedling roots of three soybean cultivars grown under three-foot zone temperatures. *J. Agron. Crop Sci.* **1998**, *180*, 77–82. <https://doi.org/10.1111/j.1439-037X.1998.tb00374.x>
53. de Lira Juniora, M.; Lima, A.S.T.; Arruda, J.R.F.; Smith, D.L. Effect of root temperature on nodule development of bean, lentil and pea. *Soil Biol. Biochem.* **2005**, *37*, 235–239. <https://doi.org/10.1016/j.soilbio.2004.07.032>
54. Chammakhi, C.; Boscari, A.; Pacoud, M.; Aubert, G.; Mhadhbi, H.; Brouquisse, R. Nitric oxide metabolic pathway in drought-stressed nodules of faba bean (*Vicia faba* L.). *Int. J. Mol. Sci.* **2022**, *23*, 13057. <https://doi.org/10.3390/ijms232113057>
55. Álvarez-Aragón, R.; Palacios, J.M.; Ramírez-Parra, E. Rhizobial symbiosis promotes drought tolerance in *Vicia sativa* and *Pisum sativum*. *Environ. Exp. Bot.* **2023**, *208*, 105268. <https://doi.org/10.1016/j.envexpbot.2023.105268>
56. Bogati, K.; Walczak, M. The impact of drought stress on soil microbial community, enzyme activities and plants. *Agronomy*. **2022**, *12*, 189. <https://doi.org/10.3390/agronomy12010189>
57. Chieb, M.; Gachomo, E.W. The role of plant growth promoting rhizobacteria in plant drought stress responses. *BMC Plant Biol.* **2023**, *23*, 407. <https://doi.org/10.1186/s12870-023-04403-8>
58. Flexas, J.; Barbour, M.M.; Brendel, O.; Cabrera, H.M.; Carriqui, M.; Díaz-Espejo, A.; Douthe, C.; Dreyer, E.; Ferrio, J.P.; Gago, J.; Gallé, A.; Galmés, J.; Kodama, N.; Medrano, H.; Niinemets, Ü.; Peguero-Pina, J.J.; Pou, A.; Ribas-Carbó, M.; Tomás, M.; Tosens, T.; Warren, C.R. Mesophyll diffusion conductance to CO₂: an unappreciated central player in photosynthesis. *Plant Sci.* **2012**, *193–194*, 70–84. <https://doi.org/10.1016/j.plantsci.2012.05.009>
59. Miransari, M.; Riahi, H.; Eftekhari, F.; Minaie, A.; Smith, D. L. Improving soybean (*Glycine max* L.) N₂ fixation under stress. *J. Plant Growth Regul.*, **2013**, *32*, 909–921. DOI 10.1007/s00344-013-9335-7
60. Thilakarathna, M.S.; Cope, K.R. Split-root assays for studying legume-rhizobia symbioses, rhizodeposition, and belowground nitrogen transfer in legumes. *J Exp Bot.* **2021**, *72*, 5285–5299. <https://doi.org/10.1093/jxb/erab198>
61. de Freitas, I.C.; Ferreira, E.A.; Alves, M.A.; de Oliveira, J.C.; Frazão, L.A. Growth, nodulation, production, and physiology of leguminous plants in integrated production systems. *Agrosyst Geosci Environ* **2023**, *6*, e20343. <https://doi.org/10.1002/agg2.20343>
62. Furlan, A.; Llanes, A.; Luna, V.; Castro, S. Physiological and biochemical responses to drought stress and subsequent rehydration in the symbiotic association peanut-*bradyrhizobium* sp. *Int. Sch. Res. Notices*. **2012**, 318083. <https://doi.org/10.5402/2012/318083>
63. Kasper, S.; Christoffersen, B.; Soti, P.; Racelis, A. Abiotic and biotic limitations to nodulation by leguminous cover crops in south Texas. *Agric.* **2019**, *9*, 209. <https://doi.org/10.3390/agriculture9100209>
64. Herrmann, L.; Chotte, J.L.; Thuita, M.; Lesueur, D. Effects of cropping systems, maize residues application and N fertilization on promiscuous soybean yields and diversity of native rhizobia in Central Kenya. *Pedobiologia (Jena)* **2014**, *57*, 75–85. <https://doi.org/10.1016/j.pedobi.2013.12.004>
65. McCulloch, L. A.; Piotto, D.; Porder, S. Drought and soil nutrients effects on symbiotic nitrogen fixation in seedlings from eight Neotropical legume species. *Biotropica*. **2021**, *53*, 703–713. <https://doi.org/10.1111/btp.12911>
66. Nishida, H.; Suzaki, T. Nitrate-mediated control of root nodule symbiosis. *Curr. Opin. Plant Biol.* **2018**, *44*, 129–136. <https://doi.org/10.1016/j.pbi.2018.04.006>
67. Zargar, S.M.; Gupta, N.; Nazir, M.; Mahajan, R.; Malik, F.A.; Sofi, N.R.; Shikari, A.B.; Salgotra, R.K. Impact of drought on photosynthesis: molecular perspective. *Plant Gene* **2017**, *11*, 154–159. <https://doi.org/10.1016/j.plgene.2017.04.003>
68. Thilakarathna, M.S.; Raizada, M.N. A meta-analysis of the effectiveness of diverse rhizobia inoculants on soybean traits under field conditions. *Soil Biol Biochem* **2017**, *105*, 177–196. <https://doi.org/10.1016/j.soilbio.2016.11.022>

69. Lumactud, R.A.; Dollete, D.; Liyanage, D.K.; Szczyglowski, K.; Hill, B.; Thilakarathna, M.S. The effect of drought stress on nodulation, plant growth, and nitrogen fixation in soybean during early plant growth. *J Agron Crop Sci* **2023**, *209*, 345–354. <https://doi.org/10.1111/jac.12627>
70. Marquez-Garcia, B.; Shaw, D.; Cooper, J.W.; Karpinska, B.; Quain, M.D.; Makgopa, E.M.; Kunert, K.; Foyer, C.H. Redox markers for drought-induced nodule senescence, a process occurring after drought-induced senescence of the lowest leaves in soybean (*Glycine max*). *Ann Bot.* **2015**, *116*, 497–510. <https://doi.org/10.1093/aob/mcv030>
71. Michalek, S. The impact of drought on nodulation, root growth, and shoot development in Polish soybean varieties. *Zesz. Probl. Postę. Nauk Rol.* **2004**, *496*, 267–274. [in Polish]
72. Sadeghipour, O.; Abbasi, S. Soybean Response to Drought and Seed Inoculation. *World Appl. Sci. J.* **2012**, *17*, 55–60.
73. Golińska, B.; Maźrzak, C. Competition between rhizobial strains nodulating leguminous plants. *Biotechnologia*, **1999**, *3*, 46, 106–121. [in Polish]
74. Robson, R. L.; Postgate, J. R. Oxygen and hydrogen in biological nitrogen fixation. *Annu. Rev. Microbiol.* **1980**, *34*, 183–207. <https://doi.org/10.1146/annurev.mi.34.100180.001151>
75. Pérez-Fernández, M.; De Lara-Del Rey, I.A.; Magadlela, A. Shining a light on symbiosis: N-fixing bacteria boost legume growth under varied light conditions. *Agriculture.* **2024**, *14*, 164. <https://doi.org/10.3390/agriculture14020164>
76. Kuzyakov, Y.; Domanski, G. Carbon input by plants into the soil. Review. *J. Plant Nutr. Soil Sci.* **2000**, *163*, 421–431. [https://doi.org/10.1002/1522-2624\(200008\)163:4<421::AID-JPLN421>3.0.CO;2-R](https://doi.org/10.1002/1522-2624(200008)163:4<421::AID-JPLN421>3.0.CO;2-R)
77. Haichar, F.E.Z.; Santaella, C.; Heulin, T.; Achouak, W. Root exudates mediated interactions belowground. *Soil Biology and Biochemistry.* **2014**, *77*, 69–80. <https://doi.org/10.1016/j.soilbio.2014.06.017>
78. de Bruijn, F.J. Biological nitrogen fixation. in Principles of plant-microbe interactions. In *Microbes for sustainable agriculture*. Cham: Springer International Publishing. Switzerland. **2015**, 215–224. https://doi.org/10.1007/978-3-319-08575-3_23
79. Saturno, D.F.; Cerezini, P.; Moreira da Silva, P.M.; Oliveira, A.B.; de Oliveira, M.C.N.; de Hungria, M.; Nogueira, M.A. Mineral nitrogen impairs the biological nitrogen fixation in soybean of determinate and indeterminate growth types. *J. Plant Nutr.* **2017**, *40*, 1690–1701. <https://doi.org/10.1080/01904167.2017.1310890>
80. Salvagiotti, F.; Cassman, K.G.; Specht, J.E.; Walters, D.T.; Weiss, A.; Dobermann, A. Nitrogen uptake, fixation and response to fertilizer N in soybeans: a review. *Field Crops Res.* **2008**, *108*, 1–13. <https://doi.org/10.1016/j.fcr.2008.03.001>
81. Ramakrishnan, V.V.G.A. Nitrogen sources and cycling in the ecosystem and its role in air, water and soil pollution: a critical review. *J. Pollut. Eff. Cont.* **2015**, *3*, 1–28. <https://doi.org/10.4172/2375-4397.1000136>
82. Abd-Alla, M.H.; Al-Amri, S.M.; El-Enany, A.-W.E. Enhancing rhizobium–legume symbiosis and reducing nitrogen fertilizer use are potential options for mitigating climate change. *Agriculture.* **2023**, *13*, 2092. <https://doi.org/10.3390/agriculture13112092>
83. Liu, Y.; Yin, X.; Xiao, J.; Tang, L.; Zheng, Y. Interactive influences of intercropping by nitrogen on flavonoid exudation and nodulation in faba bean. *Scientific Reports.* **2019**, *9*, 4818. <https://doi.org/10.1038/s41598-019-41146-9>
84. Ralston, E.J.; Imsande, J. Nodulation of hydroponically grown soybean plants and inhibition of nodule development by nitrate. *J. Exp. Bot.* **1983**, *34*, 1371–1378. <https://doi.org/10.1093/jxb/34.10.1371>
85. Ono, Y.; Fukasawa, M.; Sueyoshi, K.; Ohtake, N.; Sato, T.; Tanabata, S.; Toyota, R.; Higuchi, K.; Saito, A.; Ohyama, T. Application of nitrate, ammonium, or urea changes the concentrations of ureides, urea, amino acids and other metabolites in xylem sap and in the organs of soybean plants (*Glycine max* (L.) Merr.). *Int. J. Mol. Sci.* **2021**, *22*, 4573. <https://doi.org/10.3390/ijms22094573>
86. Li, S.; Wu, C.; Liu, H.; Lyu, X.; Xiao, F.; Zhao, S.; Ma, C.; Yan, C.; Liu, Z.; Li, H.; Wang, X.; Gong, Z. Systemic regulation of nodule structure and assimilated carbon distribution by nitrate in soybean. *Front. Plant Sci.* **2023**, *14*, 1101074. <https://doi.org/10.3389/fpls.2023.1101074>
87. Zhou, S.; Zhang, C.; Huang, Y.; Chen, H.; Yuan, S.; Zhou, X. Characteristics and research progress of legume nodule senescence. *Plants.* **2021**, *10*, 1103. <https://doi.org/10.3390/plants10061103>
88. Salvagiotti, F.; Specht, J.E.; Cassman, K.G.; Walters, D.T.; Weiss, A.; Dobermann, A. Growth and nitrogen fixation in high-yielding soybean: Impact of nitrogen fertilization. *J. Agron.* **2009**, *101*(4), 958–970. <https://doi.org/10.2134/agronj2008.0173x>
89. Nlu, Y.F.; Chai, R.S.; Jin, G.L.; Wang, H.; Tang, C.X.; Zhang, Y.S. Responses of root architecture development to low phosphorus availability: a review. *Ann. Bot.* **2013**, *112*, 391–408. <https://doi.org/10.1093/aob/mcs285>
90. Qin, L.; Zhao, J.; Tian, J.; Chen, L.; Sun, Z.; Guo, Y.; Lu, X.; Gu, M.; Xu, G.; Liao, H. The high-affinity phosphate transporter GmPT5 regulates phosphate transport to nodules and nodulation in soybean. *Plant Physiol.* **2012**, *159*, 1634–1643. <https://doi.org/10.1104/pp.112.199786>

91. Vauclare, P.; Bligny, R.; Gout, E.; Widmer, F. An overview of the metabolic differences between *Bradyrhizobium japonicum* 110 bacteria and differentiated bacteroids from soybean (*Glycine max*) root nodules: an in vitro ¹³C-and ³¹P-nuclear magnetic resonance spectroscopy study. *FEMS Microbiology Letters* **2013**, *343*, 49–56. <https://doi.org/10.1111/1574-6968.12124>
92. Chen, L.; Qin, L.; Zhou, L.; Li, X.; Chen, Z.; Sun, L.; Wang, W.; Lin, Z.; Zhao, J.; Yamaji, N.; Ma, J.F.; Gu, M.; Xu, G.; Liao, H.A. Nodule-localized phosphate transporter GmPT7 plays an important role in enhancing symbiotic N₂ fixation and yield in soybean. *New Phytol.* **2019**, *221*, 2013–2025. <https://doi.org/10.1111/nph.15541>
93. Pueyo, J.J.; Quiñones, M.A.; de la Peña T.C.; Fedorova, E.E.; Lucas, M.M. Nitrogen and phosphorus interplay in lupin root nodules and cluster roots. *Front. Plant Sci.* **2021**, *12*, 644218. <https://doi.org/10.3389/fpls.2021.644218>
94. Yang, Z.; Gao, Z.; Zhou, H.; He, Y.; Liu, Y.; Lai, Y.; Zheng, J.; Li, X.; Liao, H. GmPTF1 modifies root architecture responses to phosphate starvation primarily through regulating *GmEXPB2* expression in soybean. *Plant J.* **2021**, *107*, 525–543. <https://doi.org/10.1111/tpj.15307>
95. Zhong, Y.; Tian, J.; Li, X.; Liao, H. Cooperative interactions between nitrogen fixation and phosphorus nutrition in legumes. *New Phytologist.* **2023**, *237*, 734–745. <https://doi.org/10.1111/nph.18593>
96. Li, H.; Wang, L.; Zhang, Z.; Yang, A.; Liu, D. Effect of phosphorus supply levels on nodule nitrogen fixation and nitrogen accumulation in soybean (*Glycine max* L.). *Agronomy.* **2022**, *12*, 2802. <https://doi.org/10.3390/agronomy12112802>
97. Sulieman, S.; Tran, L.S.P. Phosphorus homeostasis in legume nodules as an adaptive strategy to phosphorus deficiency. *Plant Sci.* **2015**, *239*, 36–43. <https://doi.org/10.1016/j.plantsci.2015.06.018>
98. Korir, H.; Mungai, N.W.; Thuita, M.; Hamba, Y.; Masso, C. Co-inoculation effect of rhizobia and plant growth promoting rhizobacteria on common bean growth in a low phosphorus soil. *Front. Plant Sci.* **2017**, *8*, 141. <https://doi.org/10.3389/fpls.2017.00141>
99. Zhao, J.; Fu, J.; Liao, H.; He, Y.; Nian, H.; Hu, Y.; Qiu, L.; Dong, Y.; Yan, X. Characterization of root architecture in an applied core collection for phosphorus efficiency of soybean germplasm. *Chin. Sci. Bull.* **2004**, *49*, 1611–1620.
100. Kuang, R.; Liao, H.; Yan, X.L.; Dong, Y. Phosphorus and nitrogen interactions in field-grown soybean as related to genetic attributes of root morphological and nodular traits. *J. Integr. Plant Biol.* **2005**, *47*, 549–559. <https://doi.org/10.1111/j.1744-7909.2005.00072.x>
101. Yeremko, L.; Hanhur, V.; Staniak, M. Effect of mineral fertilization and seed inoculation with microbial preparation on seed and protein yield of pea (*Pisum sativum* L.). *Agronomy.* **2024**, *14*, 1004. <https://doi.org/10.3390/agronomy14051004>
102. Domínguez-Ferreras, A.; Muñoz, S.; Olivares, J.; Soto, M.J.; Sanjuan, J. Role of potassium uptake systems in *Sinorhizobium meliloty* osmoadaptation and symbiotic performance. *J. Bacteriol.* **2009**, *191*, 2133–2143. <https://doi.org/10.1128/jb.01567-08>
103. Saleem, M.; Elahi, E.; Gandahi, A.W.; Bhatti, S.M.; Saleem, M.B.; Ibrahim H.; Ali, M. Effect of sulphur application on growth, oil content and yield of sunflower. *Sarhad J. Agric.* **2019**, *35*, 1198–1203. <http://dx.doi.org/10.17582/journal.sja/2019/35.4.1198.1203>
104. Rawal V.K. Role of sulphur in legume crops. *J. Pharm. Innov.* **2022**, *11*, 2127–2131.
105. Becana, M.; Wienkoop, S.; Matamoros, M.A. Sulfur transport and metabolism in legume root nodules. *Front. Plant Sci.* **2018**, *9*, 1434. <https://doi.org/10.3389/fpls.2018.01434>
106. Speck, J.J.; James, E.K.; Sugawara, M.; Sadowsky, M.J.; Gyaneshwar, P. An alkane sulfonate monooxygenase is required for symbiotic nitrogen fixation by *Bradyrhizobium diazoefficiens* (syn. *Bradyrhizobium japonicum*) USDA110^T. *App. Environ. Microbiol.* **2019**, *85*, e01552-19. <https://doi.org/10.1128/AEM.01552-19>
107. Dawar, R.; Karan, S.; Bhardwaj, S.; Meena, D.K.; Padhan, S.R.; Reddy, K.S.; Bana, R.S. Role of sulphur fertilization in legume crops: a comprehensive review. *Int. J. Plant Sci.* **2023**, *35*, 718–727. <https://doi.org/10.9734/ijpss/2023/v35i214033>
108. Barczak, B. Sulfur as a nutrient shaping the yield and quality of selected cultivated crops.. *Rozprawy nr 144, Uniwersytet Technologiczno-Przyrodniczy, Bydgoszcz*, **2010**, pp. 131. [in Polish]
109. Pacyna, S.; Schulz, M.; Scherer, H.W. Influence of sulphur supply on glucose and ATP concentrations of inoculated broad beans (*Vicia faba minor* L.). *Biol. Fertil. Soils* **2006**, *42*, 324–329. <https://doi.org/10.1007/s00374-005-0030-0>
110. Miwa, H.; Sun, J.; Oldroyd, G.E.D.; Downie, J.A. Analysis of nod-factor-induced calcium signaling in root hairs of symbiotically defective mutants of *Lotus japonicus*. *Mol. plant-microb. interact.* **2006**, *19*, 914–923. <https://doi.org/10.1094/MPMI-19-0914>
111. Capoen, W.; Den Herder, J.; Sun, J.; Verplancke, C.; De Keyser, A.; De Rycke, R.; Goormachtig, S.; Oldroyd, G.; Holsters, M. Calcium spiking patterns and the role of the calcium/calmodulin-dependent kinase C_{CaMK} in lateral root base nodulation of *Sesbania rostrata*. *Plant Cell.* **2009**, *21*, 1526–1540. <https://doi.org/10.1105/tpc.109.066233>

112. Cao, H.-R.; Peng, W.-T.; Nie, M.-M.; Bai, S.; Chen, C.-Q.; Liu, Q.; Guo, Z.-L.; Liao, H.; Chen, Z.-H. Carbon-nitrogen trading in symbiotic nodules depends on magnesium import. *Curr. Biol.* **2022**, *32*, 4337–4349.e5. <https://doi.org/10.1016/j.cub.2022.08.019>
113. Ramesh, A.; Winkler, W.C. Magnesium-sensing riboswitches in bacteria. *RNA Biology.* **2010**, *7*, 77–83. <https://doi.org/10.4161/rna.7.1.10490>
114. Peng, W.T.; Zhang, L.D.; Zhou, Z.; Fu, C.; Chen, Z.C.; Liao, H. Magnesium promotes root nodulation through facilitation of carbohydrate allocation in soybean. *Physiol. Plant.* **2018**, *163*, 372–385. <https://doi.org/10.1111/pp1.12730>
115. Brear, E.M.; Day, D.A.; Smith, P.M.S. Iron: an essential micronutrient for the legume–rhizobium symbiosis. *Front. Plant Sci.* **2013**, *4*, 352. <https://doi.org/10.3389/fpls.2013.00359>
116. Liu, Y.; Xiong, Z.; Wu, W.; Ling, H.-Q.; Kong, D. Iron in the symbiosis of plants and microorganisms. *Plants.* **2023**, *12*, 1958. <https://doi.org/10.3390/plants12101958>
117. Banerjee, P.; Nath, R. Prospects of molybdenum fertilization in grain legumes – A review. *J. Plant Nutr.* **2022**, *45*, 1425–1440. <https://doi.org/10.1080/01904167.2021.2020831>
118. Bittner, F. Molybdenum metabolism in plants and crosstalk to iron. *Front. Plant Sci.* **2014**, *5*, 28. <https://doi.org/10.3389/fpls.2014.00028>
119. Bochniarz, J.; Bochniarz, M.; Lenartowicz, W. Effect of potassium and nitrogen fertilization on productivity of faba bean (*Vicia faba minor*) grown for seeds. *Pam. Pul.*, 1987, 89. [in Polish]
120. Zhou, J.; Sun, X.; Chen, C.; Chen, J. The effect of molybdenum fertilizer on the growth of grass-legume mixtures related to symbiotic rhizobium. *Agronomy.* **2023**, *13*, 495. <https://doi.org/10.3390/agronomy13020495>
121. Delgado, M.J.; Bedmar, E.J.; Downie, J.A. Genes involved in the formation and assembly of rhizobial cytochromes and their role in symbiotic nitrogen fixation. *Adv. Microb. Physiol.* **1998**, *40*, 191–231. [https://doi.org/10.1016/S0065-2911\(08\)60132-0](https://doi.org/10.1016/S0065-2911(08)60132-0)
122. Carranca, C.; De Varennes, A.; Rolston, D. Biological nitrogen fixation by fababean, pea and chickpea, under field conditions, estimated by the 15N isotope dilution technique. *Eur. J. Agron.* **1999**, *10*(1), 49–56. [https://doi.org/10.1016/S1161-0301\(98\)00049-5](https://doi.org/10.1016/S1161-0301(98)00049-5)
123. Redondo-Nieto, M.; Wilmort, A.R.; El-Hamdaoui, A.; Bonilla, I.; Bolaños, L. Relationship between boron and calcium in the N₂-fixing legume–rhizobia symbiosis. *Plant Cell Environ.* **2003**, *26*, 1905–1915. <https://doi.org/10.1046/j.1365-3040.2003.01107.x>
124. Fabiano, C.C.; Tezotto, T.; Favarin, J.L.; Polacco, J.C.; Mazzafera, P. Essentiality of nickel in plants: a role in plant stresses. *Frontiers in Plant Science.* **2015**, *6*, 754. <https://doi.org/10.3389/fpls.2015.00754>
125. Li, Y.; Liu, Q.; Zhang, D.-X.; Zhang, Z.-Y.; Xu, A.; Jiang, Y.-L.; Chen, Z.-C. Metal nutrition and transport in the process of symbiotic nitrogen fixation. *Plant Commun.* **2024**, *5*, 100829. <https://doi.org/10.1016/j.xplc.2024.100829>
126. Hu, X.; Wei, X.; Ling, J.; Chen, J. Cobalt: an essential micronutrient for plant growth? *Front. Plant Sci.* **2021**, *12*, 768523. <https://doi.org/10.3389/fpls.2021.768523>
127. Ma, J.; Song, Z.; Yang, J.; Wang, Y.; Han, H. (2021). Cobalt ferrite nanozyme for efficient symbiotic nitrogen fixation via regulating reactive oxygen metabolism. *Environmental Science: Nano.* **2021**, *8*, 188–203. <https://doi.org/10.1039/D0EN00935K>
128. León-Mediavilla, J.; Senovilla, M.; Montiel, J.; Gil-Díez, P.; Saez, Á.; Kryvoruchko, I.S.; Reguera, M.; Udvardi, M.K.; Imperial, J.; González-Guerrero, M. MtMTP2-facilitated zinc transport into intracellular compartments is essential for nodule development in *Medicago truncatula*. *Front. Plant Sci.* **2018**, *9*, 990. <https://doi.org/10.3389/fpls.2018.00990>
129. Broadley, M. R.; White, P. J.; Hammond, J. P.; Zelko, I.; Lux, A. (2007). Zinc in plants. *New Phytologist.* **2007**, *173*, 677–702. <https://doi.org/10.1111/j.1469-8137.2007.01996.x>
130. O'Hara, G.W. Nutritional constraints on root nodule bacteria affecting symbiotic nitrogen fixation: a review. *Aust. J. Exp. Agric.* **2001**, *41*, 417–433. <https://doi.org/10.1071/EA00087>
131. Weisany, W.; Raei, Y.; Allahverdipoor, K. H. Role of some of mineral nutrients in biological nitrogen fixation. *Bull. Env. Pharmacol. Life Sci.* **2013**, *2*, 77–84.
132. Alejandro, S.; Höller, S.; Meier, B.; Peiter, E. (2020). Manganese in plants: from acquisition to subcellular allocation. *Front. Plant Sci.* **2020**, *11*, 300. <https://doi.org/10.3389/fpls.2020.00300>
133. Rubio, L.M.; Ludden, P.W. Biosynthesis of the iron-molybdenum cofactor of nitrogenase. *Annu. Rev. Microbiol.* **2008**, *62*, 93–111. <https://doi.org/10.1146/annurev.micro.62.081307.162737>
134. Tripathi, R.; Tewari, R.; Singh, K.P.; Keswani, C.; Minkina, T.; Srivastava, A.K.; De Corato, U.; Sansinenea, E. Plant mineral nutrition and disease resistance: A significant linkage for sustainable crop protection. *Front. Plant Sci.* **2022**, *13*, 3116. <https://doi.org/10.3389/fpls.2022.883970>
135. Emamverdian, A.; Ding, Y.; Mokhberdoran, F.; Xie, Y. Heavy metal stress and some mechanisms of plant defense response. *Sci. World J.* **2015**, 756120. <http://dx.doi.org/10.1155/2015/756120>
136. Andersen, E.; Peiter, E.; Küpper, H. Trace metal metabolism in plants. *J. Exp. Bot.* **2018**, *69*, 909–954. <https://doi.org/10.1093/jxb/erx465>

137. Bethelenfalvay, G.; Franson, R. Manganese toxicity alleviated by mycorrhizae in soybean. *J. Plant Nutr.* **1989**, *12*(8), 953–970. DOI: 10.1080/01904168909364006
138. Nogueira, M.; Magalhaes, G.; Cardoso, E. Manganese toxicity in mycorrhizal and phosphorus-fertilized soybean plants. *J. Plant Nutr.* **2007**, *27*(1), 141–156. DOI: 10.1081/PLN-120027552
139. El-Jaoual, T.; Cox, D.A. Manganese toxicity in plants. *J. Plant Nutr.* **2008**, *21*(2), 353–386. DOI: 10.1080/01904169809365409
140. Sanchez-Pardo, B.; Fernandez-Pascual, M.; Zornoza, P. Copper microlocalisation, ultrastructural alterations and antioxidant responses in the nodules of white lupin and soybean plants grown under conditions of copper excess. *Environ. Exp. Botany* **2012**, *84*, 52–60. DOI: 10.1016/j.envexpbot.2012.04.017
141. Swędrzyńska, D.; Sawicka, A. The effect of copper on bacteria of the genus *Azospirillum* in the rhizosphere of maize and wheatseedlings. *Water-Environ.-Rural Areas* **2010**, *10*(2), 167–178.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.