

Review

Not peer-reviewed version

Signaling Pathways Regulating Dimorphism in Medically Relevant Fungal Species

[Uriel Ramírez-Sotelo](#) , [Manuela Gómez-Gaviria](#) , [Héctor M. Mora-Montes](#) *

Posted Date: 18 February 2025

doi: 10.20944/preprints202502.1154.v1

Keywords: transcription factor; environmental stimuli; yeast; hypha; kinase; antifungal drug



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Review

Signaling Pathways Regulating Dimorphism in Medically Relevant Fungal Species

Uriel Ramírez-Sotelo, Manuela Gómez-Gaviria and Héctor M. Mora-Montes *

Departamento de Biología, División de Ciencias Naturales y Exactas, Campus Guanajuato, Universidad de Guanajuato, Noria Alta s/n, col. Noria Alta, C.P., Guanajuato Gto. 36050, Mexico

* Correspondence: hmora@ugto.mx; Tel.: +52-473-732-0006 (ext. 8193)

Abstract: Pathogenic fungi exhibit morphological plasticity, a phenomenon known as dimorphism, which allows them to switch between yeast and hyphal forms in response to environmental stimuli. Thermodimorphic fungi undergo this transition based on temperature changes, playing a crucial role in their virulence and life cycle. Under saprobic conditions, some fungi exist as filamentous hyphae, producing conidia. When conidia are inhaled by mammals or traumatically inoculated body temperature (37°C) triggers their dimorphism into yeast cells. This shift promotes fungal dissemination and immune evasion. Some fungal pathogens undergo dimorphism in the contrary way, forming pseudohyphae and hyphae within the host. While temperature is a major driver of dimorphism, other factors, including CO₂ concentration, pH, nitrogen sources, and quorum-sensing molecules, also contribute to morphological shifts. This morphological transition is associated with increased expression of virulence factors that aid in adhesion, colonization, and immune evasion. This review examines the signaling pathways regulating fungal dimorphism, focusing on *H. capsulatum* as a model for mycelium-to-yeast transitions and *C. albicans* for yeast-to-hypha transitions. Understanding these pathways is essential for advancing therapeutic approaches against systemic fungal infections.

Keywords: transcription factor; environmental stimuli; yeast; hypha; kinase; antifungal drug

1. Introduction

The ability of pathogenic fungi to change their morphology during their life cycle is known as dimorphism and is carried out in response to specific environmental stimuli. Even though this trait is present across the whole fungi kingdom, few fungi can make the transition from yeasts to hyphae and vice versa [1]. Morphological changes are critical for the pathogenesis, virulence, stress adaptation, and life cycle of dimorphic fungi, and some pathogenic species can shift their morphology based on temperature changes, known as thermodimorphic fungi. Worldwide, thermodimorphic fungi are responsible for millions of human infections annually [2,3]. Among the best-known medically relevant fungal species with the ability to undergo dimorphism are *Histoplasma capsulatum*, *Paracoccidioides brasiliensis*, *Sporothrix schenckii*, *Candida albicans*, *Cryptococcus neoformans*, and *Blastomyces dermatitis*. These species are responsible for potentially fatal systemic mycoses in humans, particularly in immunocompromised individuals, although healthy individuals are also susceptible to these mycoses [4,5].

The reversible morphological transition between hypha and yeast, known as phase transition, is one of the most important features in the biology and lifestyle of dimorphic fungi. For those species naturally growing on soil or vegetal debris, at a temperature range between 22-25 °C, they usually grow as septate hyphae that produce conidia. Soil disturbance by different types of activities can aerosolize conidia and hyphal fragments [1,3]. This process can cause the inhalation of conidia and thus reach different hosts, such as mammals, which have an internal body temperature of 37 °C,

which favors the dissemination of the pathogen and the change of its morphology [1]. However, there are exceptions to this rule, since not all pathogenic fungi are saprophytes, and others that are classified as opportunistic that infect immunocompromised patients, such as species of the genera *Aspergillus*, *Fusarium*, *Scedosporium*, and *Candida* [6]. Although temperature is the predominant stimulus that influences the phase transition, other stimuli favor these changes, such as the availability of CO₂, pH, exogenous cysteine, estradiol, nitrogen sources, lipoxygenases, cyclooxygenases, and quorum sensing molecules [1,7,8].

Fungal dimorphism has also important implications for pathogenesis and immune evasion. For those species forming yeast cells within the host, this morphology has features that favor infection, such as the ability to survive in macrophages, immune cells that normally destroy pathogens [1,9,10]. In a similar line, those species that grow as hypha within the host have advantages in changing to this morphology, as these cells result big enough to be phagocytosed or the immune cells may be pierced by hypha growing inside the phagolysosome [11,12]. During morphological change, dimorphic fungi may undergo changes in the cell wall that lead to the reorganization of components, thus preventing the recognition of pathogen-associated molecular patterns by host immune cells [13–17]. Furthermore, morphological change is also associated with the upregulation of specific virulence factors that promote adhesion to host tissues and growth.

In this review, we focused on analyzing the main signaling pathways that govern fungal dimorphism in the transition of mycelial saprobic fungi that change to yeast-like morphology, taking *H. capsulatum* as a model, as well as those fungi that transition oppositely, that is, from yeast-like morphology to the formation of true hyphae, considering *C. albicans* as a biological model. The study of these pathways is crucial not only to understand the biology of these organisms, but also to improve prevention, diagnosis, and treatment strategies for systemic fungal infections.-

2. Fungal Dimorphism Regulators

2.1. Temperature-Responsive Genes

Thermally dimorphic fungi possess the ability to transit between two distinct morphological states, mycelium, and yeast. Currently, in all species studied, dimorphism occurs in response to environmental stimuli. Thermoresistant dimorphic pathogenic fungi, such as *H. capsulatum* grow on decaying organic matter with a filamentous morphology with the ability to produce vegetative conidia. Following inhalation of hyphal fragments or conidia by the host and subsequent growth at mammalian body temperature, *H. capsulatum* undergoes dimorphism to a yeast-like form capable of growing and spreading as a pathogen in mammals [18]. Because thermally dimorphic fungi can prevail in mammals after an acute infection resolves, it is thought that the parasitic form can return to the saprophytic environment after the death of infected animal hosts, facilitating a transition to the filamentous form, which helps maintain a latent infectious reservoir ready to continue this cycle [18]. In thermodimorphic fungi, genes that control growth in the yeast and filamentous phases have been identified and are considered master regulators governing the transitions, such as Ryp (Ryp1, Ryp2, and Ryp3), the histidine kinase Drk1 (dimorphism-regulating kinase 1), and the GATA family transcriptional regulator Sre1/SreB [19–22]. In contrast, fungal dimorphism can also occur as a morphological transition from a yeast form to a mycelium, which is necessary for dissemination, and *C. albicans* is the best fungal representative with this ability, where the yeast morphology favors its dissemination, and its mycelial morphology is tissue invasion. *C. albicans* dimorphism stimulation is also influenced by temperature and the first step for the transition consists of the germ tube formation [23]. In this organism, the family of cytoskeletal filament-forming proteins, septins, has been reported to contribute to morphogenesis by forming a scaffold to capture other proteins that promote septum formation, playing relevant roles during hyphal growth and the maintenance of polarized morphology, as well as proper selection of germ tube formation sites [23]. In *C. albicans*, at least seven genes encoding septins have been reported, where *CDC3* and *CDC12* are essential, but *CDC10*, *CDC11*, and *SEP7*, despite not being considered essential genes, contribute to the formation of the

septum and coordinate morphogenesis [23–25]. A *C. albicans* heterozygous *cdc12Δ/CDC12* mutant grew normally at 30 °C but showed an inability to grow at the temperature that induces hyphal formation (37 °C). Although this mutant formed germ tubes, a strongly polarized hyphal morphogenesis was not observed, as the cells acquired a pseudohyphal morphology, suggesting that septins play a special role in maintaining highly polarized growth [23]. Heat shock proteins (HSPs) also play a vital role in thermomorphism. When growing in temperatures that cause stress, such as 37°C, the main physiological role of HSPs is to promote protein folding, and one of the essential players is Hsp90 [26]. Hsp90, unlike other HSPs, is associated with a specific subset of proteins in the cell, called "client proteins", promoting their folding and maturation, such that Hsp90 generates protein folding reservoirs that can compensate for the phenotypic impacts of mutations in client proteins, facilitating evolutionary change [27]. *C. albicans* Hsp90 governs morphogenesis, such that engagement of Hsp90 function induces the transition from yeast to mycelium growth in the absence of any additional inducing signal and makes it temperature-dependent by inhibiting signaling through the cAMP-protein kinase A (PKA) pathway [27]. Furthermore, Hsp90 stabilizes other kinases of the cell wall integrity pathway, including Pkc1, Bck1, Mkk2, and Mkc1. Other proteins linked to Hsp90 include the cyclin Pcl1, the cyclin-dependent kinase Pho85, and the transcription factor Hms1, which are required for filamentation in response to Hsp90 inhibition and elevated temperature [27]. On the other hand, the heat shock transcription factor Hsf1 also plays a central role in regulating transcriptional responses following heat stress, including the upregulation of genes encoding other chaperones, such as *HSP70*, *HSP104*, and the same *HSP90*. Thus, modulation of *HSF1* levels also has a profound impact on morphogenesis [27].

Mucins are transmembrane glycoproteins involved in eukaryotic signaling [28]. In fungi, Msb2 is a regulator of environmental stress, cell wall biosynthesis, and the CEK pathway [28]. Msb2 plays a key role in *C. albicans* thermal adaptation, since Msb2 is essential for the hyphal formation and growth at 42 °C, regulating the high-temperature response through the CEK pathway [28]. Ssa1, an Hsp70 chaperone, and abundant cell wall component, was also found to be critical for thermal adaptation and the regulation of the MAP kinase pathways (CEK and PKC) [28].

It is known that in *H. capsulatum* the transcription factors Ryp1–4 (required for yeast-phase growth) are responsible for regulating the dimorphic change in response to temperature [29]. Through insertional mutagenesis, Nguyen & Sil [19] identified *RYP1*, which is required for *H. capsulatum* growth at 37 °C, and the mutants in this gene grew with filamentous morphology constitutively regardless of temperature. Furthermore, *RYP1* expression analysis revealed that it is required for the regulation of yeast phase-specific genes, as well as the regulation of two key genes in *H. capsulatum* virulence: the virulence factor *CBP1* and *YPS3*, which are required for the colonization of organs, such as the mouse lungs, liver, and spleen [19]. Ryp2 and Ryp3 are two additional regulators in *H. capsulatum* necessary for its growth as yeast at 37°C [20]. Mutants in *RYP2* and *RYP3* genes also lost their ability to grow as yeast at 37 °C and had defects in conidiation. Moreover, when murine bone marrow-derived macrophages were infected with conidia of the *RYP2* or *RYP3* mutant strains, most of the conidia did not germinate or were nonviable [20]. The fourth transcriptional regulator RYP was identified as a Zn(II)²Cys6 zinc binuclear cluster domain protein, whose expression analysis revealed that its transcription levels are higher in the yeast phase at 37 °C than in the mycelial morphology [30]. The transcription factors Ryp1–3 interact directly with DNA, particularly with a large set of genes related to virulence and cellular morphology [30]. Even though Ryp4 acts synergistically with Ryp1–3, its regulation is independent of them [30]. This temperature-responsive positive regulation circuit governs *H. capsulatum* dimorphism, generating the infective phase [30].

The HSPs play a relevant role in fungal dimorphism and pathogenesis because they modulate cell signaling in response to temperature variations associated with the transition from the saprobic stage to the warm environment into the host [31]. In *H. capsulatum*, it has been identified that Hsp70 is involved in the change from mycelium to yeast depending on the temperature change [32]. The *H. capsulatum* *HSP70* gene shares sequence homology with *HSP70* genes from other dimorphic fungi,

such as *Paracoccidioides brasiliensis* [33]. The expression of this gene is also inducible by the change in the *P. brasiliensis* growth temperature, as well as in the transition from mycelium to yeast, although its expression is differentially higher in yeast morphology [33]. Surprisingly, when the incubation of *P. brasiliensis* yeast was increased to 42 °C, a downregulation of the expression of several genes encoding other cellular proteins occurred, but there was a higher expression of four genes possibly corresponding to HSPs [33]. This fact suggests that Hsp70 expression is closely related to cell differentiation as a response of *P. brasiliensis* to temperature changes [33]. Other proteins that belong to this family are Hsp90 and have already been described for *H. capsulatum*, *C. neoformans*, *A. fumigatus*, *P. brasiliensis*, among others. *H. capsulatum* Hsp90 (Hsp82) is also heat-inducible and is predominantly expressed at 37°C once morphogenesis begins in the mycelial phase [34]. Insertion mutants affected in the Hsp82 expression levels showed attenuated virulence when confronted with bone marrow-derived macrophages, even though the growth of this mutant at 37 °C was normal [35,36]. The gene encoding Hsp90 has been identified as a single copy in the *P. brasiliensis* genome, is upregulated in yeast morphology, and its expression is upregulated in the initial stage of the mycelium-to-yeast transition [37]. Furthermore, Hsp90 from *P. brasiliensis* acts synergistically with calcineurin to promote dimorphism and prevents the generation of reactive oxygen species (ROS) by stimulating antioxidant defenses during thermal stress [38].

2.2. Mating is Associated with Dimorphism

Sexual reproduction in fungi is directed by a specialized region of the genome known as the mating-type (*MAT*) locus that confers cell type identity and controls fungal development during the sexual cycle [39]. Fungi can have self-compatibility to mating, meaning they do not need a genetically different partner (homothallic), or have one of two idiomorphs to mating (heterothallic) [39]. *C. neoformans* is a heterothallic fungus and its sexual reproduction is governed by this sex locus *MAT*, containing one of two possible idiomorphs, *MAT α* or *MATa* [40], which determine sexual compatibility, that is, one locus encodes homeodomain transcription factors and the other encodes pheromones and pheromone receptors [41]. Fungal cells with either the *MATa* or *MAT α* idiomorph are morphologically indistinguishable, their growth is similar in laboratory cultures and their mating occurs with the same ease in both cases, obtaining offspring in the same *MATa* and *MAT α* proportions [40]. In *C. neoformans*, only strains with the *MAT α* idiomorph develop conjugation tubes that elongate and fuse with *MATa* cells during mating. Then, *MATa* cells develop dikaryotic hyphae whose meiotic products give rise to basidiospores, in a similar way as occurs in dioecious animals and plants [40]. The mating type *MATa* and *MAT α* in *C. neoformans* has a connection with the global transcriptional repressor *TUP1* involved in the regulation of fungal growth and dimorphism, since the disruption of *TUP1* in a wild-type *MATa* strain crossed with another test strain (JEC32) resulted in extensive mating hyphae after 2 days of incubation, conversely, the hyphal development of the cross between the reintegrating strain and the test strain JEC32 was significantly reduced [40]. The frequency of mating by *TUP1* reintegrants was restored, whereas *tup1* null mutants transformed with the vector were not. These results indicate that loss of *TUP1* reduces mating efficiency and overall, *TUP1* has an important effect on the *C. neoformans* sexual reproduction [40]. Another transcriptional factor associated with mating type and that also regulates the signal transduction pathway of the mitogen-activated protein (MAP) kinase cascade is Ste12p [42]. It is also known that the MAPK pathway is directly involved in pheromone detection, which in turn, the main elements of this pathway coordinate morphogenesis in several fungal species, including *C. neoformans* [43]. This transcription factor regulates mating, pseudohyphal development, and haploid invasive growth [44]. A *C. neoformans* *ste12 α* null mutant did not show defects in yeast growth; however, it lost its ability to carry out haploid fruiting in a culture medium that favored filamentation [45]. In addition, the *STE12 α* deletion altered the expression of some virulence-associated genes, such as *CNLAC1*, *CAP59*, *CAP60*, and *CAP64*. Consequently, the *C. neoformans* *ste12 α* showed a global reduction in virulence [42,45]. Even genes that are not in the *MAT* locus, such as *CPR2*, *STE7*, and *CPK1*, are also involved in dimorphism and mating [43]. The *C. neoformans* homeodomain proteins Sxi1 α or Sxi2a are elements

that control newly fused cells from being unable to initiate a new developmental pathway, contributing to the specification of the $a + \alpha$ state of the newly formed dikaryon. The absence of these proteins causes fused cells to arrest and be unable to form filaments, basidia or dikaryotic spores [46].

Ascomycetes belonging to the *Candida* genus, such as the opportunistic pathogen *C. albicans*, were previously considered asexual or imperfect yeasts. However, there is currently evidence demonstrating their capacity for sexual or parasexual reproduction [47]. In the *Candida* genus, there are haploid or diploid species, where the haploids have the potential to carry out complete sexual cycles, such as *Candida lusitanae*. Despite not having the molecular machinery typically necessary to carry out meiosis, Ste12 seems to be the transcription factor essential for mating and meiosis [48–50]. On the other hand, in *C. albicans*, which is representative of the diploid species, *MTL* was identified, and it was shown that mating also involves a and α idiomorphs, where successful mating requires that the cells undergo a phenotypic change from the preestablished white state to the competent opaque state [48]. The *C. albicans* change of white-opaque state regulates mating to such an extent that it influences other aspects, including filamentation, interaction with the host immune-cells, and even virulence [48,51–53].

2.3. Signalling Cascades in Dimorphic Fungi

All living organisms share an elementary characteristic of their environment, that is, the ability to respond to external stimuli. Cells transform almost all external signals from their environment into required cellular responses through the activation of suitable signaling pathways [54]. The evolution of multicellular organisms, together with the specialization of the cells that shape them, has considerably increased the complexity of signaling pathways. These types of cellular responses are associated with basic elements such as growth and proliferation, to more complex processes like the development of structural support elements, transport, and supply of nutrients, among others [54]. When talking of fungal pathogens, the host represents a hostile environment, and signaling pathways help the fungal cell to adapt to these unfavorable conditions, such as body temperature, pH, and the particular host milieu. In addition, for growth as any other living organism, pathogenic fungi require essential nutrients for their development, such as carbohydrates, amino acids, nitrogen, and trace elements [55]. In fungi, the pathway for nutrient sensing is closely related to pheromone responsiveness for mating, growth, and cell morphology [56]. The G protein-coupled receptors (GPCRs) are a family of transmembrane receptors that are relevant to mediate the detection and transduction of signals in response to different environmental factors and their presence is evolutionarily conserved in practically all organisms [57]. The main objective of any organism in a constantly changing environment is adaptation to the availability of nutrients and stress, so the detection of these external variables is strictly necessary to adapt the morphology, metabolism, and activation of sexual reproduction and the virulence of fungi [55]. In this section, we will discuss the main key pathways in fungal growth and activation of dimorphism, with emphasis on the stages of these pathways that coordinate the transcriptional regulators directly involved in the change of fungal morphology.

2.3.1. Two-Component Regulatory Systems

These signal transduction pathways, strongly conserved in plants and microorganisms, are based on phosphorylation cascades between histidine kinases that end in the phosphorylation of a canonical aspartate residue in the response regulatory protein [58,59]. The two-component systems govern the dimorphism in several pathogenic fungi, such as *C. albicans*, *P. brasiliensis*, *Coccidioides immitis*, *Sporothrix schenckii*, *C. neoformans*, *Blastomyces dermatitidis*, *H. capsulatum* and *Penicillium marneffei* [29,59–62]. In addition, these pathways have been associated with morphogenesis control, response to oxidative and osmotic stress, quorum sensing, and virulence at least in *C. albicans* [59]. A classic two-component signaling system consists of a membrane-bound histidine protein kinase (HPK) and a response regulator (RR) protein, hence the name "two-component" [59]. The HPK is a dimer composed of two subunits, each of which has an ATP-binding domain, a dimerization domain,

and a kinase domain (containing the phosphorylation site, Figure 1, Panel A). Once the HPK input domain is stimulated by an external signal, the dimerization domain of one subunit approaches the kinase domain of the other subunit to promote phosphorylation (Figure 1, Panel A). The degree of HPK phosphorylation affects the level of RR phosphorylation [59]. In this respect, many HPKs could regulate one RR, or an HPK could regulate multiple RRs [59]. The HPKs are distributed and closely conserved in fungi that present the two-component system as a signal transduction pathway [60]. On the other hand, RR consists of a receptor module and an output domain, where the receptor module regulates the activity of the output domain through the phosphorylation of aspartate residues. In *C. albicans*, three RR, namely *SSK1*, *SKN7*, and *SRR1* have been identified [63,64]. Ssk1 and Skn7 play a notable role in morphogenesis and the response to oxidative stress, while Srr1 has implications in hyphal development and its loss makes *C. albicans* cells sensitive to osmotic stress and oxidative [63]. The domain or output structure of a two-component regulatory pathway can be a transcription factor that in turn controls gene expression or influences the regulation of target protein activity [59].




Figure 1. Schematic representation of the five signaling pathways associated with the regulation of fungal dimorphism. A) Two-component regulatory systems. B) Ca^{2+} /Calcineurin pathway. C) Mitogen-activated protein kinase (MAPK). D) Cyclic AMP (cAMP)/protein kinase A (PKA). E) Pal/Rim pathway. Detailed descriptions of each pathway can be found in the main text. Proteins separated by a slash (/) are functionally redundant. Arrows indicate activation, whereas inverted T-shaped bars represent inhibition. Double-headed arrows indicate physical interactions between proteins. Not all known pathway elements or interactions are shown in this representative model. Figure created with Biorender.com.

The two-component system is a multi-step phosphate transduction system in eukaryotes. However, in fungi, it occurs with slight variations. *C. albicans* contains three HPK genes (*SLN1*, *NIK1*, and *HK1*) and there are seven in *C. neoformans* [65,66]. The action mechanism consists of the donation of a phosphate from ATP to phosphorylate a conserved histidine residue (H box) after HPK perceives the stimulant signal. The phosphate is then transferred to the aspartate residue of the same HPK receptor domain, followed by transfer to the aspartate residue of the RR receptor domain via the histidine residue of the intermediate transfer protein, Ypd1 [64]. In total, four phosphorylation events occur in an ordered sequence, forming a four-step phosphate transfer system [59,65]. The relevance

that Ypd1 plays is its essentiality for the viability of eukaryotic microorganisms, such as *C. neoformans*, and its interaction with the response regulators Ssk1 and Skn7 has been demonstrated [67]. However, in *C. albicans*, *YPD1* is not essential for viability and its role seems to be linked to the activation of two signal transduction pathways: the Hog1 mitogen-activated protein kinase (MAPK) pathway, as well as Ssk1 and Skn7 that belong to the two-component pathway [64]. The three *C. albicans* genes seem to have a role in filamentation due to the loss of *NIK1* brought a severe defect in hyphal formation even when the *nik1Δ* mutant was grown in a culture medium supplemented with serum, an inducer of hyphal development [66]. In general, *SLN1* and *HK1* loss also affected hyphal formation compared to the wild-type strain. However, the most striking effect was observed in the *hk1Δ* mutant, where the hyphal formation was abolished [66]. Downstream of the pathway, Hog1 and Skn7 sense the Sln1 signal and regulate later gene expression [68]. The RR Skn7 is a specific and highly conserved transcription factor in ascomycete and basidiomycete and as we mentioned above, is required for *C. albicans* hyphal morphogenesis, since it was reported that the loss of *SKN7* has an impact on hyphal formation of *C. albicans* in solid but not liquid media, suggesting that Skn7 plays a highly specific role in surface-induced morphogenesis [58]. In addition, it has also been shown that *SKN7* overexpression induces the formation of hyphae regardless of their environment [69]. Skn7 is a positive regulator of the expression of other transcription factors that positively modulate morphogenesis, such as Brg1, Cph1, Czf1, Eed1, Sfl2, Tec1, and Ume6 [58]. This regulatory network seems to be associated with hyphal differentiation because the formation of biofilms is closely linked to hyphal growth and some of these transcription factors are responsible for this phenomenon [58].

A variant of the classical two-component system is the so-called phosphorelay, which begins with a hybrid kinase, an alternative form that contains a receptor domain fused to its C-terminal end. The hybrid kinase autophosphorylates and performs intramolecular phosphorylation of the receptor domain, then the phosphoryl group is transported to a histidine phosphotransferase (HPT) and this transfers it to a terminal RR to effect an output [70]. *DRK1* (dimorphism-regulating kinase 1) is a hybrid histidine kinase (HHK) transmembrane receptor highly conserved in some fungi. It is characterized by containing the catalytic and receptor domains in the same polypeptide, where the extracellular sensing domain that detects environmental signals is located at the amino-terminal end and the cytosolic signaling domain is at the C-terminal end [60,62,70]. Accordingly, a report revealed that there was a high percentage of similarity of the *DRK1* sequences of the species *H. capsulatum*, *B. dermatitidis*, *C. immitis*, *S. schenckii*, *C. neoformans*, *C. albicans*, and *P. brasiliensis* [60]. In *B. dermatitidis* and *H. capsulatum*, *DRK1* acts as a global regulator of dimorphism, and its activity is required for the transition from mycelium to yeast [61]. Disruption of *B. dermatitidis* *DRK1* resulted in the arrest of the fungus in mycelial morphology even under conditions that normally induce its transition to the yeast form and this has repercussions on the yeast phase-specific *BYS1* [61]. A similar gene regulation downstream of the pathway occurs in *H. capsulatum* where yeast phase-specific genes such as *CBP1*, *AGS1*, and *yps-3* are affected by the lack of *DRK1* [61]. In *S. schenckii*, *DRK1* lacks transmembrane domains, but it has PAS and GAF domains belonging to the families of cytoplasmic sensor domains located at the amino-terminal end [61]. Silencing of *S. schenckii* *DRK1* caused growth defects in the mutants, but no alterations in dimorphism were obtained because the mutants achieved the transition from mycelium to yeast, although with notable changes in yeast morphology [62]. The only correlation of *SsDRK1* with dimorphism is that its silencing caused a decrease in the expression of *Ste20*, a gene related to morphogenesis when *SsDRK1-i* yeasts were compared to wild-type strain [62]. Inhibition of *P. brasiliensis* *Drk1* with iprodione only delayed the change from mycelium to yeast without causing toxicity or inhibiting fungal growth. However, the inhibitor effect transiently maintained *Drk1* activity, but no overwhelming effect on arrest was obtained in any cell morphology [60]. Unlike previously described dimorphic fungi, *P. marneffei* has two genes, *DRKA* and *SLNA*, which encode histidine kinases that conform to its two-component system, and it was observed that the non-lethal deletion of *DRKA* and *SLNA* reduced hyphal formation, changed the number of septa, had abnormal chitin deposits in the cell wall of the hyphae and showed a higher rate of phagocytosis [29]. Current antifungal drugs are becoming ineffective against mycoses over time and new

alternatives are continually required to ensure adequate cure. However, the development of these drugs has been impeded by the structural and functional similarity of mammalian and fungal cells, which has questionable consequences if the drugs do not discriminate between host and pathogen cells. Interestingly, hybrid histidine kinases, such as the classic two-component system of fungi in general, are not elements shared with human cells or those of other mammals, which makes them more suitable candidates for therapeutic targets [71,72], hence the importance of continuing the study of the two-component signal transduction pathway in clinically relevant fungi.

2.3.2. Calcium/Calcineurin Pathway

Adaptation to different environmental conditions, which includes the response to various cations, is important for the survival and proliferation of microorganisms. Cells have intracellular differentiation systems, which identify these stimuli and can implement responses to counteract stress conditions [73]. Calcium (Ca^{2+}) signaling is preserved in eukaryotes and fulfills several functions, such as the detection of environmental stimuli, the transmission of extracellular signals to the nucleus to modulate the expression of some genes, morphological regulation, responses to abiotic and biotic stress and defense against pathogenicity and virulence [74,75]. When changes in cytosolic Ca^{2+} levels occur, several detecting proteins such as calmodulin and calcineurin can be activated, leading to the induction of different downstream signal transduction pathways [76,77].

Calcineurin (CaN) is known as a heterodimer that has two subunits, the catalytic subunit A (CnA) and the regulatory subunit B (CnB) (Figure 1, Panel B). Although it is well preserved in the different types of eukaryotes, the CaN plays diverse and different roles in organisms [77–80]. In the pathogenic species *C. neoformans* and *C. albicans*, calcineurin regulates growth at alkaline pH and high temperatures. In addition, it is also responsible for regulating dimorphism, mating, and virulence. In filamentous fungi, it regulates the hyphae growth, adaptation to stress, and cell wall integrity [77,81–83]. In *H. capsulatum*, this pathway is essential for the regulation of several cellular functions, especially those related to adaptation to stress conditions, contributing to fungal survival at high temperatures, and allowing adaptation to the host milieu [8,84].

In *C. albicans*, Ca^{2+} homeostasis plays an important role in survival and pathogenicity. Under different conditions, the Cch1-Mid1 calcium channel of the plasma membrane is activated, contributing to the entry of Ca^{2+} into the cells (Figure 1, Panel B) [85]. The increase in Ca^{2+} levels is subsequently detected by calmodulin (CaM) and then activates CaN, which in turn dephosphorylates Crz1 (Figure 1, Panel B), causing the expression of genes downstream that are dependent on Ca^{2+} -CaN, including *FKS2*, *CEK1*, *BMT3*, *CCH1*, *PMC1*, and *ERG26* [86,87]. In *C. albicans*, the study of these genes has been of great interest, because it has been shown that mutant strains in these genes have sensitivity to different antifungal drugs. In addition, several compounds directed to the proteins of this signaling pathway, such as CsA inhibitors, FK506 and Hsp90 (Figure 1, Panel B) exhibit antifungal activity alone or in combination with other antifungal drugs [85,88–90]. It has been reported that changes in *MID1*, *CCH1*, *YVC1*, and *SPF1* that are involved in this pathway, bring defects in virulence, sensitivity to drugs, such as fluconazole, terbinafine, and changes in hyphal development. In *C. albicans*, it has been shown that *PMC1* has an important role in morphological transition, biofilm formation, and virulence [91–95]. The CaN pathway plays a crucial role in dimorphism, allowing *C. albicans* to respond to environmental signals such as high temperature and pH, which are factors that favor hypha formation [82,96]. Different studies have shown that CaN is important for hyphal morphogenesis under stress conditions, such as high temperature and cell wall damage. CaN-deficient mutants (*cna1Δ* or *cnb1Δ*) showed a significant defect in hypha formation, suggesting that this pathway is important for dimorphism under conditions that mimic those of the human host [86]. CaN also regulates the ability of *C. albicans* to survive stressful conditions found within the host, such as osmotic stress, cell wall damage, and thermal stress. These factors are important for the survival of the fungus during infection. CaN activates genes that reinforce the cell wall integrity, allowing the fungus to resist the host's immune defenses and the attack by antifungal agents [85,97,98].

In *C. neoformans*, CaN is important for survival at high temperatures, and it is known that, in this fungus, CaN regulates genes that are essential for the response to oxidative stress, osmotic stress, and cell wall integrity. The survival of *C. neoformans* at a temperature of 37 °C favors its ability to proliferate in the host's internal environment. Mutants that lack the genes involved in this pathway are unable to survive at physiological temperatures and show changes in virulence in different infection models [99,100]. In *A. fumigatus*, mutations in *cnaA* and *cnaB* cause defects in germination, the morphology of the hyphae, and the formation of the septa. In addition, it is thought that these mutations are related to changes in the cell wall [101–103]. It is known that in this species the Ca^{2+} -CaN pathway regulates the polarized growth of the hyphae, which is essential for colonization and invasion in the host tissues. Unlike other species, in *A. fumigatus* growth is carried out through the extension of apical hyphae, which require effective machinery for transporting and assembling components of the cell wall. The CaN can regulate this process, since it controls the dynamics of the cytoskeleton, thus facilitating directional growth [103]. In *S. schenckii* and *P. brasiliensis*, the activation of this pathway is critical in the dimorphic transition from mycelium to yeast and facilitates the survival of the fungus within the host. In addition to its role in dimorphism, the Ca^{2+} -CaN pathway regulates mechanisms that allow the survival of these species in stress conditions, such as oxidative and osmotic stress, and promotes the integrity of the cell wall [104,105].

Despite the important role that Ca^{2+} and CaN play in fungal pathogenicity, their exploitation as a therapeutic target presents significant challenges due to their presence in human cells, where they regulate critical immune functions. However, efforts to develop specific inhibitors for fungal CaN, which do not interfere with the human version of the enzyme, are an active area of research. Combining CaN inhibitors with conventional antifungals, such as those that affect cell wall synthesis, could offer a synergistic strategy to treat resistant fungal infections. In the future, a deeper understanding of the molecular interactions of CaN with other signaling pathways in fungi could open new avenues for the development of more specific and effective therapies against clinically important fungi [79,101,103].

Finally, the Ca^{2+} -CaN pathway also interacts with other signaling pathways that regulate morphogenesis in different fungal species such as the cAMP/PKA and MAPK pathways, which will be described below. These interactions ensure that the morphological transition and response are effectively coordinated.

2.3.3. Mitogen-Activated Protein Kinase (MAPK)

In eukaryotic organisms, the family of serine/threonine protein kinases, known as mitogen-activated protein kinases (MAPKs) are involved in the transduction of a wide variety of extracellular signals and the regulation of different developmental processes [106]. Sequential activation of the MAPK cascade results in the activation of transcription factors and expression of specific gene sets in response to environmental stimuli [106]. In addition, this pathway responds to mating pheromones, maintenance of cell wall integrity, changes in osmolarity, and nutrient sensing [107].

In *Saccharomyces cerevisiae*, the HOG pathway consists of multiple, repetitive upstream osmosensing mechanisms leading to Hog1 activation. Briefly, the upstream osmosensing signaling of the HOG pathway consists of the *SLN1* and *SHO1* branches (Figure 1, Panel C). Ste11 is an upstream activator of the Hog1 MAPK in the *SHO1* branch, which is activated by phosphorylation by the Ste20/Cla4 kinases when osmotic stress is sensed [108]. In the *SHO1* branch of the HOG pathway, several non-kinase proteins such as Hkr1, Msb2, Sho1, Opy2, Ahk1, Bem1, and Ste50 are involved in the activation and/or regulation of the Hog1 MAPK [108]. The two HOG pathways converge in the activation of Pbs2, which is subsequently responsible for activating Hog1, which then activates genes responsible for promoting cellular osmoadaptation (Figure 1, Panel C) [108].

In *C. albicans*, the MAPK pathway is known to be essential for adaptation to different environmental conditions and virulence. In addition, it is involved in many cellular processes such as stress response, morphogenesis, biofilm formation, and antifungal resistance [106,109]. Signalling through the MAPK pathway activates different transcription factors, such as Cph1 and Efg1 that are

responsible for promoting the transition from the yeast to the filamentous form. Mutations in these genes disrupted the development of hyphae or pseudohyphae in response to many stimuli [106,110,111]. The *CEK1* pathway is also one of the key components of this regulation, by activating Cph1. The *C. albicans* *CEK1* deletion showed defects in hyphal formation in certain solid media, such as SLAD medium, where the nitrogen source is limited [112,113]. *HOG1* also responds to other stress conditions and participates in two important morphogenetic moments of *C. albicans*, filamentation and chlamydoconidia formation [109]. It has been shown that *HOG1* is activated in response to changes in osmotic potential, such as exposure to hypertonic solutions or changes in salt concentration. Such activation allows *C. albicans* to adjust its metabolism and cell structure to cope with adverse conditions. This metabolic adaptation allows the fungus to survive during the transition from the yeast- to the filamentous form. *HOG1* works together with other genes, such as *CPH1* to regulate dimorphism and interacts in different signaling pathways that regulate the morphological transition [109,114,115]. Mutants in *C. albicans* *HOG1* showed morphological alterations, as a result of a failure to complete the final stages of cytokinesis, with parallel defects in the budding pattern [116]. In addition, these mutants have different colonial morphologies compared to the parental strain in some media that promote morphological transition [116].

The *CEK1* and *HOG1* pathways also play an important role in *A. fumigatus*. Activation of *CEK1* is necessary for hyphal formation and conidial production in response to environmental changes. The transition from filamentous to more compact morphologies favors colonization of host tissues and biofilm formation [117,118]. On the other hand, the *HOG1*-mediated pathway is involved in the stress response, especially under conditions of high osmolarity. These changes allow the fungus to adapt and survive in unfavorable environments. *HOG1* is also related to other signaling pathways that regulate morphogenesis, for example, interacting with *CEK1* [118]. *C. neoformans* has different MAPKs, including Cpk1, Hog1, and Mpk1. However, these are mainly involved in cell wall integrity and maintenance [119]. In *P. brasiliensis* and *S. schenckii*, the activation of MAPK pathways is associated with the transition between yeast-like cells and filamentous forms. In particular, the MAPK pathway appears to be crucial for the transition to the filamentous form in response to environmental factors, such as temperature, pH, and nutrient availability [106,120]. In *H. capsulatum*, there are opposing regulatory pathways that control dimorphism in response to temperature, the *Msb2* regulon that is part of the MAPK pathway has implications for the regulation of yeast-specific genes. During the yeast-to-hyphal transition, in which cells transition from 37 °C to ambient temperatures, about 1,100 genes typical of the yeast phase reduce their expression [121,122]. However, in *msb2* mutants of *H. capsulatum* that were in yeast morphology, 40 of these genes continued to be inappropriately expressed. This group of 40 genes mainly includes direct transcriptional targets of Ryp proteins, indicating that these proteins are still present in the *msb2* mutant even at room temperature. Previous studies showed that, although Ryp protein levels decrease markedly in wild-type cells at room temperature, in the *msb2* mutant these proteins remain abundant. These results suggest that the *Msb2* pathway is crucial in preventing the accumulation of Ryp proteins at low temperatures. However, at 37 °C, the transcription factor Ryp3 binds to the *MSB2* promoter region and reduces its expression, implying that the circuit of Ryp proteins counteracts the *Msb2* pathway at high temperatures [30,121]. Thus, *Histoplasma* manages to switch between states in response to temperature through these opposing regulatory pathways [30,121].

The MAPK pathway is a key component in the biology of pathogenic fungi and represents an important target in the development of new antifungal drugs. The modulation of this pathway may offer innovative strategies to combat fungal infections, tackle drug resistance, and improve the effectiveness of existing therapies.

2.3.4. Cyclic AMP-Dependent Protein Kinase A

The cyclic AMP(cAMP)-dependent protein kinase A (PKA) pathway is involved in the perception and adaptation of fungi to the inhospitable environmental changes that occur during their life cycle, particularly in adaptation to nutrient accessibility [123]. Roughly, the cAMP/PKA pathway

is activated in response to nutrient availability in the environment, especially against carbon and nitrogen sources, such as glucose and amino acids, which are sensed by G protein-coupled receptors (GPCR) located on the cell surface (Figure 1, Panel D) [123]. The detected signal results in the activation of guanine nucleotide-binding proteins (G proteins) that induce an increase in the production of intracellular cAMP mediated by the adenylate cyclase enzyme [123]. The high cAMP concentration, known as the second messenger, activates cAMP-dependent protein kinase A, which subsequently phosphorylates target proteins located downstream (Figure 1, Panel D). The identity of these proteins in the final part of the pathway includes transcription factors, structural proteins, or even enzymes that carry out an immeasurable number of responses due to the activation of this pathway [123]. The Morphological transition and the expression of virulence factors are among the responses derived from the activation of this signaling pathway in medically relevant fungal pathogens [123,124]. This pathway has been identified in *C. albicans*, *C. neoformans*, *P. brasiliensis*, *M. rouxii*, and *P. marneffei* [125–129].

In *C. albicans*, there are two Ras GTPases, namely Ras1 and Ras2, a single adenylate cyclase, Cyr1 (also named as Cdc35), the low- and high-affinity cyclic nucleotide phosphodiesterases, Pde1 and Pde2, respectively, and protein kinase A conformed of regulatory and catalytic subunits (Figure 1, Panel D) [130]. Cdc25 is a guanine nucleotide exchange activator of Ras1 and Ras2, while Ira2 is its inactivator [130,131]. Similarly, in *C. neoformans* there are seven GPCR that detect glucose (Ste3/Cpr, Cpr2 y Gpr1–5), and several G proteins acting downstream GPCR have been identified, these are three subunits $G\alpha$ (Gpa1/3), Gpb1, Gib2, and Gpg1/2, where Gpa1 y Gib2 specifically participate in the cAMP/PKA pathway, as well Cac1 is the only adenylate cyclase, Aca1 induces Cac1 activity, Ras1/2 are Ras GTPases, the PKA is conformed by two catalytic subunits and one regulatory subunit [125]. In *C. albicans*, Ras1 is located near the plasma membrane through its C-terminal motif and is relevant to hyphal development and virulence [132], otherwise, Ras2 is special because it has low sequence similarity to other Ras proteins, but maintains the common CCIIT tag for membrane anchoring, suggesting that it is also associated with the plasma membrane [133]. Notably, Ras2 seems to have a lesser impact on morphogenesis, however, it acts synergistically with Ras1 and both proteins have an antagonistic role in the concentration of cAMP since mutants lacking Ras1/2 present a cell morphology affected by both a cAMP-dependent and cAMP-independent mechanisms [130,131,133]. The activation of this pathway is mediated by Cdc25, which activates Ras1/2. In *C. albicans*, loss of Cdc25 affected the growth rate in both liquid and solid medium, and on the cytotoxic effect against oral epithelial cells, despite the fact these cells are not capable of filamentation [131]. In this regard, filamentation is a *Candida* evasion strategy against the macrophage-mediated immune response [134] and unexpectedly it was observed that the hypofilamentous *cd25* Δ null mutant had an even higher survival rate than the wild-type strain, suggesting that other factors in addition to morphogenesis participate in the immune evasion mechanism [131]. The *C. neoformans* pathway activation consists of the coordination between the Gpa1 and the adenylate cyclase-associated protein Aca1, and both induce the Cac1 activity. The role of Aca1 is relevant for activation, and its elimination leads to problems in mating, capsule formation, and melanin synthesis. However, these phenotypes are restored with the external addition of cAMP [125,135]. An alternative way for Cac1 activation is by the transformation of CO₂ to bicarbonate, a reaction catalyzed by the carbonic anhydrase Can2. Therefore, the cAMP/PKA pathway can be alternatively triggered by host stimuli [125,136]. Both Cyr1 and Cac1 synthesize cAMP through cyclization of adenosine triphosphate (ATP) and this second messenger binds to the PKA regulatory subunit for its activation [125,131]. Similarly, the transient increase in Cyr1 is associated with the onset of dimorphic change in *P. brasiliensis*, suggesting that this pathway is actively involved in controlling dimorphic change [126]. The cAMP level is regulated by phosphodiesterases and *C. neoformans* genome contains *PDE1* and *PDE2* genes with such predicted activity. However, only Pde1 actively participates in the control of cAMP concentration, whereas in *C. albicans* both Pde1 (with low-affinity) and Pde2 (with high-affinity) phosphodiesterases participate in cAMP regulation and are associated with hyphal growth (Figure 1, Panel D) [137,138]. *C. neoformans* *PKA1* and *PKA2* encode for the catalytic subunits of PKA, whereas *PKR1* encodes for

its regulatory subunit and the whole holoenzyme is involved in melanin synthesis and capsule formation [125]. On the other hand, *C. albicans* PKA is integrated by catalytic subunits Tpk1 and Tpk2 as well as the regulatory subunit Bcy1 and the protein complex acts on transcription factors associated with filamentation (Figure 1, Panel D) [139]. Recently, it has been proposed that a dual-specificity tyrosine phosphorylation-regulated kinase, Yak1, acts downstream or in synergy with PKA to regulate filamentation in *C. albicans* since Yak1 is required for the yeast to hyphae transition (Figure 1, Panel D) [139]. On the opposite way, the methyltransferase Ctm1 is a crucial regulator in hyphal morphogenesis in *C. albicans*, as unmethylated Ctm1 binds to Tpk1 and Tpk2 inhibiting their kinase activity, which suppresses downstream regulation of the pathway by preventing hyphal development [128]. Once the PKA holoenzyme is activated its catalytic subunits activate several effector proteins (this encompasses transcription factors) downstream of the pathway through phosphorylation [131]. A transcription factor highly associated with this pathway is Efg1 (Figure 1, Panel D), which regulates the expression of genes involved in morphogenesis, the white-opaque phenotypic change, among other traits related to *C. albicans* virulence [140,141]. *C. albicans* Efg1 is homologous to *Saccharomyces cerevisiae* Sok2 and Pdh1, which are other transcription factors with functions in pseudohyphal growth, all of them belonging to the APSES family (Asm1, Phd1, Sok2, Efg1, and StuA) of transcriptional regulators (Figure 1, Panel D) [142]. *C. albicans* Efg1 has a dual negative and positive impact on hyphal development based on environmental stimulus and signaling cascades, such as cAMP/PKA because it directly regulates hyphal-associated gene expression [142]. Nrg1 is another transcription factor whose decreased expression depends on the cAMP/PKA pathway and plays an important role in repressing hyphal formation [143]. *C. neoformans* Nrg 1 is distantly related to the *C. albicans* ortholog, sharing only PKA consensus sequence similarity for its phosphorylation, but it is positively involved in cAMP-associated phenotypes, such as capsule formation and mating but not with melanin synthesis [144,145]. The *C. neoformans* genes downstream of PKA are mostly related to environmental stress responses, such as heat-shock proteins acting in response to stress and susceptibility to azole drugs, but not with hyphae formation [144,145]. In conclusion, the cAMP/PKA signalling pathway of *C. albicans* is associated with the control of the yeast to hyphal phenotypic change; however, in *C. neoformans* this pathway contributes more to maintaining pathogenesis against of stressful environmental factors and is not involved in hypha formation [128,146].

2.3.5. Pal/Rim Pathway

Environmental pH and its variations have a significant influence on microorganisms. Therefore, they have to sense and respond to these stimuli to maintain survival [147]. Different fungal species as well as other microorganisms share signaling pathways in response to environmental pH, and one of them acts particularly against the alkaline pH signal, namely Pal/Rim, which is very important for the fungal pathogenesis and secretion of various low molecular mass molecules into the extracellular medium [147,148]. The Rim/Pal pathway involves the activation of a zinc finger transcription factor, PacC in filamentous fungi, or Rim101 in yeast-like fungi, through a proteolytic cascade [148]. This master regulator in turn positively or negatively regulates the transcription of other genes involved in many physiological tasks, including dimorphism [148,149]. The general components of this pathway are conserved among filamentous and yeast-like cells fungi, such as *C. neoformans* and *C. albicans* [149–151]. However, among fungal species, there are notable particularities, especially in the transcriptional regulation downstream of this pathway [147]. This transcriptional specificity probably allows fungi a distinctive adaptation to species-specific environmental niches [147]. For example, pathogenic fungi of humans and other mammals can survive the unfavorable conditions of the host microenvironments where the pH can range from 2 to 10 because of signaling pathways that are essential for proper cellular functioning [147]. In human commensal fungi that colonize mucosal anatomical cavities, such as the gastrointestinal, oral-pharyngeal, and genitourinary tracts and that are also opportunistic with the ability to spread until reaching any organ, it is a challenge to recognize extracellular pH changes to survive, therefore it is not surprising that some of these fungi, such as *C.*

C. albicans have the Rim101 pathway [147,149,152] where the acidic pH favors the growth of its cells in the form of yeast, and, on the other hand, the neutral-alkaline pH promotes hypha formation [153]. However, the control of dimorphism depending on pH can occur oppositely, for example, in *Trichosporon cutaneum* the acidic pH drives filamentation, while the neutral-basic pH induces the growth of cells in yeast-like shape [154]. Extracellular pH recognition is mediated by the plasma membrane-associated receptor proteins Rim21/PalH and Dfg16/PalI (Figure 1, Panel E), where alkaline pH activates the Rim101/PacC pathway and in contrast, the acidic pH does not favor proteolysis-mediated activation of Rim101/PacC or is activated differently [152,155] (Figure 1, panel E). The activation of Rim21/PalH and Dfg16/PalI by neutral-alkaline pH promotes the ubiquitination of the Rim21-associated protein Rim8, which contributes to endocytosis [152] (Figure 1, panel E). *C. neoformans* Rim101 is intimately connected to the cAMP/PKA signal transduction cascade where Rim101 activation requires phosphorylation by PKA. The relationship of these two pathways seems to assign Rim101 a flexible role in the upstream signals that allow its activation. *C. neoformans* Rim101 does not seem to be only a pH-responsive factor but can respond to multiple host stimuli through the cAMP/PKA pathway, including other environmental signals, such as low iron levels or tissue culture medium [156].

Subsequently, Snf7 is oligomerized on the endosome surface and takes on Rim13/PalB, a calpain-like protease, and Rim20/PalA, another protease that binds to C-terminal inhibitory domain of the inactive transcription factor Rim101/PacC-long, which is engaged in the endosomal membrane [152,155]. In addition, Rim20 interacts with Vps4, which in turn interacts with Snf7 [155] (Figure 1, panel E). Bro1 was later found to interact synergistically with Vps4 and Snf7 in the transport of multivesicular bodies (MVB) to the vacuole [155]. The enrapture of MVB to the vacuole needs three protein complexes termed ESCRT-I-III (Endosomal Sorting Complex Required for Transport) [155]. For the operation of these complexes is required of Snf7, a cytoplasmatic protein that is recruited to endosomal membranes; Vps4, which is a cytoplasmic AAA-type ATPase that is present in the three ESCRT complexes to recycle components of the pathway; and Bro1 is also cytoplasmic protein that is associated with MVB and required Vps4 for to regulate endosomal dissociation of ESCRT-III [157]. Returning to Rim101/PacC, once Rim13/PalB performs the proteolytic processing on the C-terminal inhibitory domain of Rim101/PacC, its active form is obtained, which is suitable for translocating to the nucleus and regulating gene transcription associated with responses dependent on neutral-alkaline pH [149,152]. The identification of this pathway in *C. albicans* began with a differential expression analysis that identified two genes induced by alkaline pH, namely, PRA1 and PHR1 [149]. The PRA1 (pH-regulated antigen) gene encodes a protein located in the cell wall whose role is involved in alkali-induced filamentation, being that the PRA1 null mutant had a temperature-dependent defect in the initiation of filamentation, indicating its role in morphogenesis [158]. PHR1 is another pH-associated gene in *C. albicans*, whose expression was induced at alkaline pH and was repressed when pH values were lower than 5.5 [159]. The mutant lacking the two PHR1 alleles showed a defect in its morphology conditioned by pH, since at alkaline pH, this mutant could not form hyphae or yeasts as the parental strain used as control [159]. RIM101, RIM 20, and RIM8 are components of the only signaling pH-pathway Rim101p/PacC in *C. albicans* [149]. RIM101 encodes a pH-inducible zinc finger transcription factor that is dependent on Rim20 and Rim8 [149]. The operation of this pathway begins when the alkaline pH of the medium induces Rim8 and Rim20 activity on Rim101-long by proteolysis until it is processed into its Rim101p-short form, ready to carry out a response, that is, the activation (such as PHR1) or repression (e.g. PHR2) of genes that coordinate induced filamentation by alkali [149,160]. Finally, Rim101 activates filamentation by interacting with either the positive regulators Cph1 and Efg1 or the negative regulator Tup1 [149]. Efg1 is required for Rim101-induced filamentation but is not necessary for alkali-induced gene expression and Tup1 is not regulated during filamentous growth; otherwise, their associated DNA-binding proteins are likely regulated [160]. Thus, in *C. albicans*, Rim101 is a positive regulator of filamentation, as it acts strongly on ECE1, CSA1, CSA2, SAP5, HYR1, HWP1, RBT1, and IHD1 [161]. Unlike *C. albicans*, in *C. neoformans* the role of Rim101 is related to the formation of the capsule and

the fungal growth in various stressful environments, such as low iron concentrations, high salinity, or alkaline pH [151,156].

3. Transcriptional Regulation of Fungal Dimorphism

Signals derived from the environment or the host, such as nutrient availability, light, gases, stress, temperature, and pH, mainly activate the change in the morphology of fungal cells, through the signaling pathways previously described above, which ends in the transcription of genes that direct dimorphism [29,56,160]. The ability of fungi to transition from yeast to hyphal or mycelium to yeast growth in response to environmental factors is not only immanent to their pathogenicity, but this phenomenon is an excellent model for understanding how signaling pathways govern the growth and development of these organisms [160]. Activation of phase-specific genes in either hyphae or yeast contributes to cell cycle deregulation, establishment of polarity, and cell wall changes [58]. Fungal morphogenesis is under the control of a complex network of transcriptional regulators that act negatively or positively on the expression of phase-specific genes [58]. The negative regulation of hyphal-specific genes (HSGs) as it appears in *C. albicans* is driven by the transcriptional repressor Tup1, in association with the DNA-binding proteins Nrg1, Rfg1, and Sfl1 in synergy with the corepressor Ssn6 [58,162]. The positive regulation of HSGs is controlled under almost all conditions that induce hyphal formation by the transcription factors Skn7, Efg1, Flo8, and Ndt80, while other transcriptional factors, such as Cph1, Czf1, Tec1, Sfl2, and Rim101, positively regulate HSGs in a condition-specific manner or for each activated signaling pathway [58]. These transcriptional factors act mainly at the beginning of morphogenesis and their participation continues during hyphal growth, however, the transcriptional factor Ume6 has a major role in hyphal development [58]. Chromatin remodeling is a phenomenon that is also involved in morphogenesis since it is necessary for the transcriptional activation of phase-specific genes. In this sense, it has been shown that Brg1 binds to HSG promoters to counteract the repression mediated by the Nrg1-Tup1 complex, which promotes chromatin remodeling and favors HSG transcription [58].

In dimorphic fungal pathogens such as *C. albicans*, the role of genes upregulated in the yeast morphotype is fundamental and has been extensively studied, but genes that are silenced in yeast morphology, i.e., mold-specific genes in filamentous fungi that transition to yeast, also play a critical role in dimorphism and have not been described in the same detail.

4. Dimorphism Regulators with Therapeutic Potential

Current therapy against fungal infections is focused on target cellular components or processes such as ergosterol, sterol 14 α -demethylase, β -glucan synthesis, and interference with protein, RNA, or DNA synthesis [163]. However, each of these strategies has its drawbacks and limitations, such as high toxicity as well as the latent risk of developing drug resistance [163,164]. For this reason, new therapeutic approaches continue to be sought and proposed, such as drug repositioning, the use of natural or synthetic compounds modified to target other structural components of the cell wall, the plasma membrane, the regulation of gene expression, virulence factors, and even the signal transduction pathways [163]. Considering the drawbacks of conventional antifungal therapy, as well as the relevance of sanitary control of mycoses, the signaling pathways involved in the dimorphism of pathogenic fungi of clinical interest could have a wide potential for the development of new drugs against some elements that regulate these cascades [163]. The dimorphic regulatory histidine kinase 1, Drk1, (described in detail in the Two-component regulatory systems section) has been proposed as a key factor for inhibitory analysis in other clinically relevant species, such as *S. schenckii*, *P. marneffei*, and *P. brasiliensis* [62,165]. In a study involving fludioxonil, a group III histidine kinase inhibitor (iDrk1), it was suggested that *P. brasiliensis* Drk1 functions in fungal resistance to different cell wall disrupting agents by reducing viability after iDrk1 treatment [166]. In addition, a significantly higher phagocytic index was also observed in iDrk1-treated yeasts than in the control group through phagocytosis assays, suggesting the possible role of PbDrk1 in cell wall modulation, making it a

relevant target for further investigation [166]. Taking into consideration these multifaceted peculiarities of Drk1, a study was recently performed using a phage display technique to identify peptides capable of interacting with Drk1 of *P. brasiliensis*, and the effects of these interactions on the association with the *P. brasiliensis* host were examined [163]. The findings of the study were promising, as the peptides were found to be able to inhibit the morphological phase transition of *P. brasiliensis*. Furthermore, a substantial proportion of these peptides did not allow adhesion to pneumocytes. Although these peptides may not have inherent antifungal properties, they potentiate the effects of some antifungal agents. The efficacy of the peptides was also evaluated in the alternative host *Galleria mellonella* and was shown to contribute to improved larval survival rates [163]. Due to the broad role of the histidine kinase Drk1 is remarkably restricted to bacteria and fungi, it is of interest to continue its physiological research, but even more important would be to persist with developing new therapeutic proposals directed against Drk1.

The calcium-calcineurin signaling pathway of *C. albicans* has also been proposed as a candidate for the development of inhibitors to control infection by this opportunistic fungus [85]. Specifically, proteins that modulate the intracellular concentration of Ca^{2+} in both *C. albicans* and humans are of interest, since they have a low level of identity at the amino acid sequence level. For example, Cch1 has a sequence identity of 11.2% with its human counterpart. Also, the functional domains III S1, S2, and S3 of *C. albicans* Cch1 share little identity, with 11.1%, 34.8%, and 20.0%, respectively, compared to the domains of human Cch1 [85]. It is relevant to highlight that Mid1 and Yvc1 are fungi-specific proteins of the Ca^{2+} -CaN signaling pathway, which gives them the potential to be desirable targets for the development of drugs for the control of candidiasis [85,167]. Deletion of the *SPF1* results in defects in hyphal development and strongly attenuates *C. albicans* virulence [91]. Pmc1 also plays an important role in the morphological transition, biofilm formation, and virulence in a mouse model of disseminated infection. It was shown that the reduced susceptibility of the homozygous *pmc1* mutant to fluconazole was temperature dependent and therefore may be related to a form of azole tolerance [95]. CaN plays a central role in this and other signal transduction pathways, however in *C. albicans* it has been reported that the affectation of this gene produces in the mutants a greater susceptibility to fluconazole and a decrease in its virulence during a pulmonary infection [97,168]. Therefore, CaN appears to have a favorable potential to be targeted by antifungal drugs that cause its dysfunction. Despite the complete CaN protein of humans and fungi sharing a similarity greater than 40%, some functional domains of fungal CaN have clear differences, which is encouraging for its use in the development of inhibitory compounds that specifically bind to these domains and serve to combat *C. albicans* infections [85]. The transcriptional factor Crz1 and the protein Hsp90 also show promising potential as targets for the development of antifungal drugs and their possible use is also due to the low levels of similarity with the human orthologs [85]. The Hsp90 chaperone has promising potential for drug design, as the determination of the crystal structure of the N-terminal domain of *C. albicans* Hsp90 provided relevant information about the notable structural differences between fungal Hsp90 and its mammalian counterpart, despite the high sequence conservation within the nucleotide-binding domain [169].

The signaling involved in pH changes has also been addressed with pharmacological approaches to combat fungal infections caused by *A. fumigatus*, *C. albicans*, and *C. neoformans*, among others, specifically, attention has been paid to the Rim/Pal pathway as a source of targets to propose antifungal strategies [161]. Evidence of this is that the disruption of *VPS28* and *VPS32*, two Vps factors required for Rim pathway activation, increased the susceptibility of *C. albicans* to drugs directly involved with cell wall assembly at the level of chitin synthesis and deposition or the level of the construction of the β -1,3-glucan layer independently of the Rim pathway [170]. Recently, research on the Rim pathway has continued and it is now known that this entire cascade is associated with tolerance to azole compounds and echinocandins since the elimination of any of its elements brings with it hypersensitivity [171,172].

Within the protein kinase pathways, there are also distinguishable therapeutic targets, such as the cyclic AMP protein kinase A (cAMP-PKA) pathway, where the catalytic subunits Tpk2 and Cyr1

play an important role in the regulation of tolerance to drugs through the regulation of efflux pumps expression [172]. The relevance of the entire cAMP-PKA pathway has been studied together with its correlation with the regulation of susceptibility to antifungals in *C. auris*, and it is known that the pathway is closely related to antifungal resistance, such as amphotericin B, fluconazole, 5-fluorocytosine, caspofungin, and the fungicide fludioxonil [173]. Thus, it has been proposed that the development of inhibitors of the cAMP/PKA pathway could be favorable for the treatment of *C. auris* infection, in combination with some currently available antifungal drugs. Future perspectives are focused on the characterization of possible upstream regulators of Cyr1, such as a G protein-coupled receptor or small or heterotrimeric GTP-binding proteins; the identification of Cyr-independent upstream regulators of PKA; and the clarification of Cyr1- and PKA-dependent downstream signaling networks [173].

5. Conclusions

Many of the mechanisms described above support the hypothesis that some key elements of these pathways could be targets to develop new therapeutic approaches against fungal infections. We propose that it is necessary to continue investigating in detail each of the master regulators of dimorphism, especially those that are not shared with the mammalian hosts, that are conserved throughout the fungal kingdom, and that have promising potential as therapeutic targets. Understanding the key master regulators of these pathways in detail is a crucial step, since although there are many proteins involved in these processes with the potential to be inhibited, only those target elements that are highly conserved in pathogenic fungi and specific to the fungi kingdom should be selected. These relevant fungal-specific elements should be characterized from a genetic, biochemical, physiological, and pharmaceutical perspective to achieve greater efficiency in the development of new drugs. Some of the signaling pathways and their mechanisms described here are very well understood. However, others have not been well studied and this makes it difficult to know the true relevance of the components that comprise them. It is also crucial to know precisely the interweaving of the genetic networks that exist with other metabolic pathways that could be shared in humans and other mammals to guarantee the efficacy of the drugs that are designed based on this knowledge.

Author Contributions: The three authors contributed to the Conceptualization of the review work, Literature revision and analysis, and Draft preparation. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Secretaría de Ciencia, Humanidades, Tecnología e Innovación [Ciencia de Frontera 2019-6380 and CBF2023-2024-655]. The funding sources that supported this work did not have any involvement in the design, acquisition, and analysis of data and writing of the manuscript.

Institutional Review Board Statement Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Gauthier, G.M. Dimorphism in fungal pathogens of mammals, plants, and insects. *PLoS Pathog* **2015**, *11*, e1004608, doi:10.1371/journal.ppat.1004608.
2. Schwartz, I.S.; Govender, N.P.; Corcoran, C.; Dlamini, S.; Prozesky, H.; Burton, R.; Mendelson, M.; Taljaard, J.; Lehloeny, R.; Calligaro, G.; et al. Clinical characteristics, diagnosis, management, and outcomes of disseminated emmonsiosis: a retrospective case series. *Clin Infect Dis* **2015**, *61*, 1004-1012, doi:10.1093/cid/civ439.
3. Gauthier, G.M. Fungal dimorphism and virulence: molecular mechanisms for temperature adaptation, immune evasion, and in vivo survival. *Mediators Inflamm* **2017**, *2017*, 8491383, doi:10.1155/2017/8491383.

4. Gauthier, G.; Klein, B.S. Insights into fungal morphogenesis and immune evasion: fungal conidia, when situated in mammalian lungs, may switch from mold to pathogenic yeasts or spore-forming spherules. *Microbe Wash DC* **2008**, *3*, 416-423, doi:10.1128/microbe.3.416.1.
5. Pappas, P.G.; Alexander, B.D.; Andes, D.R.; Hadley, S.; Kauffman, C.A.; Freifeld, A.; Anaissie, E.J.; Brumble, L.M.; Herwaldt, L.; Ito, J.; et al. Invasive fungal infections among organ transplant recipients: results of the Transplant-Associated Infection Surveillance Network (TRANSNET). *Clin Infect Dis* **2010**, *50*, 1101-1111, doi:10.1086/651262.
6. Lopes, J.P.; Lionakis, M.S. Pathogenesis and virulence of *Candida albicans*. *Virulence* **2022**, *13*, 89-121, doi:10.1080/21505594.2021.2019950.
7. Ribes, J.A.; Vanover-Sams, C.L.; Baker, D.J. Zygomycetes in human disease. *Clin Microbiol Rev* **2000**, *13*, 236-301, doi:10.1128/cmr.13.2.236.
8. Lee, S.C.; Li, A.; Calo, S.; Heitman, J. Calcineurin plays key roles in the dimorphic transition and virulence of the human pathogenic zygomycete *Mucor circinelloides*. *PLoS Pathog* **2013**, *9*, e1003625, doi:10.1371/journal.ppat.1003625.
9. Jiménez Mdel, P.; Restrepo, A.; Radzioch, D.; Cano, L.E.; García, L.F. Importance of complement 3 and mannose receptors in phagocytosis of *Paracoccidioides brasiliensis* conidia by Nramp1 congenic macrophages lines. *FEMS Immunol Med Microbiol* **2006**, *47*, 56-66, doi:10.1111/j.1574-695X.2006.00059.x.
10. Guzman-Beltran, S.; Perez-Torres, A.; Coronel-Cruz, C.; Torres-Guerrero, H. Phagocytic receptors on macrophages distinguish between different *Sporothrix schenckii* morphotypes. *Microbes Infect* **2012**, *14*, 1093-1101, doi:10.1016/j.micinf.2012.06.001.
11. McKenzie, C.G.; Koser, U.; Lewis, L.E.; Bain, J.M.; Mora-Montes, H.M.; Barker, R.N.; Gow, N.A.; Erwig, L.P. Contribution of *Candida albicans* cell wall components to recognition by and escape from murine macrophages. *Infect Immun* **2010**, *78*, 1650-1658, doi:10.1128/IAI.00001-10.
12. Alonso, M.F.; Bain, J.M.; Erwig, L.P.; Brown, A.J.P.; Gow, N.A.R. Hyphal swelling induced in the phagosome of macrophages. *Fungal Biology* **2024**, *128*, 2148-2156, doi:<https://doi.org/10.1016/j.funbio.2024.08.011>.
13. García-Carnero, L.C.; Martínez-Duncker, I.; Gómez-Gaviria, M.; Mora-Montes, H.M. Differential recognition of clinically relevant *Sporothrix* species by human mononuclear cells. *J Fungi (Basel)* **2023**, *9*, 448, doi:10.3390/jof9040448.
14. Martínez-Álvarez, J.A.; Pérez-García, L.A.; Mellado-Mojica, E.; López, M.G.; Martínez-Duncker, I.; López-Bezerra, L.M.; Mora-Montes, H.M. *Sporothrix schenckii sensu stricto* and *Sporothrix brasiliensis* are differentially recognized by human peripheral blood mononuclear cells. *Front Microbiol* **2017**, *8*, 843, doi:10.3389/fmicb.2017.00843.
15. Mukaremera, L.; Lee, K.K.; Mora-Montes, H.M.; Gow, N.A.R. *Candida albicans* yeast, pseudohyphal, and hyphal morphogenesis differentially affects immune recognition. *Front Immunol* **2017**, *8*, 629, doi:10.3389/fimmu.2017.00629.
16. Villalobos-Duno, H.L.; Barreto, L.A.; Alvarez-Aular, Á.; Mora-Montes, H.M.; Lozoya-Pérez, N.E.; Franco, B.; Lopes-Bezerra, L.M.; Niño-Vega, G.A. Comparison of cell wall polysaccharide composition and structure between strains of *Sporothrix schenckii* and *Sporothrix brasiliensis*. *Front Microbiol* **2021**, *12*, 726958, doi:10.3389/fmicb.2021.726958.
17. Gantner, B.N.; Simmons, R.M.; Underhill, D.M. Dectin-1 mediates macrophage recognition of *Candida albicans* yeast but not filaments. *EMBO J* **2005**, *24*, 1277-1286, doi:10.1038/sj.emboj.7600594.
18. Gilmore, S.A.; Naseem, S.; Konopka, J.B.; Sil, A. N-acetylglucosamine (GlcNAc) triggers a rapid, temperature-responsive morphogenetic program in thermally dimorphic fungi. *PLoS Genet* **2013**, *9*, e1003799, doi:10.1371/journal.pgen.1003799.
19. Nguyen, V.Q.; Sil, A. Temperature-induced switch to the pathogenic yeast form of *Histoplasma capsulatum* requires Ryp1, a conserved transcriptional regulator. *Proc Natl Acad Sci U S A* **2008**, *105*, 4880-4885, doi:10.1073/pnas.0710448105.
20. Webster, R.H.; Sil, A. Conserved factors Ryp2 and Ryp3 control cell morphology and infectious spore formation in the fungal pathogen *Histoplasma capsulatum*. *Proc Natl Acad Sci U S A* **2008**, *105*, 14573-14578, doi:10.1073/pnas.0806221105.

21. Gauthier, G.M.; Sullivan, T.D.; Gallardo, S.S.; Brandhorst, T.T.; Vanden Wymelenberg, A.J.; Cuomo, C.A.; Suen, G.; Currie, C.R.; Klein, B.S. *SREB*, a GATA transcription factor that directs disparate fates in *Blastomyces dermatitidis* including morphogenesis and siderophore biosynthesis. *PLoS Pathog* **2010**, *6*, e1000846, doi:10.1371/journal.ppat.1000846.
22. Chao, L.Y.; Marletta, M.A.; Rine, J. Sre1, an iron-modulated GATA DNA-binding protein of iron-uptake genes in the fungal pathogen *Histoplasma capsulatum*. *Biochemistry* **2008**, *47*, 7274-7283, doi:10.1021/bi800066s.
23. Li, L.; Zhang, C.; Konopka, J.B. A *Candida albicans* temperature-sensitive cdc12-6 mutant identifies roles for septins in selection of sites of germ tube formation and hyphal morphogenesis. *Eukaryot Cell* **2012**, *11*, 1210-1218, doi:10.1128/ec.00216-12.
24. González-Novo, A.; Correa-Bordes, J.; Labrador, L.; Sánchez, M.; Vázquez de Aldana, C.R.; Jiménez, J. Sep7 is essential to modify septin ring dynamics and inhibit cell separation during *Candida albicans* hyphal growth. *Mol Biol Cell* **2008**, *19*, 1509-1518, doi:10.1091/mbc.e07-09-0876.
25. Warena, A.J.; Konopka, J.B. Septin function in *Candida albicans* morphogenesis. *Mol Biol Cell* **2002**, *13*, 2732-2746, doi:10.1091/mbc.e02-01-0013.
26. Shapiro, R.S.; Sellam, A.; Tebbji, F.; Whiteway, M.; Nantel, A.; Cowen, L.E. Pho85, Pcl1, and Hms1 signaling governs *Candida albicans* morphogenesis induced by high temperature or Hsp90 compromise. *Curr Biol* **2012**, *22*, 461-470, doi:10.1016/j.cub.2012.01.062.
27. Robbins, N.; Cowen, L.E. Roles of Hsp90 in *Candida albicans* morphogenesis and virulence. *Curr Opin Microbiol* **2023**, *75*, 102351, doi:10.1016/j.mib.2023.102351.
28. Saraswat, D.; Kumar, R.; Pande, T.; Edgerton, M.; Cullen, P.J. Signalling mucin Msb2 regulates adaptation to thermal stress in *Candida albicans*. *Mol Microbiol* **2016**, *100*, 425-441, doi:10.1111/mmi.13326.
29. Boyce, K.J.; Andrianopoulos, A. Fungal dimorphism: the switch from hyphae to yeast is a specialized morphogenetic adaptation allowing colonization of a host. *FEMS Microbiology Reviews* **2015**, *39*, 797-811, doi:10.1093/femsre/fuv035.
30. Beyhan, S.; Gutierrez, M.; Voorhies, M.; Sil, A. A temperature-responsive network links cell shape and virulence traits in a primary fungal pathogen. *PLoS Biol* **2013**, *11*, e1001614, doi:10.1371/journal.pbio.1001614.
31. Cleare, L.G.; Zamith-Miranda, D.; Nosanchuk, J.D. Heat shock proteins in *Histoplasma* and *Paracoccidioides*. *Clin Vaccine Immunol* **2017**, *24*, e00221-00217, doi:10.1128/cvi.00221-17.
32. Caruso, M.; Sacco, M.; Medoff, G.; Maresca, B. Heat shock 70 gene is differentially expressed in *Histoplasma capsulatum* strains with different levels of thermotolerance and pathogenicity. *Mol Microbiol* **1987**, *1*, 151-158, doi:10.1111/j.1365-2958.1987.tb00507.x.
33. da Silva, S.P.; Borges-Walmsley, M.I.; Pereira, I.S.; Soares, C.M.; Walmsley, A.R.; Felipe, M.S. Differential expression of an *hsp70* gene during transition from the mycelial to the infective yeast form of the human pathogenic fungus *Paracoccidioides brasiliensis*. *Mol Microbiol* **1999**, *31*, 1039-1050, doi:10.1046/j.1365-2958.1999.01236.x.
34. Minchiotti, G.; Gargano, S.; Maresca, B. The intron-containing hsp82 gene of the dimorphic pathogenic fungus *Histoplasma capsulatum* is properly spliced in severe heat shock conditions. *Mol Cell Biol* **1991**, *11*, 5624-5630, doi:10.1128/mcb.11.11.5624-5630.1991.
35. Edwards, J.A.; Zemska, O.; Rappleye, C.A. Discovery of a role for Hsp82 in *Histoplasma* virulence through a quantitative screen for macrophage lethality. *Infect Immun* **2011**, *79*, 3348-3357, doi:10.1128/iai.05124-11.
36. Rocha, M.C.; Minari, K.; Fabri, J.; Kerkaert, J.D.; Gava, L.M.; da Cunha, A.F.; Cramer, R.A.; Borges, J.C.; Malavazi, I. *Aspergillus fumigatus* Hsp90 interacts with the main components of the cell wall integrity pathway and cooperates in heat shock and cell wall stress adaptation. *Cell Microbiol* **2021**, *23*, e13273, doi:10.1111/cmi.13273.
37. Nicola, A.M.; Andrade, R.V.; Dantas, A.S.; Andrade, P.A.; Arraes, F.B.; Fernandes, L.; Silva-Pereira, I.; Felipe, M.S. The stress responsive and morphologically regulated hsp90 gene from *Paracoccidioides brasiliensis* is essential to cell viability. *BMC Microbiol* **2008**, *8*, 158, doi:10.1186/1471-2180-8-158.

38. Matos, T.G.; Morais, F.V.; Campos, C.B. Hsp90 regulates *Paracoccidioides brasiliensis* proliferation and ROS levels under thermal stress and cooperates with calcineurin to control yeast to mycelium dimorphism. *Med Mycol* **2013**, *51*, 413-421, doi:10.3109/13693786.2012.725481.
39. Morrow, C.A.; Fraser, J.A. Sexual reproduction and dimorphism in the pathogenic basidiomycetes. *FEMS Yeast Res* **2009**, *9*, 161-177, doi:10.1111/j.1567-1364.2008.00475.x.
40. Lee, H.; Chang, Y.C.; Kwon-Chung, K.J. *TUP1* disruption reveals biological differences between *MATa* and *MATα* strains of *Cryptococcus neoformans*. *Mol Microbiol* **2005**, *55*, 1222-1232, doi:<https://doi.org/10.1111/j.1365-2958.2004.04458.x>.
41. Fraser, J.A.; Heitman, J. Fungal mating-type loci. *Curr Biol* **2003**, *13*, R792-R795, doi:10.1016/j.cub.2003.09.046.
42. Chang, Y.C.; Penoyer, L.A.; Kwon-Chung, K.J. The second *STE12* homologue of *Cryptococcus neoformans* is *MATa*-specific and plays an important role in virulence. *Proc Natl Acad Sci U S A* **2001**, *98*, 3258-3263, doi:10.1073/pnas.061031998.
43. Lin, X. *Cryptococcus neoformans*: morphogenesis, infection, and evolution. *Infect Genet Evol* **2009**, *9*, 401-416, doi:10.1016/j.meegid.2009.01.013.
44. Roberts, R.L.; Fink, G.R. Elements of a single MAP kinase cascade in *Saccharomyces cerevisiae* mediate two developmental programs in the same cell type: mating and invasive growth. *Genes Dev* **1994**, *8*, 2974-2985, doi:10.1101/gad.8.24.2974.
45. Chang, Y.C.; Wickes, B.L.; Miller, G.F.; Penoyer, L.A.; Kwon-Chung, K.J. *Cryptococcus neoformans* *STE12α* regulates virulence but is not essential for mating. *J Exp Med* **2000**, *191*, 871-882, doi:10.1084/jem.191.5.871.
46. Hull, C.M.; Boily, M.J.; Heitman, J. Sex-specific homeodomain proteins Sxi1α and Sxi2α coordinately regulate sexual development in *Cryptococcus neoformans*. *Eukaryot Cell* **2005**, *4*, 526-535, doi:10.1128/ec.4.3.526-535.2005.
47. Butler, G.; Rasmussen, M.D.; Lin, M.F.; Santos, M.A.; Sakthikumar, S.; Munro, C.A.; Rheinbay, E.; Grabherr, M.; Forche, A.; Reedy, J.L.; et al. Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. *Nature* **2009**, *459*, 657-662, doi:10.1038/nature08064.
48. Bennett, R.J.; Turgeon, B.G. Fungal sex: the Ascomycota. *Microbiol Spectr* **2016**, *4*, doi:10.1128/microbiolspec.FUNK-0005-2016.
49. Reedy, J.L.; Floyd, A.M.; Heitman, J. Mechanistic plasticity of sexual reproduction and meiosis in the *Candida* pathogenic species complex. *Curr Biol* **2009**, *19*, 891-899, doi:10.1016/j.cub.2009.04.058.
50. Sherwood, R.K.; Scaduto, C.M.; Torres, S.E.; Bennett, R.J. Convergent evolution of a fused sexual cycle promotes the haploid lifestyle. *Nature* **2014**, *506*, 387-390, doi:10.1038/nature12891.
51. Kvaal, C.A.; Srikantha, T.; Soll, D.R. Misexpression of the white-phase-specific gene *WH11* in the opaque phase of *Candida albicans* affects switching and virulence. *Infect Immun* **1997**, *65*, 4468-4475, doi:10.1128/iai.65.11.4468-4475.1997.
52. Lohse, M.B.; Johnson, A.D. Differential phagocytosis of white versus opaque *Candida albicans* by *Drosophila* and mouse phagocytes. *PLoS One* **2008**, *3*, e1473, doi:10.1371/journal.pone.0001473.
53. Si, H.; Hernday, A.D.; Hirakawa, M.P.; Johnson, A.D.; Bennett, R.J. *Candida albicans* white and opaque cells undergo distinct programs of filamentous growth. *PLoS Pathog* **2013**, *9*, e1003210, doi:10.1371/journal.ppat.1003210.
54. Ma, Y.; Nicolet, J. Specificity models in MAPK cascade signaling. *FEBS Open Bio* **2023**, *13*, 1177-1192, doi:10.1002/2211-5463.13619.
55. Braunsdorf, C.; Mailänder-Sánchez, D.; Schaller, M. Fungal sensing of host environment. *Cell Microbiol* **2016**, *18*, 1188-1200, doi:10.1111/cmi.12610.
56. Bahn, Y.S.; Xue, C.; Idnurm, A.; Rutherford, J.C.; Heitman, J.; Cardenas, M.E. Sensing the environment: lessons from fungi. *Nat Rev Microbiol* **2007**, *5*, 57-69, doi:10.1038/nrmicro1578.
57. Xue, C.; Hsueh, Y.P.; Heitman, J. Magnificent seven: roles of G protein-coupled receptors in extracellular sensing in fungi. *FEMS Microbiol Rev* **2008**, *32*, 1010-1032, doi:10.1111/j.1574-6976.2008.00131.x.
58. Basso, V.; Znaidi, S.; Lagage, V.; Cabral, V.; Schoenherr, F.; LeibundGut-Landmann, S.; d'Enfert, C.; Bachellier-Bassi, S. The two-component response regulator Skn7 belongs to a network of transcription

- factors regulating morphogenesis in *Candida albicans* and independently limits morphogenesis-induced ROS accumulation. *Mol Microbiol* **2017**, *106*, 157-182, doi:10.1111/mmi.13758.
59. Liao, B.; Ye, X.; Chen, X.; Zhou, Y.; Cheng, L.; Zhou, X.; Ren, B. The two-component signal transduction system and its regulation in *Candida albicans*. *Virulence* **2021**, *12*, 1884-1899, doi:10.1080/21505594.2021.1949883.
 60. Chaves, A.F.; Navarro, M.V.; Castilho, D.G.; Calado, J.C.; Conceição, P.M.; Batista, W.L. A conserved dimorphism-regulating histidine kinase controls the dimorphic switching in *Paracoccidioides brasiliensis*. *FEMS Yeast Res* **2016**, *16*, fow047, doi:10.1093/femsyr/fow047.
 61. Hou, B.; Zhang, Z.; Zheng, F.; Liu, X. Molecular cloning, characterization and differential expression of *DRK1* in *Sporothrix schenckii*. *Int J Mol Med* **2013**, *31*, 99-104, doi:10.3892/ijmm.2012.1193.
 62. Zhang, Z.; Hou, B.; Wu, Y.Z.; Wang, Y.; Liu, X.; Han, S. Two-component histidine kinase *DRK1* is required for pathogenesis in *Sporothrix schenckii*. *Mol Med Rep* **2018**, *17*, 721-728, doi:10.3892/mmr.2017.8005.
 63. Desai, C.; Mavrianos, J.; Chauhan, N. *Candida albicans* *SRR1*, a putative two-component response regulator gene, is required for stress adaptation, morphogenesis, and virulence. *Eukaryot Cell* **2011**, *10*, 1370-1374, doi:10.1128/ec.05188-11.
 64. Mavrianos, J.; Desai, C.; Chauhan, N. Two-component histidine phosphotransfer protein Ypd1 is not essential for viability in *Candida albicans*. *Eukaryot Cell* **2014**, *13*, 452-460, doi:10.1128/ec.00243-13.
 65. Kruppa, M.; Calderone, R. Two-component signal transduction in human fungal pathogens. *FEMS Yeast Res* **2006**, *6*, 149-159, doi:10.1111/j.1567-1364.2006.00024.x.
 66. Yamada-Okabe, T.; Mio, T.; Ono, N.; Kashima, Y.; Matsui, M.; Arisawa, M.; Yamada-Okabe, H. Roles of three histidine kinase genes in hyphal development and virulence of the pathogenic fungus *Candida albicans*. *J Bacteriol* **1999**, *181*, 7243-7247, doi:10.1128/jb.181.23.7243-7247.1999.
 67. Lee, J.W.; Ko, Y.J.; Kim, S.Y.; Bahn, Y.S. Multiple roles of Ypd1 phosphotransfer protein in viability, stress response, and virulence factor regulation in *Cryptococcus neoformans*. *Eukaryot Cell* **2011**, *10*, 998-1002, doi:10.1128/ec.05124-11.
 68. Li, S.; Ault, A.; Malone, C.L.; Raitt, D.; Dean, S.; Johnston, L.H.; Deschenes, R.J.; Fassler, J.S. The yeast histidine protein kinase, Sln1p, mediates phosphotransfer to two response regulators, Ssk1p and Skn7p. *Embo j* **1998**, *17*, 6952-6962, doi:10.1093/emboj/17.23.6952.
 69. Chauvel, M.; Nesser, A.; Cabral, V.; Znaidi, S.; Goyard, S.; Bachellier-Bassi, S.; Firon, A.; Legrand, M.; Diogo, D.; Naulleau, C.; et al. A versatile overexpression strategy in the pathogenic yeast *Candida albicans*: identification of regulators of morphogenesis and fitness. *PLoS One* **2012**, *7*, e45912, doi:10.1371/journal.pone.0045912.
 70. Capra, E.J.; Laub, M.T. Evolution of two-component signal transduction systems. *Annu Rev Microbiol* **2012**, *66*, 325-347, doi:10.1146/annurev-micro-092611-150039.
 71. Tebbets, B.; Yu, Z.; Stewart, D.; Zhao, L.X.; Jiang, Y.; Xu, L.H.; Andes, D.; Shen, B.; Klein, B. Identification of antifungal natural products via *Saccharomyces cerevisiae* bioassay: insights into macrotetrolide drug spectrum, potency and mode of action. *Med Mycol* **2013**, *51*, 280-289, doi:10.3109/13693786.2012.710917.
 72. Kennedy, E.N.; Menon, S.K.; West, A.H. Extended N-terminal region of the essential phosphorelay signaling protein Ypd1 from *Cryptococcus neoformans* contributes to structural stability, phosphostability and binding of calcium ions. *FEMS Yeast Res* **2016**, *16*, fow068, doi:10.1093/femsyr/fow068.
 73. Colinet, A.S.; Sengottaiyan, P.; Deschamps, A.; Colsoul, M.L.; Thines, L.; Demaegd, D.; Duchêne, M.C.; Foulquier, F.; Hols, P.; Morsomme, P. Yeast Gdt1 is a Golgi-localized calcium transporter required for stress-induced calcium signaling and protein glycosylation. *Sci Rep* **2016**, *6*, 24282, doi:10.1038/srep24282.
 74. Xu, J.R. Map kinases in fungal pathogens. *Fungal Genet Biol* **2000**, *31*, 137-152, doi:10.1006/fgbi.2000.1237.
 75. Miskei, M.; Karányi, Z.; Pócsi, I. Annotation of stress-response proteins in the aspergilli. *Fungal Genet Biol* **2009**, *46 Suppl 1*, S105-S120, doi:10.1016/j.fgb.2008.07.013.
 76. Matsumoto, T.K.; Ellsmore, A.J.; Cessna, S.G.; Low, P.S.; Pardo, J.M.; Bressan, R.A.; Hasegawa, P.M. An osmotically induced cytosolic Ca²⁺ transient activates calcineurin signaling to mediate ion homeostasis and salt tolerance of *Saccharomyces cerevisiae*. *J Biol Chem* **2002**, *277*, 33075-33080, doi:10.1074/jbc.M205037200.

77. Wang, S.; Zhou, H.; Wu, J.; Han, J.; Li, S.; Shao, S. Transcriptomic analysis reveals genes mediating salt tolerance through calcineurin/CchA-Independent signaling in *Aspergillus nidulans*. *Biomed Res Int* **2017**, *2017*, 4378627, doi:10.1155/2017/4378627.
78. Mendoza, I.; Rubio, F.; Rodriguez-Navarro, A.; Pardo, J.M. The protein phosphatase calcineurin is essential for NaCl tolerance of *Saccharomyces cerevisiae*. *J Biol Chem* **1994**, *269*, 8792-8796.
79. Fox, D.S.; Cruz, M.C.; Sia, R.A.; Ke, H.; Cox, G.M.; Cardenas, M.E.; Heitman, J. Calcineurin regulatory subunit is essential for virulence and mediates interactions with FKBP12-FK506 in *Cryptococcus neoformans*. *Mol Microbiol* **2001**, *39*, 835-849, doi:10.1046/j.1365-2958.2001.02295.x.
80. Stie, J.; Fox, D. Calcineurin regulation in fungi and beyond. *Eukaryot Cell* **2008**, *7*, 177-186, doi:10.1128/ec.00326-07.
81. Kothe, G.O.; Free, S.J. Calcineurin subunit B is required for normal vegetative growth in *Neurospora crassa*. *Fungal Genet Biol* **1998**, *23*, 248-258, doi:10.1006/fgbi.1998.1037.
82. Cruz, M.C.; Goldstein, A.L.; Blankenship, J.R.; Del Poeta, M.; Davis, D.; Cardenas, M.E.; Perfect, J.R.; McCusker, J.H.; Heitman, J. Calcineurin is essential for survival during membrane stress in *Candida albicans*. *Embo j* **2002**, *21*, 546-559, doi:10.1093/emboj/21.4.546.
83. Cervantes-Chávez, J.A.; Ali, S.; Bakkeren, G. Response to environmental stresses, cell-wall integrity, and virulence are orchestrated through the calcineurin pathway in *Ustilago hordei*. *Mol Plant Microbe Interact* **2011**, *24*, 219-232, doi:10.1094/mpmi-09-10-0202.
84. Garfoot, A.L.; Rappleye, C.A. *Histoplasma capsulatum* surmounts obstacles to intracellular pathogenesis. *Febs j* **2016**, *283*, 619-633, doi:10.1111/febs.13389.
85. Li, W.; Shrivastava, M.; Lu, H.; Jiang, Y. Calcium-calcineurin signaling pathway in *Candida albicans*: A potential drug target. *Microbiol Res* **2021**, *249*, 126786, doi:10.1016/j.micres.2021.126786.
86. Bader, T.; Bodendorfer, B.; Schröppel, K.; Morschhäuser, J. Calcineurin is essential for virulence in *Candida albicans*. *Infect Immun* **2003**, *71*, 5344-5354, doi:10.1128/iai.71.9.5344-5354.2003.
87. Liu, S.; Hou, Y.; Liu, W.; Lu, C.; Wang, W.; Sun, S. Components of the calcium-calcineurin signaling pathway in fungal cells and their potential as antifungal targets. *Eukaryot Cell* **2015**, *14*, 324-334, doi:10.1128/ec.00271-14.
88. Young, J.C.; Moarefi, I.; Hartl, F.U. Hsp90: a specialized but essential protein-folding tool. *J Cell Biol* **2001**, *154*, 267-273, doi:10.1083/jcb.200104079.
89. Onyewu, C.; Blankenship, J.R.; Del Poeta, M.; Heitman, J. Ergosterol biosynthesis inhibitors become fungicidal when combined with calcineurin inhibitors against *Candida albicans*, *Candida glabrata*, and *Candida krusei*. *Antimicrob Agents Chemother* **2003**, *47*, 956-964, doi:10.1128/aac.47.3.956-964.2003.
90. Li, L.; An, M.; Shen, H.; Huang, X.; Yao, X.; Liu, J.; Zhu, F.; Zhang, S.; Chen, S.; He, L.; et al. The non-Geldanamycin Hsp90 inhibitors enhanced the antifungal activity of fluconazole. *Am J Transl Res* **2015**, *7*, 2589-2602.
91. Yu, Q.; Wang, H.; Cheng, X.; Xu, N.; Ding, X.; Xing, L.; Li, M. Roles of Cch1 and Mid1 in morphogenesis, oxidative stress response and virulence in *Candida albicans*. *Mycopathologia* **2012**, *174*, 359-369, doi:10.1007/s11046-012-9569-0.
92. Chang, Y.; Schlenstedt, G.; Flockerzi, V.; Beck, A. Properties of the intracellular transient receptor potential (TRP) channel in yeast, Yvc1. *FEBS Lett* **2010**, *584*, 2028-2032, doi:10.1016/j.febslet.2009.12.035.
93. Chen, Y.; Mallick, J.; Maqnas, A.; Sun, Y.; Choudhury, B.I.; Côte, P.; Yan, L.; Ni, T.J.; Li, Y.; Zhang, D.; et al. Chemogenomic profiling of the fungal pathogen *Candida albicans*. *Antimicrob Agents Chemother* **2018**, *62*, e02365-02317, doi:10.1128/aac.02365-17.
94. Jiang, L.; Xu, D.; Hameed, A.; Fang, T.; Bakr Ahmad Fazili, A.; Asghar, F. The plasma membrane protein Rch1 and the Golgi/ER calcium pump Pmr1 have an additive effect on filamentation in *Candida albicans*. *Fungal Genet Biol* **2018**, *115*, 1-8, doi:10.1016/j.fgb.2018.04.001.
95. Luna-Tapia, A.; DeJarnette, C.; Sansevere, E.; Reitler, P.; Butts, A.; Hevener, K.E.; Palmer, G.E. The vacuolar Ca(2+) ATPase pump Pmc1p is required for *Candida albicans* pathogenesis. *mSphere* **2019**, *4*, e00715-00718, doi:10.1128/mSphere.00715-18.
96. Karababa, M.; Valentino, E.; Pardini, G.; Coste, A.T.; Bille, J.; Sanglard, D. CRZ1, a target of the calcineurin pathway in *Candida albicans*. *Mol Microbiol* **2006**, *59*, 1429-1451, doi:10.1111/j.1365-2958.2005.05037.x.

97. Blankenship, J.R.; Wormley, F.L.; Boyce, M.K.; Schell, W.A.; Filler, S.G.; Perfect, J.R.; Heitman, J. Calcineurin is essential for *Candida albicans* survival in serum and virulence. *Eukaryot Cell* **2003**, *2*, 422-430, doi:10.1128/ec.2.3.422-430.2003.
98. Chen, Y.L.; Brand, A.; Morrison, E.L.; Silao, F.G.; Bigol, U.G.; Malbas, F.F., Jr.; Nett, J.E.; Andes, D.R.; Solis, N.V.; Filler, S.G.; et al. Calcineurin controls drug tolerance, hyphal growth, and virulence in *Candida dubliniensis*. *Eukaryot Cell* **2011**, *10*, 803-819, doi:10.1128/ec.00310-10.
99. Odom, A.; Muir, S.; Lim, E.; Toffaletti, D.L.; Perfect, J.; Heitman, J. Calcineurin is required for virulence of *Cryptococcus neoformans*. *Embo j* **1997**, *16*, 2576-2589, doi:10.1093/emboj/16.10.2576.
100. Kraus, P.R.; Nichols, C.B.; Heitman, J. Calcium- and calcineurin-independent roles for calmodulin in *Cryptococcus neoformans* morphogenesis and high-temperature growth. *Eukaryot Cell* **2005**, *4*, 1079-1087, doi:10.1128/ec.4.6.1079-1087.2005.
101. Steinbach, W.J.; Cramer, R.A., Jr.; Perfect, B.Z.; Asfaw, Y.G.; Sauer, T.C.; Najvar, L.K.; Kirkpatrick, W.R.; Patterson, T.F.; Benjamin, D.K., Jr.; Heitman, J.; et al. Calcineurin controls growth, morphology, and pathogenicity in *Aspergillus fumigatus*. *Eukaryot Cell* **2006**, *5*, 1091-1103, doi:10.1128/ec.00139-06.
102. Juvvadi, P.R.; Fortwendel, J.R.; Rogg, L.E.; Burns, K.A.; Randell, S.H.; Steinbach, W.J. Localization and activity of the calcineurin catalytic and regulatory subunit complex at the septum is essential for hyphal elongation and proper septation in *Aspergillus fumigatus*. *Mol Microbiol* **2011**, *82*, 1235-1259, doi:10.1111/j.1365-2958.2011.07886.x.
103. Juvvadi, P.R.; Lamothe, F.; Steinbach, W.J. Calcineurin as a multifunctional regulator: unraveling novel functions in fungal stress responses, hyphal growth, drug resistance, and pathogenesis. *Fungal Biol Rev* **2014**, *28*, 56-69, doi:10.1016/j.fbr.2014.02.004.
104. Campos, C.B.; Di Benedetto, J.P.; Morais, F.V.; Ovalle, R.; Nobrega, M.P. Evidence for the role of calcineurin in morphogenesis and calcium homeostasis during mycelium-to-yeast dimorphism of *Paracoccidioides brasiliensis*. *Eukaryot Cell* **2008**, *7*, 1856-1864, doi:10.1128/ec.00110-08.
105. García-Carnero, L.C.; Martínez-Álvarez, J.A. Virulence factors of *Sporothrix schenckii*. *J Fungi (Basel)* **2022**, *8*, 318, doi:10.3390/jof8030318.
106. Zhao, X.; Mehrabi, R.; Xu, J.R. Mitogen-activated protein kinase pathways and fungal pathogenesis. *Eukaryot Cell* **2007**, *6*, 1701-1714, doi:10.1128/ec.00216-07.
107. May, G.S.; Xue, T.; Kontoyiannis, D.P.; Gustin, M.C. Mitogen activated protein kinases of *Aspergillus fumigatus*. *Medical Mycology* **2005**, *43*, S83-S86, doi:10.1080/13693780400024784.
108. Takayama, T.; Yamamoto, K.; Saito, H.; Tatebayashi, K. Interaction between the transmembrane domains of Sho1 and Opy2 enhances the signaling efficiency of the Hog1 MAP kinase cascade in *Saccharomyces cerevisiae*. *PLoS One* **2019**, *14*, e0211380, doi:10.1371/journal.pone.0211380.
109. Monge, R.A.; Román, E.; Nombela, C.; Pla, J. The MAP kinase signal transduction network in *Candida albicans*. *Microbiology (Reading)* **2006**, *152*, 905-912, doi:10.1099/mic.0.28616-0.
110. Lo, H.J.; Köhler, J.R.; DiDomenico, B.; Loebenberg, D.; Cacciapuoti, A.; Fink, G.R. Nonfilamentous *C. albicans* mutants are avirulent. *Cell* **1997**, *90*, 939-949, doi:10.1016/s0092-8674(00)80358-x.
111. Braun, B.R.; Head, W.S.; Wang, M.X.; Johnson, A.D. Identification and characterization of *TUP1*-regulated genes in *Candida albicans*. *Genetics* **2000**, *156*, 31-44, doi:10.1093/genetics/156.1.31.
112. Csank, C.; Schröppel, K.; Leberer, E.; Harcus, D.; Mohamed, O.; Meloche, S.; Thomas, D.Y.; Whiteway, M. Roles of the *Candida albicans* mitogen-activated protein kinase homolog, Cek1p, in hyphal development and systemic candidiasis. *Infect Immun* **1998**, *66*, 2713-2721, doi:10.1128/iai.66.6.2713-2721.1998.
113. Srinivasa, K.; Kim, J.; Yee, S.; Kim, W.; Choi, W. A MAP kinase pathway is implicated in the pseudohyphal induction by hydrogen peroxide in *Candida albicans*. *Mol Cells* **2012**, *33*, 183-193, doi:10.1007/s10059-012-2244-y.
114. Arana, D.M.; Nombela, C.; Alonso-Monge, R.; Pla, J. The Pbs2 MAP kinase kinase is essential for the oxidative-stress response in the fungal pathogen *Candida albicans*. *Microbiology (Reading)* **2005**, *151*, 1033-1049, doi:10.1099/mic.0.27723-0.
115. Román, E.; Nombela, C.; Pla, J. The Sho1 adaptor protein links oxidative stress to morphogenesis and cell wall biosynthesis in the fungal pathogen *Candida albicans*. *Mol Cell Biol* **2005**, *25*, 10611-10627, doi:10.1128/mcb.25.23.10611-10627.2005.

116. Alonso-Monge, R.; Navarro-García, F.; Molero, G.; Diez-Orejas, R.; Gustin, M.; Pla, J.; Sánchez, M.; Nombela, C. Role of the mitogen-activated protein kinase Hog1p in morphogenesis and virulence of *Candida albicans*. *J Bacteriol* **1999**, *181*, 3058-3068, doi:10.1128/jb.181.10.3058-3068.1999.
117. Du, C.; Sarfati, J.; Latge, J.P.; Calderone, R. The role of the sakA (Hog1) and tcsB (sln1) genes in the oxidant adaptation of *Aspergillus fumigatus*. *Med Mycol* **2006**, *44*, 211-218, doi:10.1080/13693780500338886.
118. Ma, D.; Li, R. Current understanding of HOG-MAPK pathway in *Aspergillus fumigatus*. *Mycopathologia* **2013**, *175*, 13-23, doi:10.1007/s11046-012-9600-5.
119. -Jang, Y.-B.; Kim, J.-Y.; Bahn, Y.-S. Unraveling the cryptic functions of mitogen-activated protein kinases Cpk2 and Mpk2 in *Cryptococcus neoformans*. *mBio* **2024**, *15*, e01156-01124, doi:doi:10.1128/mbio.01156-24.
120. González-Rubio, G.; Fernández-Acero, T.; Martín, H.; Molina, M. Mitogen-activated protein kinase phosphatases (MKPs) in fungal signaling: conservation, function, and regulation. *Int J Mol Sci* **2019**, *20*, 1709, doi:10.3390/ijms20071709.
121. Rodriguez, L.; Voorhies, M.; Gilmore, S.; Beyhan, S.; Myint, A.; Sil, A. Opposing signaling pathways regulate morphology in response to temperature in the fungal pathogen *Histoplasma capsulatum*. *PLoS Biol* **2019**, *17*, e3000168, doi:10.1371/journal.pbio.3000168.
122. Sil, A. Molecular regulation of *Histoplasma dimorphism*. *Curr Opin Microbiol* **2019**, *52*, 151-157, doi:10.1016/j.mib.2019.10.011.
123. Choi, J.; Jung, W.H.; Kronstad, J.W. The cAMP/protein kinase A signaling pathway in pathogenic basidiomycete fungi: Connections with iron homeostasis. *J Microbiol* **2015**, *53*, 579-587, doi:10.1007/s12275-015-5247-5.
124. Leberer, E.; Marcus, D.; Dignard, D.; Johnson, L.; Ushinsky, S.; Thomas, D.Y.; Schröppel, K. Ras links cellular morphogenesis to virulence by regulation of the MAP kinase and cAMP signalling pathways in the pathogenic fungus *Candida albicans*. *Mol Microbiol* **2001**, *42*, 673-687, doi:10.1046/j.1365-2958.2001.02672.x.
125. Caza, M.; Kronstad, J.W. The cAMP/protein kinase a pathway regulates virulence and adaptation to host conditions in *Cryptococcus neoformans*. *Front Cell Infect Microbiol* **2019**, *9*, 212, doi:10.3389/fcimb.2019.00212.
126. Chen, D.; Janganan, T.K.; Chen, G.; Marques, E.R.; Kress, M.R.; Goldman, G.H.; Walmsley, A.R.; Borges-Walmsley, M.I. The cAMP pathway is important for controlling the morphological switch to the pathogenic yeast form of *Paracoccidioides brasiliensis*. *Mol Microbiol* **2007**, *65*, 761-779, doi:10.1111/j.1365-2958.2007.05824.x.
127. Zaremborg, V.; Donella-Deana, A.; Moreno, S. Mechanism of activation of cAMP-dependent protein kinase: in *Mucor rouxii* the apparent specific activity of the cAMP-activated holoenzyme is different than that of its free catalytic subunit. *Arch Biochem Biophys* **2000**, *381*, 74-82, doi:10.1006/abbi.2000.1948.
128. Zeng, G.; Xu, X.; Kok, Y.J.; Deng, F.S.; Ling Chow, E.W.; Gao, J.; Bi, X.; Wang, Y. Cytochrome c regulates hyphal morphogenesis by interfering with cAMP-PKA signaling in *Candida albicans*. *Cell Rep* **2023**, *42*, 113473, doi:10.1016/j.celrep.2023.113473.
129. Zuber, S.; Hynes, M.J.; Andrianopoulos, A. G-protein signaling mediates asexual development at 25 degrees C but has no effect on yeast-like growth at 37 degrees C in the dimorphic fungus *Penicillium mameffeii*. *Eukaryot Cell* **2002**, *1*, 440-447, doi:10.1128/ec.1.3.440-447.2002.
130. Huang, G.; Huang, Q.; Wei, Y.; Wang, Y.; Du, H. Multiple roles and diverse regulation of the Ras/cAMP/protein kinase A pathway in *Candida albicans*. *Mol Microbiol* **2019**, *111*, 6-16, doi:10.1111/mmi.14148.
131. Wijnants, S.; Vreys, J.; Nysten, J.; Van Dijck, P. The Cdc25 and Ras1 proteins of *Candida albicans* Influence epithelial toxicity in a niche-specific way. *J Fungi (Basel)* **2023**, *9*, 201, doi:10.3390/jof9020201.
132. Piispanen, A.E.; Bonnefoi, O.; Carden, S.; Deveau, A.; Bassilana, M.; Hogan, D.A. Roles of Ras1 membrane localization during *Candida albicans* hyphal growth and farnesol response. *Eukaryot Cell* **2011**, *10*, 1473-1484, doi:10.1128/ec.05153-11.
133. Zhu, Y.; Fang, H.M.; Wang, Y.M.; Zeng, G.S.; Zheng, X.D.; Wang, Y. Ras1 and Ras2 play antagonistic roles in regulating cellular cAMP level, stationary-phase entry and stress response in *Candida albicans*. *Mol Microbiol* **2009**, *74*, 862-875, doi:10.1111/j.1365-2958.2009.06898.x.

134. Gómez-Gaviria, M.; Ramírez-Sotelo, U.; Mora-Montes, H.M. Non-*albicans* *Candida* species: immune response, evasion mechanisms, and new plant-derived alternative therapies. *J Fungi (Basel)* **2022**, *9*, 11, doi:10.3390/jof9010011.
135. Bahn, Y.S.; Hicks, J.K.; Giles, S.S.; Cox, G.M.; Heitman, J. Adenylyl cyclase-associated protein Aca1 regulates virulence and differentiation of *Cryptococcus neoformans* via the cyclic AMP-protein kinase A cascade. *Eukaryot Cell* **2004**, *3*, 1476-1491, doi:10.1128/ec.3.6.1476-1491.2004.
136. Mogensen, E.G.; Janbon, G.; Chaloupka, J.; Steegborn, C.; Fu, M.S.; Moyrand, F.; Klengel, T.; Pearson, D.S.; Geeves, M.A.; Buck, J.; et al. *Cryptococcus neoformans* senses CO₂ through the carbonic anhydrase Can2 and the adenylyl cyclase Cac1. *Eukaryot Cell* **2006**, *5*, 103-111, doi:10.1128/ec.5.1.103-111.2006.
137. Hicks, J.K.; Bahn, Y.S.; Heitman, J. Pde1 phosphodiesterase modulates cyclic AMP levels through a protein kinase A-mediated negative feedback loop in *Cryptococcus neoformans*. *Eukaryot Cell* **2005**, *4*, 1971-1981, doi:10.1128/ec.4.12.1971-1981.2005.
138. Davis-Hanna, A.; Piispanen, A.E.; Stateva, L.I.; Hogan, D.A. Farnesol and dodecanol effects on the *Candida albicans* Ras1-cAMP signalling pathway and the regulation of morphogenesis. *Mol Microbiol* **2008**, *67*, 47-62, doi:10.1111/j.1365-2958.2007.06013.x.
139. MacAlpine, J.; Liu, Z.; Hossain, S.; Whitesell, L.; Robbins, N.; Cowen, L.E. DYRK-family kinases regulate *Candida albicans* morphogenesis and virulence through the Ras1/PKA pathway. *mBio* **2023**, *14*, e0218323, doi:10.1128/mbio.02183-23.
140. Giusani, A.D.; Vines, M.; Kumamoto, C.A. Invasive filamentous growth of *Candida albicans* is promoted by Czf1p-dependent relief of Efg1p-mediated repression. *Genetics* **2002**, *160*, 1749-1753, doi:10.1093/genetics/160.4.1749.
141. Biswas, S.; Van Dijk, P.; Datta, A. Environmental sensing and signal transduction pathways regulating morphopathogenic determinants of *Candida albicans*. *Microbiol Mol Biol Rev* **2007**, *71*, 348-376, doi:10.1128/mmbr.00009-06.
142. Glazier, V.E. *EFG1*, everyone's favorite gene in *Candida albicans*: a comprehensive literature review. *Front Cell Infect Microbiol* **2022**, *12*, 855229, doi:10.3389/fcimb.2022.855229.
143. Su, C.; Yu, J.; Lu, Y. Hyphal development in *Candida albicans* from different cell states. *Curr Genet* **2018**, *64*, 1239-1243, doi:10.1007/s00294-018-0845-5.
144. Maeng, S.; Ko, Y.J.; Kim, G.B.; Jung, K.W.; Floyd, A.; Heitman, J.; Bahn, Y.S. Comparative transcriptome analysis reveals novel roles of the Ras and cyclic AMP signaling pathways in environmental stress response and antifungal drug sensitivity in *Cryptococcus neoformans*. *Eukaryot Cell* **2010**, *9*, 360-378, doi:10.1128/ec.00309-09.
145. Bahn, Y.S.; Jung, K.W. Stress signaling pathways for the pathogenicity of *Cryptococcus*. *Eukaryot Cell* **2013**, *12*, 1564-1577, doi:10.1128/ec.00218-13.
146. Maliehe, M.; Ntoi, M.A.; Lahiri, S.; Folorunso, O.S.; Ogundeji, A.O.; Pohl, C.H.; Sebolai, O.M. Environmental factors that contribute to the maintenance of *Cryptococcus neoformans* pathogenesis. *Microorganisms* **2020**, *8*, doi:10.3390/microorganisms8020180.
147. Selvig, K.; Alspaugh, J.A. pH Response pathways in fungi: adapting to host-derived and environmental signals. *Mycobiology* **2011**, *39*, 249-256, doi:10.5941/myco.2011.39.4.249.
148. Cervantes-Chávez, J.A.; Ortiz-Castellanos, L.; Tejeda-Sartorius, M.; Gold, S.; Ruiz-Herrera, J. Functional analysis of the pH responsive pathway Pal/Rim in the phytopathogenic basidiomycete *Ustilago maydis*. *Fungal Genet Biol* **2010**, *47*, 446-457, doi:10.1016/j.fgb.2010.02.004.
149. Davis, D.; Wilson, R.B.; Mitchell, A.P. *RIM101*-dependent and-independent pathways govern pH responses in *Candida albicans*. *Mol Cell Biol* **2000**, *20*, 971-978, doi:10.1128/mcb.20.3.971-978.2000.
150. Davis, D.; Edwards, J.E., Jr.; Mitchell, A.P.; Ibrahim, A.S. *Candida albicans* *RIM101* pH response pathway is required for host-pathogen interactions. *Infect Immun* **2000**, *68*, 5953-5959, doi:10.1128/iai.68.10.5953-5959.2000.
151. Ost, K.S.; O'Meara, T.R.; Huda, N.; Esher, S.K.; Alspaugh, J.A. The *Cryptococcus neoformans* alkaline response pathway: identification of a novel rim pathway activator. *PLoS Genet* **2015**, *11*, e1005159, doi:10.1371/journal.pgen.1005159.

152. Davis, D.A. How human pathogenic fungi sense and adapt to pH: the link to virulence. *Curr Opin Microbiol* **2009**, *12*, 365-370, doi:10.1016/j.mib.2009.05.006.
153. Shu, T.; He, X.Y.; Chen, J.W.; Mao, Y.S.; Gao, X.D. The pH-responsive transcription factors YlRim101 and Mhy1 regulate alkaline pH-induced filamentation in the dimorphic yeast *Yarrowia lipolytica*. *mSphere* **2021**, *6*, e00179-00121, doi:10.1128/mSphere.00179-21.
154. Zhu, L.B.; Wang, Y.; Zhang, Z.B.; Yang, H.L.; Yan, R.M.; Zhu, D. Influence of environmental and nutritional conditions on yeast–mycelial dimorphic transition in *Trichosporon cutaneum*. *Biotechnology & Biotechnological Equipment* **2017**, *31*, 516-526, doi:10.1080/13102818.2017.1292149.
155. Kullas, A.L.; Li, M.; Davis, D.A. Snf7p, a component of the ESCRT-III protein complex, is an upstream member of the RIM101 pathway in *Candida albicans*. *Eukaryot Cell* **2004**, *3*, 1609-1618, doi:10.1128/ec.3.6.1609-1618.2004.
156. O'Meara, T.R.; Norton, D.; Price, M.S.; Hay, C.; Clements, M.F.; Nichols, C.B.; Alspaugh, J.A. Interaction of *Cryptococcus neoformans* Rim101 and protein kinase A regulates capsule. *PLoS Pathog* **2010**, *6*, e1000776, doi:10.1371/journal.ppat.1000776.
157. Odorizzi, G.; Katzmann, D.J.; Babst, M.; Audhya, A.; Emr, S.D. Bro1 is an endosome-associated protein that functions in the MVB pathway in *Saccharomyces cerevisiae*. *J Cell Sci* **2003**, *116*, 1893-1903, doi:10.1242/jcs.00395.
158. Sentandreu, M.; Elorza, M.V.; Sentandreu, R.; Fonzi, W.A. Cloning and characterization of PRA1, a gene encoding a novel pH-regulated antigen of *Candida albicans*. *J Bacteriol* **1998**, *180*, 282-289, doi:10.1128/jb.180.2.282-289.1998.
159. Saporito-Irwin, S.M.; Birse, C.E.; Sypherd, P.S.; Fonzi, W.A. PHR1, a pH-regulated gene of *Candida albicans*, is required for morphogenesis. *Mol Cell Biol* **1995**, *15*, 601-613, doi:10.1128/mcb.15.2.601.
160. Liu, H. Transcriptional control of dimorphism in *Candida albicans*. *Curr Opin Microbiol* **2001**, *4*, 728-735, doi:10.1016/s1369-5274(01)00275-2.
161. Cornet, M.; Gaillardin, C. pH signaling in human fungal pathogens: a new target for antifungal strategies. *Eukaryot Cell* **2014**, *13*, 342-352, doi:10.1128/ec.00313-13.
162. Zhao, R.; Lockhart, S.R.; Daniels, K.; Soll, D.R. Roles of TUP1 in switching, phase maintenance, and phase-specific gene expression in *Candida albicans*. *Eukaryot Cell* **2002**, *1*, 353-365, doi:10.1128/ec.1.3.353-365.2002.
163. Marcos, C.M.; de Oliveira, H.C.; Assato, P.A.; de Oliveira, L.T.; Fregonezi, N.; Dos Santos, K.S.; Costa-Orlandi, C.B.; Fusco-Almeida, A.M.; Mendes-Giannini, M.J.S. Polypeptides targeting *Paracoccidioides brasiliensis* Drk1. *J Fungi (Basel)* **2023**, *9*, 980, doi:10.3390/jof9100980.
164. de Oliveira, H.C.; Bezerra, B.T.; Rodrigues, M.L. Antifungal development and the urgency of minimizing the impact of fungal diseases on public health. *ACS Bio Med Chem Au* **2023**, *3*, 137-146, doi:10.1021/acsbiomedchemau.2c00055.
165. Marcos, C.M.; de Oliveira, H.C.; Assato, P.A.; Castelli, R.F.; Fusco-Almeida, A.M.; Mendes-Giannini, M.J.S. Drk1, a Dimorphism Histidine Kinase, contributes to morphology, virulence, and stress adaptation in *Paracoccidioides brasiliensis*. *J Fungi (Basel)* **2021**, *7*, 852, doi:10.3390/jof7100852.
166. Navarro, M.V.; de Barros, Y.N.; Segura, W.D.; Chaves, A.F.A.; Jannuzzi, G.P.; Ferreira, K.S.; Xander, P.; Batista, W.L. The Role of Dimorphism Regulating Histidine Kinase (Drk1) in the pathogenic fungus *Paracoccidioides brasiliensis* cell wall. *J Fungi (Basel)* **2021**, *7*, 1014, doi:10.3390/jof7121014.
167. Martin, D.C.; Kim, H.; Mackin, N.A.; Maldonado-Báez, L.; Evangelista, C.C., Jr.; Beaudry, V.G.; Dudgeon, D.D.; Naiman, D.Q.; Erdman, S.E.; Cunningham, K.W. New regulators of a high affinity Ca²⁺ influx system revealed through a genome-wide screen in yeast. *J Biol Chem* **2011**, *286*, 10744-10754, doi:10.1074/jbc.M110.177451.
168. Sanglard, D.; Ischer, F.; Marchetti, O.; Entenza, J.; Bille, J. Calcineurin A of *Candida albicans*: involvement in antifungal tolerance, cell morphogenesis and virulence. *Mol Microbiol* **2003**, *48*, 959-976, doi:10.1046/j.1365-2958.2003.03495.x.
169. Whitesell, L.; Robbins, N.; Huang, D.S.; McLellan, C.A.; Shekhar-Guturja, T.; LeBlanc, E.V.; Nation, C.S.; Hui, R.; Hutchinson, A.; Collins, C.; et al. Structural basis for species-selective targeting of Hsp90 in a pathogenic fungus. *Nat Commun* **2019**, *10*, 402, doi:10.1038/s41467-018-08248-w.

170. Cornet, M.; Gaillardin, C.; Richard, M.L. Deletions of the endocytic components *VPS28* and *VPS32* in *Candida albicans* lead to echinocandin and azole hypersensitivity. *Antimicrob Agents Chemother* **2006**, *50*, 3492-3495, doi:10.1128/aac.00391-06.
171. Garnaud, C.; García-Oliver, E.; Wang, Y.; Maubon, D.; Bailly, S.; Despinasse, Q.; Champlébourg, M.; Govin, J.; Cornet, M. The Rim pathway mediates antifungal tolerance in *Candida albicans* through newly identified Rim101 transcriptional targets, including Hsp90 and Ipt1. *Antimicrob Agents Chemother* **2018**, *62*, e01785-01717, doi:10.1128/aac.01785-17.
172. Iyer, K.R.; Robbins, N.; Cowen, L.E. The role of *Candida albicans* stress response pathways in antifungal tolerance and resistance. *iScience* **2022**, *25*, 103953, doi:10.1016/j.isci.2022.103953.
173. Kim, J.S.; Lee, K.T.; Lee, M.H.; Cheong, E.; Bahn, Y.S. Adenylyl cyclase and protein kinase A play redundant and distinct roles in growth, differentiation, antifungal drug resistance, and pathogenicity of *Candida auris*. *mBio* **2021**, *12*, e0272921, doi:10.1128/mBio.02729-21.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.