
Enhancing Pork Quality: Impact of *Jathropha Tangorensis* and *Psidium guajava* Leaf Meals on Carcass Traits, Physicochemical Properties and Sensory Appeal

Joseph S. Ekpo^{*}, [Utibe M. Okon](#), Opeyemi A. Adediran

Posted Date: 20 February 2025

doi: 10.20944/preprints202502.1561.v1

Keywords: Pork quality; Meat quality; Physicochemical properties; Sustainable Feed Additives; Psidium guajava



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Enhancing Pork Quality: Impact of *Jathropa tangorensis* and *Psidium guajava* Leaf Meals on Carcass Traits, Physicochemical Properties and Sensory Appeal

J. S. EKPO ^{1,*}, U. M. OKON ¹ and O. A. ADEDIRAN ²

¹ Department of Animal Science, Akwa Ibom State University, Akwa Ibom State, Nigeria

² Department of Animal Science, University of Ibadan, Ibadan, Nigeria

* Correspondence: jsekpo2000@gmail.com

Abstract: *Jathropa tangorensis* and *Psidium guajava* are plants of interest containing promising phyto-additives with the potentials to improve pork quality and carcass traits. This study evaluated the effects of *Jathropa tangorensis* and *Psidium guajava* leaf meals on pork quality in terms of sensory attributes, chemical composition, and carcass traits. Twenty-four grower pigs (21.81±2.15 kg) were randomly assigned to three dietary regimens: the basal diet group, which served as the control, the *Jathropa tangorensis* Leaf Meal (JTLM) and *Psidium guajava* Leaf Meal (PGLM) groups. The additives were included at 3% of body weight, with two replicates of four pigs per regimen. Results showed JTLM and PGLM significantly improved carcass traits ($p<0.05$) compared to control. Preslaughter weight, dressed weight, shank, belly, breast, and thick rib chop were higher ($p<0.05$) in JTLM and PGLM groups. Test ingredients also elicited increase in organ weights. Abdominal fat differed significantly ($p<0.05$), with the control group showing the highest value (10.00) compared to JTLM (0.00) and PGLM (0.00). Pork physicochemical properties improved significantly ($p<0.05$) in the JTLM and PGLM groups. Cooking loss, cooking yield, and water holding capacity showed significant differences ($p=0.001$). Taste panelists preferred pork from JTLM and PGLM groups for flavour, juiciness, and tenderness. Conclusively, JTLM and PGLM have good application values as potential feed additives for growing pigs. Dietary supplementation with *Jathropa tangorensis* and *Psidium guajava* at 3% of body weight can improve pork quality including carcass traits, physicochemical properties, and sensory appeal. Higher inclusion levels of the leaf meals is encouraged in future studies.

Keywords: pork quality; meat quality; physicochemical properties; sustainable feed additives; *psidium guajava*

Introduction

The global demand for high-quality pork remains strong because of its significant role as a primary protein source in the human diet [1]. However, modern pork production faces increasing scrutiny regarding its sustainability, environmental impact, and alignment with consumer preferences for healthier, more ethically and naturally produced meat [2,3]. These challenges have prompted the exploration of alternative feed ingredients that can improve pork quality, while supporting environmentally friendly and sustainable practices [4]. One promising avenue is the use of plant-based feed additives from underutilised plants, such as *Jatropha tangorensis* and *Psidium guajava* leaf meal, which are known for their bioactive properties and nutritional benefits [5,6].

Jatropha tangorensis (hospital is too far) and *Psidium guajava* (apple guava), which are rich in phytochemicals, such as flavonoids, phenolic acids, tannins, and other antioxidants, have demonstrated potential for improving livestock growth performance and immune responses [7].

Although their functional attributes are well documented, their specific effects on pork quality, particularly sensory appeal, physicochemical properties, and carcass traits, have received limited attention. Addressing this gap is crucial for optimising pork production in a sustainable and consumer-focused manner. *Psidium guajava* is particularly abundant in polyphenols, including tannins (2-4 %), which exhibit strong antioxidant properties and may contribute to improved meat quality [5,8]. Additionally, it contains vitamin C and carotenoids that are also noted for their antioxidant action [9]. Similarly, *Jatropha tangorensis* is a valuable source of nutrients and bioactive compounds such as tannins, alkaloids, phenols, flavonoids, and saponins [6]. The antioxidative potential of polyphenols has been widely studied [10,11], with evidence suggesting their role in enhancing muscle polyphenol deposition, delaying lipid oxidation, and improving colour, flavour, and overall meat quality. For instance, dietary supplementation with quebracho (*Schinopsis lorentzii*) extract, rich in tannins, increased the total phenolic content of *longissimus dorsi* muscle in lambs [12]. Comparable findings were reported in pigs fed *Vernonia amygdalina* and *Jatropha tangorensis* leaf meal, where significant improvements in growth performance, carcass traits, and meat quality attributes were observed [6,12]. Furthermore, polyphenol supplementation in meat products has been shown to extend shelf life by mitigating lipid oxidation and enhancing colour and flavour stability [14].

Physicochemical properties such as pH, water-holding capacity, and cooking yield are critical determinants of pork quality and consumer acceptance. Both *Jatropha tangorensis* and *Psidium guajava* leaf meals have been shown to contain high levels of antioxidants, which can enhance meat quality by mitigating oxidative stress during production and storage [15]. Additionally, the sensory appeal of pork, which is characterised by attributes such as flavour, tenderness, and juiciness, is a key driver of consumer preference. Despite the known nutritional and bioactive benefits of these plant-based meals, their direct influence on the sensory attributes of pork has remained underexplored.

Carcass traits, including fat distribution, muscle development, and overall yield, are important indicators of pork quality. Previous studies have suggested that dietary supplementation with bioactive plant-based feed additives can influence the balance between fat and muscle deposition, potentially improving carcass composition [7,16]. However, a detailed examination of how *Jatropha tangorensis* and *Psidium guajava* leaf meals affect these traits has not been previously undertaken, leaving a significant knowledge gap. Taking into consideration the many benefits these phyto-additives offer, this study therefore sought to investigate the effects of *Jatropha tangorensis* and *Psidium guajava* leaf meals on the sensory appeal, physicochemical properties, and carcass traits of pork.

Materials and Methods

Study Site, Sourcing and Processing of Test Materials

This experiment was conducted at the Piggery Unit of the Teaching and Research Farm of Akwa Ibom State University, Obio Akpa, Nigeria. *Jatropha tangorensis* and *Psidium guajava* leaves were obtained from Abak town in Akwa Ibom State, Nigeria. The leaves were harvested, thoroughly cleaned under running water, and subsequently chopped with a kitchen knife before being air-dried for two days to mitigate the effect of any antinutritional factors. Upon drying and milling two leaf meals were obtained and designated *Jatropha tangorensis* leaf meal (JTLM) and *Psidium guajava* leaf meal (PGLM).

Experimental Diets and Design

The experimental diets consisted of the control group (basal diet), the JTLM group (basal diet + JTLM), and the PGLM group (basal diet + PGLM). Both leaf meals were incorporated into the daily rations at 3% of the experimental animals' body weights. Feed and water were provided *ad libitum* twice daily (08:00 hours and 16:00 hours). The feeding trial lasted 56 days. The experimental rations were formulated to meet the nutritional needs of the animals in line with the recommendations of NRC.

Experimental Animals, Design and Management

A total of twenty-four [24] young boars (Landrace × Large White) with an average initial weight of 21.81±2.15 kg was acquired from the pig farm of the Department of Animal Science, Akwa Ibom State University, Obio Akpa campus, and used for the experiment. The pigs were randomly distributed in a completely randomized design, with one control treatment and two other diets, all comprising of three treatments. Each treatment group contained eight pigs and was replicated twice with four pigs per replicate.

Carcass Analysis

On day 56 of the feeding trial, carcass traits were determined in two pigs per replicate (12 per treatment) after withdrawing feed 12h prior with the aim of clearing the gut content and reducing the contamination rate. The pigs were slaughtered manually via exsanguination, bleeding, and expulsion of the gut content. The dressed weight obtained was used to calculate the dressing percentage as the ratio of dressed weight to live weight multiplied by 100. Each organ (heart, liver, kidney, and lungs) and abdominal fat were removed and weighed, and weights were expressed as a percentage of the live weight. The carcass was dissected into different primal cuts (Breast, Thick rib chop, Rib chop, Loin, Chump chop, Leg fillet and Leg shank end) and weighed using an electronic scale (AS3101) accordingly. Fresh meat samples obtained from the right thigh were used to determine the meat-to-bone ratio.

Sensory Evaluation

Fresh pork samples (200g) obtained from the left thigh muscle were cooked before being subjected to sensory evaluation using a 9-point hedonic scale (1 = extremely dislike to 9 = like extremely) as described by Okon *et al.* [17]. A 24-member trained taste panel was set up to perform the sensory evaluation. The meat was rated for colour, flavour, tenderness, juiciness, and overall acceptability.

Cooking Loss

This was determined according to the procedure outlined by Jezek *et al.* [18]. The samples were individually wrapped in heat-resistant polythene bags and immersed in boiling water at 75°C for 20 min. After cooking, the meat samples were reweighed, and cooking loss was calculated as the percentage change in weight after cooking.

Thermal Shortening

The length of the meat was measured, and the meat was placed in a heat-resistant polythene bag immersed in boiling water and allowed to cook for 20 min at 80°C. The length of each sample was measured after cooking. The percentage change in the length of meat was taken as thermal loss [19].

Water Holding Capacity

This was determined using the press method [20,21]. A sample of 1 g of meat was placed between two 9 cm Whitman No. 1 filter papers and pressed between plexiglasses using a vice for one minute. The meat sample was removed and oven-dried at 105°C for 24 hours to determine the moisture content. The amount of water released from the meat samples was measured indirectly by measuring the area of filter paper wetted relative to the area of the pressed meat samples. The water-holding capacity was calculated as follows:

$$\text{WHC} = \frac{100 - (AW - AM)}{W_m \times M_c} \times 9.4$$

$$W_m \times M_c$$

Where, Aw = Area of meat samples (cm)
 Wm = weight of meat samples (g)
 Mc = Moisture content of meat samples (%)
 9.47 = a constant factor

Statistical Analysis

Data obtained were subjected to a one-way analysis of variance using GraphPad Prism (version 8.0.2, 268), while the means with significance ($p < 0.05$) were separated using Tukey's test of the same package.

Results and Discussion

Carcass Traits of Pigs Fed *Jathropha Tanjorensis* and *Psidium Guajava* Leaf Meal

The carcass traits of pigs fed *Jathropha tanjorensis* and *Psidium guajava* leaf meal are presented in Table 1. Feeding regimens indicated significant differences ($p < 0.05$) in the pre-slaughter weight, dressed weight, and cut parts including shank, belly, breast, thick rib chop loin, and chump chop. Similarly, all internal organs and abdominal fat were significantly ($p < 0.05$) influenced by the feeding regimen. Pigs fed diets containing JTLM recorded higher values, followed by those in the PGLM group, whereas the CTL group recorded the lowest values across all parameters. Apart from carcass yield being influenced by pre-slaughter weights, which are mostly true impacts of diets [22], *Jathropha tanjorensis* and *Psidium guajava* are rich in antioxidants known to reduce oxidative stress [23], which may result in better nutrient utilization and improved carcass traits obtained in the JTLM and PGLM groups. However, the increase observed in the organ weights could be related to the higher fibre content in the feed served to the pigs in the JTLM and PGLM groups, which engaged the animal's gastrointestinal tract and other organs to work more to digest and utilise the ingested fibre. The results obtained for abdominal fat showed a significant reduction in the JTLM and PGLM groups. This implies that the test ingredients which contained antioxidants such as flavonoids and polyphenols may have increased fat oxidation, regulation of blood sugar levels and reduced fat storage. Our observations are similar to earlier reports on pigs fed phyto-additives [13,24]. Nobre *et al.* [25], in a previous study on lambs reported that feeding 20% of guava leaves did not have any detrimental effect on the carcass.

Table 1. Carcass traits of pigs fed *Jathropha tanjorensis* and *Psidium guajava* leaf meals.

Parameters	Control (CTL)	JTLM	PGLM	SEM	P-values
Pre-slaughter weight (kg)	37.56 ^b	42.10 ^a	39.00 ^b	0.73	0.004
Dressed weight (kg)	20.00 ^b	25.27 ^a	22.00 ^b	0.82	0.002
Dressing (%)	53.24	60.00	56.41	1.95	0.419
Head (kg)	3.00	3.10	3.05	0.17	0.978
Shank (kg)	0.30 ^b	0.70 ^a	0.60 ^a	0.07	0.007
Belly (kg)	1.40 ^b	1.74 ^a	1.64 ^a	0.05	0.001
Breast (kg)	1.30 ^b	1.90 ^a	1.50 ^b	0.09	0.000
Thick rib chop (kg)	1.20 ^c	1.94 ^a	1.62 ^b	0.11	0.000
Rib chop (kg)	1.10 ^b	1.40 ^a	1.20 ^{ab}	0.05	0.027
Loin (kg)	0.86 ^c	1.45 ^a	1.11 ^b	0.09	0.000
Chump chop (kg)	0.90 ^b	1.34 ^a	1.20 ^a	0.07	0.005
Leg fillet end (kg)	1.00	1.30	1.10	0.17	0.819
Leg shank end (kg)	1.20	2.00	1.40	0.21	0.291

Kidney (% LW)	0.35 ^c	0.56 ^b	0.86 ^a	0.08	0.000
Lungs (% L.W)	1.30 ^b	1.80 ^a	1.70 ^a	0.08	0.002
Liver (% L.W)	1.25 ^c	1.45 ^b	1.75 ^a	0.07	0.000
Spleen (% L.W)	0.37 ^b	0.84 ^a	0.54 ^b	0.07	0.003
Heart (% L.W)	0.41 ^b	0.61 ^b	0.91 ^a	0.08	0.000
Abdominal fat (g)	10.00 ^a	0.00 ^b	0.00 ^b	1.67	0.000
Meat to bone ratio	2.00	3.57	2.67	0.37	0.235

^{abc} means on the same row with different superscripts are significantly different ($p < 0.05$) JTLM = *Jathropha tangorensis* leaf meal; PGLM = *Psidium guajava* leaf meal. LW = Live weight.

Physicochemical Properties of Pork from Pigs Fed *Jathropha Tanjorensis* and *Psidium Guajava* Leaf Meal

The physicochemical properties of pork, including pH, water-holding capacity (WHC), and cooking yield (CY), play a crucial role in determining the overall quality and consumer acceptance of pork products [26]. These properties are interconnected and influence various aspects of meat characteristics. Figure 1 presents the physicochemical properties (cooking loss, cooking yield, thermal shortening, and water-holding capacity) of pork from pigs fed diets containing *Jathropha tanjorensis* and *Psidium guajava* leaf meal.




Figure 1. (A, B, C, D): Physicochemical properties of pork from pigs fed diets containing *Jathropha tanjorensis* and *Psidium guajava* leaf meal. Different letters on top of the data bars indicate significant differences ($p < 0.05$) between the mean values. JTLM = *Jathropha tangorensis* leaf meal; PGLM = *Psidium guajava* leaf meal.

Cooking Loss

The use of phyto-additives in animal feed has shown positive effects on meat quality parameters, including cooking loss. As presented in Figure 1A, feeding regimens significantly ($p < 0.05$) reduced cooking loss (CL) in the JTLM and PGLM groups compared with the CTL (control) group. This reduction may be attributed to the presence of antioxidants in the plants. Studies have shown that antioxidants not only reduce oxidative stress and protein degradation during processing but also preserve muscle proteins (myofibrils), thus improving the ability of meat to retain moisture (lower cooking loss) [27–29]. Ayanwale and Aye [30] reported that lower cooking loss is considered an advantage in meat production because it leads to improved tenderness, juiciness, and more favourable meat quality compared to meat with high cooking loss. Generally, fluid loss from meat leads to loss of water-soluble vitamins and nutrients, which can affect juiciness and flavour.

According to a study on lambs, dietary phyto-additive supplementation significantly decreased cooking loss compared to the control group [31].

Cooking Yield

Cooking yield is inversely proportional to cooking loss and important in determining meat's final product texture and economic value [26,32]. Cooking yield is influenced by factors such as the cooking method, temperature, and duration. Higher cooking yields generally result in improved texture, juiciness, and overall meat quality, and are often preferred by consumers. As indicated in Figure 1B, pork in the JTLM and PGLM groups showed significantly ($p < 0.05$) higher CY values than the CTL group. The low cooking loss recorded in these groups, achieved by the application of a low cooking time and temperature, may have been a possible reason for this observation. This is in line with the report by Latoch [32] that marinating meat in fermented dairy products and using sous-vide cooking at lower temperatures can increase tenderness and improve cooking yield. The protein stabilization ability of some of the antinutrients in the tested ingredient, together with their antioxidant properties [30], may also have contributed to the increased cooking yield. The addition of plant-based ingredients significantly decreased cooking loss and increased moisture retention and yield of beef patties, leading to improved texture [28].

Thermal Shortening

Thermal shortening, as illustrated in Figure 1C, did not differ significantly ($p > 0.05$) across feeding regimens. However, lower values were observed in the JTLM and PGLM groups, while the CTL group exhibited the highest value. The high thermal shortening in the CTL group may be attributed to greater moisture loss, leading to muscle shrinkage. Phyto-additives contain antinutritional factors with water-binding properties, which may mitigate moisture loss during heating [33]. As meat proteins denature and contract under thermal processing, water is expelled from the muscle structure, influencing moisture retention and overall meat quality. Comparative studies have evaluated weight loss, moisture content, and water-holding capacity between sous-vide and conventionally cooked meats, demonstrating that sous-vide cooking yields lower cooking losses and higher water-holding capacity, indicative of reduced thermal shortening [19,34]. Additionally, hot-boning combined with sous-vide cooking has been reported to minimise muscle shortening compared to wet-aged meats subjected to conventional cooking methods [19]. These reports further stress the role of pre-cooking muscle state in determining thermal shortening outcomes. Our findings emphasize the influence of these dietary phyto-additives and cooking techniques on muscle integrity and moisture dynamics, which are critical determinants of meat quality.

Water-Holding Capacity

Meat's juiciness during storage and cooking is crucial for consumer's acceptability. This is achievable by maintaining a good water-holding capacity of meat. The feeding regimens in this study showed significant ($p < 0.05$) differences in WHC (Figure 1D). Higher WHC values were observed in the JTLM and PGLM groups than in the CTL (control) group. The low cooking loss values recorded for the JTLM and PGLM groups may have contributed to the observed high WHC values. Additionally, the bioavailability and efficacy of an antinutritional compound (tannins) in the tested materials may have helped stabilize the protein structures of pork, potentially leading to improved water-holding capacity [35]. Water plays a crucial role as a plasticizer and maintains the flexibility of muscle proteins. However, when meat is cooked, protein denaturation occurs, causing water to expel from the myofibrillar lattice structure. As a result, muscle fibre volume decreases, leading to moisture loss, particularly at higher cooking temperatures [36]. Our findings contrast with those of Verma *et al.* [37], who reported that dietary phyto-additive supplementation in sheep nuggets reduced water-holding capacity due to lower emulsion stability, cooking yield, and moisture content, indicating a negative impact on meat quality. Conversely, Joo *et al.* [38] observed that pigs fed conjugated linoleic

acid (CLA)-supplemented diets exhibited lower purge loss, suggesting improved WHC through increased intramuscular fat deposition.

The sensory appeal of pork from pigs fed diets containing *Jathropha tanjorensis* and *Psidium guajava* leaf meals is presented in Figure 2. The results indicated significant differences ($p < 0.05$) in flavour, juiciness, and tenderness between the treatments. Colour and overall acceptability, however, indicated no significant differences ($p > 0.05$). Pork in the PGLM and JTLM groups were preferred by the taste panelists compared to those in the CTL group. *Psidium guajava* and *Jathropha tangorensis* leaves contain phenolic and volatile compounds, such as limonene, flavonoids and tannins which can contribute to the formation of flavour precursors in meat [39]. Our results agree with those of [25], who fed lambs supplemental guava waste and reported no negative change in sensory attributes. Tannins and vitamin C in *Jathropha tangorensis* and *Psidium guajava*, respectively, may also stabilize the protein structure of pork, leading to improved water-holding capacity, tenderness, and juiciness. Our findings align with those of Tran *et al.* [27], Ekpo and Okon [6], and Sahal *et al.* [29], who reported that *Jatropha* and guava leaves positively influence meat quality. The phyto-genic compounds in these plants have been shown to enhance sensory properties and extend the shelf life of pork, thereby improving consumer acceptability.




Figure 2. Sensory appeal of pork from pigs fed diets containing *Jathropha tanjorensis* and *Psidium guajava* leaf meals.

Conclusion

This study revealed that the inclusion of *Jathropha tanjorensis* and *Psidium guajava* leaf meals in pig diets improved meat quality by reducing fat deposition while increasing dressing percentage and primal cuts. The phyto-additives reduced thermal and cooking losses while improving sensory properties such as juiciness, juiciness and tenderness. Based on the results obtained from this study, both *Jathropha tanjorensis* and *Psidium guajava* leaves are recommended for inclusion in pig diets at 3% of body weight to obtain improved accretion of lean meat in pig carcasses, higher carcass yield and better sensory appeal from the resulting meat.

Acknowledgement: The Department of Animal Science, Akwa Ibom State University, is greatly appreciated for providing the experimental animals and essential facilities that facilitated the successful completion of this study.

Conflict of Interest: This study was conducted with no conflicts of interest.

Ethical Approval: The study was approved by the Akwa Ibom State University Ethical Review Board, and all procedures involving human participants adhered to the board's ethical guidelines and standards.

References

1. Nakrachata-Amon, T.; Vorasayan, J.; Pitiruek, K.; Arunyanart, S.; Niyamosoth, T.; Pathumnakul, S. Optimizing vertically integrated pork production supply chain: A Lagrangian heuristic approach. *Heliyon*, **2024**, *10*(6), e26407. <https://doi.org/10.1016/J.HELIYON.2024.E26407>
2. Yang, P.; Yu, M.; Ma, X.; Deng, D. Carbon Footprint of the Pork Product Chain and Recent Advancements in Mitigation Strategies. *Foods (Basel, Switzerland)*, **2023**, *12*(23). <https://doi.org/10.3390/foods12234203>
3. Lin-Schilstra, L.; Backus, G.; Snoek, H.; Mörlein, D. Consumers' view on pork: Consumption motives and production preferences in ten European Union and four non-European Union countries. *Meat Science*, **2022**, *187*, 108736. <https://doi.org/10.1016/J.MEATSCI.2022.108736>
4. Ekpo, J. S.; Okon, U. M. Effects of multiple spices and processing methods on the organoleptic quality of pork. *Acta Periodica Technologica*, **2025**; <https://doi.org/10.2298/APT241022001E>
5. Costa, R. G.; Ribeiro, N. L.; Nobre, P. T.; Carvalho, F. F. R.; Medeiros, A. N.; Cruz, G. R. B.; Freire, L. F. S. Biochemical and hormonal parameters of lambs using guava (*Psidium guajava* L.) agro-industrial waste in the diet. *Tropical Animal Health and Production*, **2018**, *50*(1), 217–221. <https://doi.org/10.1007/s11250-017-1406-5>
6. Ekpo, J. S.; Okon, U. M. Organoleptic Quality, Pork Characterization and Hematological Indices of Growing Pigs Fed Supplemental Diets Containing Bitter Leaf and Hospital Too Far. *AKSU Journal of Agriculture and Food Science*, **2023**, *7*(1), 22–33.
7. Nwachukwu, C. U.; Aliyu, K. I.; Ewuola, E. O. Growth indices, intestinal histomorphology, and blood profile of rabbits fed probiotics- and prebiotics-supplemented diets. *Translational Animal Science*, **2021**, *5*(3). <https://doi.org/10.1093/tas/txab096>
8. Khalifa, I.; Barakat, H. Influencing of Guava Processing Residues Incorporation on Cupcake Characterization. *Journal of Nutrition & Food Sciences*, **2016**, *6*(4). <https://doi.org/10.4172/2155-9600.1000513>
9. Batista, P. F.; Lima, M. A. C.; de Alves, R. E.; Façanha, R. V. Bioactive compounds and antioxidant activity in tropical fruits grown in the lower-middle São Francisco Valley. *Revista Ciência Agronômica*, **2018**, *49*(4). <https://doi.org/10.5935/1806-6690.20180070>
10. Lorenzo, J. M.; Pateiro, M.; Domínguez, R.; Barba, F. J.; Putnik, P.; Kovačević, D. B.; Shpigelman, A.; Granato, D.; Franco, D. Berries extracts as natural antioxidants in meat products: A review. *Food Research International*, **2018**, *106*, 1095–1104. <https://doi.org/10.1016/j.foodres.2017.12.005>
11. Pateiro, M.; Vargas, F. C.; Chinchá, A. A. I. A.; Sant'Ana, A. S.; Strozzi, I.; Rocchetti, G.; Barba, F. J.; Domínguez, R.; Lucini, L.; do Amaral Sobral, P. J.; Lorenzo, J. M. Guarana seed extracts as a useful strategy to extend the shelf life of pork patties: UHPLC-ESI/QTOF phenolic profile and impact on microbial inactivation, lipid and protein oxidation and antioxidant capacity. *Food Research International*, **2018**, *114*, 55–63. <https://doi.org/10.1016/J.FOODRES.2018.07.047>
12. Luciano, G.; Vasta, V.; Monahan, F. J.; López-Andrés, P.; Biondi, L.; Lanza, M.; Priolo, A. Antioxidant status, colour stability and myoglobin resistance to oxidation of longissimus dorsi muscle from lambs fed a tannin-containing diet. *Food Chemistry*, **2011**, *124*(3), 1036–1042. <https://doi.org/10.1016/J.FOODCHEM.2010.07.070>
13. Ekpo, J. S.; Okon, U. M. Performance and lipid profile of growing pigs fed *Vernonia amygdalina* and *Jathropa tanjorensis* leaf meal supplementation, *Livestock Research for Rural Development*, **2022**, <https://www.lrrd.net/search>
14. Vasta, V.; Luciano, G. The effects of dietary consumption of plants secondary compounds on small ruminants' products quality. *Small Ruminant Research*, **2011**, *101*(1–3), 150–159. <https://doi.org/10.1016/J.SMALLRUMRES.2011.09.035>
15. Adeyemi, K. D.; Audu, S.; Oloke, J. A.; Oladiji, O. E.; Salawu, K. F.; Ahmed, R. A.; Sulaimon, R. O. Influence of *Crescentia cujete* and *Launaea taraxacifolia* leaves on growth, immune indices, gut microbiota, blood chemistry, carcass, and meat quality in broiler chickens. *Tropical Animal Health and Production*, **2021**, *53*(3), 365. <https://doi.org/10.1007/s11250-021-02812-1>
16. Wang, Q.; Wang, L.; Li, L.; Sun, M.; Li, P.; Yu, Y.; Zhang, Y.; Xu, Z.; Gao, P.; Ma, J.; Liu, X. Effects of dietary supplementation of fermented *Artemisia argyi* on growth performance, slaughter performance, and meat quality in broilers. *Poultry Science*, **2024**, *103*(4), 103545. <https://doi.org/10.1016/J.PSJ.2024.103545>

17. Okon, U. M.; Ekpo, J. S.; Essien, C. A.; Thabethe, F.; Nuamah, E. Serum lipid profile and organoleptic characteristics of meat from rabbits fed diets containing selim pepper (*Xylopia aethiopica*) and African nutmeg (*Monodora myristica*). *Nigerian Journal of Animal Science*. **2023**, *25*(2), 223-235
18. Ježek, F.; Kameník, J.; Macharáčková, B.; Bogdanovičová, K.; Bednář, J. Cooking of meat: effect on texture, cooking loss and microbiological quality – a review. *Acta Veterinaria Brno*, **2019**, *88*(4), 487–496. <https://doi.org/10.2754/avb201988040487>
19. Xu, B.; Zhang, Q.; Zhang, Y.; Yang, X.; Mao, Y.; Luo, X.; Hopkins, D. L.; Niu, L.; Liang, R. Sous vide cooking improved the physicochemical parameters of hot-boned bovine semimembranosus muscles. *Meat Science*, **2023**, *206*, 109326. <https://doi.org/10.1016/j.meatsci.2023.109326>
20. Irie, M.; Izumo, A.; Mohri, S. Rapid method for determining water-holding capacity in meat using video image analysis and simple formulae. *Meat Science*, **1996**, *42*(1), 95–102. [https://doi.org/10.1016/0309-1740\(95\)00009-7](https://doi.org/10.1016/0309-1740(95)00009-7)
21. Szmańko, T.; Lesiów, T.; Górecka, J. The water-holding capacity of meat: A reference analytical method. *Food Chemistry*, **2021**, *357*, 129727. <https://doi.org/10.1016/j.foodchem.2021.129727>
22. Ekpo, J. S.; Sam, I. M.; Okon, U. M. Performance, Carcass and Relative Organ Weights of Young Boars Fed Graded Dietary Levels of Raw *Icacinia manni* (Earth Ball). *Journal of Agricultural Research and Development*, **2021**, *19*(1). <https://doi.org/10.4314/jard.v19i1.3>
23. Essien, C.; Sam, I. M.; Okon, U. M. *Effect of Jatropha tanjeronsis leaf meal on performance, carcass and internal organ characteristics of broiler finisher chickens*. **2023**, <https://www.researchgate.net/publication/375693443>
24. Pla, M. Effect of nutrition and selection on meat quality proceeding, 8th world rabbit congress held at Puebla, Mexico 7th – 10th September **2004**. Pp: 1337-1348
25. Nobre, P. T.; Munekata, P. E. S.; Costa, R. G.; Carvalho, F. R.; Ribeiro, N. L.; Queiroga, R. C. R. E.; Sousa, S.; da Silva, A. C. R.; Lorenzo, J. M. The impact of dietary supplementation with guava (*Psidium guajava* L.) agroindustrial waste on growth performance and meat quality of lambs. *Meat Science*, **2020**, *164*, 108105. <https://doi.org/10.1016/J.MEATSCI.2020.108105>
26. Watanabe, G.; Motoyama, M.; Nakajima, I.; Sasaki, K. Relationship between water-holding capacity and intramuscular fat content in Japanese commercial pork loin. *Asian-Australasian Journal of Animal Sciences*, **2018**, *31*(6), 914–918. <https://doi.org/10.5713/ajas.17.0640>
27. Tran, T. T. T.; Ton, N. M. N.; Nguyen, T. T.; Le, V. V. M.; Sajeev, D.; Schilling, M. W.; Dinh, T. T. N. Application of natural antioxidant extract from guava leaves (*Psidium guajava* L.) in fresh pork sausage. *Meat Science*, **2020**, *165*, 108106. <https://doi.org/10.1016/J.MEATSCI.2020.108106>
28. Shen, Y.; Hong, S.; Du, Z.; Chao, M.; O'Quinn, T.; Li, Y. Effect of adding modified pea protein as functional extender on the physical and sensory properties of beef patties. *LWT*, **2022**, *154*, 112774. <https://doi.org/10.1016/j.lwt.2021.112774>
29. Sahal, A.; Chaudhary, S.; Hussain, A.; Arora, S.; Dobhal, A.; Ahmad, W.; Kumar, V.; Kumar, S. A comprehensive review on the nutritional composition, bioactive potential, encapsulation techniques, and food system applications of guava (*Psidium guajava* L.) leaves. *Grain & Oil Science and Technology*. **2024**, <https://doi.org/10.1016/J.GAOST.2024.12.003>
30. Ayanwale, B. A.; Aya, V. E. Nutritional Evaluation of Cornflakes Waste in Diets for Broilers. *Pakistan Journal of Nutrition*, **2006**, *5*: 485-489.
31. Zhang, M.; Guo, Y.; Su, R.; Corazzin, M.; Hou, R.; Xie, J.; Zhang, Y.; Zhao, L.; Su, L.; Jin, Y. Transcriptome analysis reveals the molecular regulatory network of muscle development and meat quality in Sunit lamb supplemented with dietary probiotic. *Meat Science*, **2022**, *194*, 108996. <https://doi.org/10.1016/j.meatsci.2022.108996>
32. Latoch, A. Effect of meat marinating in kefir, yoghurt and buttermilk on the texture and color of pork steaks cooked sous-vide. *Annals of Agricultural Sciences*, **2020**, *65*(2), 129–136. <https://doi.org/10.1016/j.aos.2020.07.003>
33. Okon, U. M.; Ekpo, J. S.; Nuamah, E. Effect of *Ficus exasperata* as feed on meat quality of rabbits. *NSAP 2024 Proceedings*, **2024**, 748–752. <https://doi.org/10.51791/njap.vi.5509>

34. Ortuño, J., Mateo, L.; Rodríguez-Estrada, M. T.; Bañón, S. Effects of sous vide vs grilling methods on lamb meat colour and lipid stability during cooking and heated display. *Meat Science*, **2021**, *171*, 108287. <https://doi.org/10.1016/J.MEATSCI.2020.108287>
35. Molino, S.; Pilar Francino, M.; Ángel Rufián Henares, J. Why is it important to understand the nature and chemistry of tannins to exploit their potential as nutraceuticals? *Food Research International*, **2023**, *173*, 113329. <https://doi.org/10.1016/J.FOODRES.2023.113329>
36. Hughes, J. M.; Oiseth, S. K.; Purslow, P. P.; Warner, R. D. A structural approach to understanding the interactions between colour, water-holding capacity and tenderness. *Meat Science*, **2014**, *98*(3), 520–532. <https://doi.org/10.1016/j.meatsci.2014.05.022>
37. Verma, A. K.; Rajkumar, V.; Banerjee, R.; Biswas, S.; Das, A. K. Guava (<italic>Psidium guajava</italic> L.) Powder as an Antioxidant Dietary Fibre in Sheep Meat Nuggets. *Asian-Australasian Journal of Animal Sciences*, **2013**, *26*(6), 886–895. <https://doi.org/10.5713/ajas.2012.12671>
38. Joo, S. T.; Ha, Y. L.; Lee, J. I.; Park, G. B. Effects of dietary conjugated linoleic acid on fatty acid composition, lipid oxidation, color, and water-holding capacity of pork loin. *Journal of Animal Science*, **2002**, *80*(1), 108–112. <https://doi.org/10.2527/2002.801108x>
39. Abd El-Ghany, W. A. The impact of dietary guava (*Psidium guajava* L.) on some livestock production systems. *CABI Reviews*, **2024**, <https://doi.org/10.1079/cabireviews.2024.0018>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.