
The Refined Space–Time Membrane Model: Deterministic Emergence of Quantum Fields and Gravity from Classical Elasticity

[Paul Swann](#)*

Posted Date: 21 January 2026

doi: 10.20944/preprints202503.0736.v7

Keywords: deterministic quantum gravity; spacetime membrane (STM) model; high-order elasticity; emergent gauge symmetry; deterministic decoherence; CKM–PMNS fit; sextic regulator; solitonic black-hole core; tree-level electroweak scattering; hubble-rate tension



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

The Refined Space–Time Membrane Model: Deterministic Emergence of Quantum Fields and Gravity from Classical Elasticity

Paul Swann 

Independent Researcher, UK; officialpaulswann@gmail.com

Abstract

We present a Space–Time Membrane (STM) model in which an eight-parameter, Planck-anchored elasticity equation is explored as a single deterministic framework for emergent quantum-like, gauge-like and gravitational phenomena. The master PDE $\rho\partial_t^2 u + T\nabla^2 u - (E_{STM}(\mu) + \Delta E)\nabla^4 u + \eta\nabla^6 u - \rho\gamma\partial_t u - \lambda u^3 - gu\Psi\Psi = 0$ is fixed by a dimensionless set $\{\rho, T, E_{STM}(\mu), \Delta E, \eta, \lambda, g, \gamma\}$ anchored once to c, G, α and Λ . A bimodal split of u furnishes an effective spinor Ψ ; local internal rephasings then support $U(1) \times SU(2) \times SU(3)$ -type gauge structures as zero-energy wave/anti-wave cycles. Coarse-graining rapid sub-Planck oscillations yields Schrödinger-like envelope dynamics, and we use non-Markovian GKSL master equations to model deterministic decoherence and obtain Born-rule-compatible statistics in representative measurement-style scenarios. In the gravitational sector, the transverse–traceless tensor modes reduce uniquely to an Einstein– Λ limit in the infrared under standard spin-2 bootstrap assumptions and acquire k^4 – k^6 corrections in the ultraviolet; enhanced short-range stiffness replaces singularities with solitonic cores whose microcanonical mode counting, within this STM solitonic-core model, yields a leading-order area-law black-hole entropy with Hawking-like temperature and grey-body factors, and suggests specific deviations in strong-field and ringdown observables. Flavour is treated as a consistency check rather than a full derivation. A calibrated $z = 3$ scalar Functional Renormalisation Group (FRG) analysis (Appendix Y.10), anchored to the STM elastic coefficients, shows that an open set of ultraviolet couplings flows to an effective triple-well scalar potential at a finite infrared scale; the three resulting elastic basins are used as generation-labelled mass scales in the Yukawa sector. A flat-prior scan over STM elastic bands, with no flavour-specific tuning, then reproduces all nine CKM moduli at PDG-2024 precision (primary band) and yields a compact PMNS parameter-space fit at few-unit χ^2 , supported by robustness tests (seed sweeps, ablations, down-scaled draws and a PMNS-target bootstrap). A minimal SMEFT bridge reproduces the tree-level $e^+e^- \rightarrow \mu^+\mu^-$ line shape, including γ - Z interference and leptonic running of $\alpha(s)$. The STM construction realises a $z=3$ Lifshitz-type scaling and, within this class, is super-renormalisable and ghost-free in the UV, admits an Osterwalder–Schrader reconstruction with a spin–statistics theorem on globally hyperbolic backgrounds and anomaly cancellation for the bimodal spinor, and remains stable under small BRST-compatible GKSL deformations. It is directly testable in principle via membrane/metamaterial interferometry, gravitational-wave dispersion and selected collider channels. Overall, in its present calibrated form STM should be viewed as a deterministic elasticity candidate for emergent quantum and gravitational phenomena, offering concrete numerical recipes and experimental benchmarks, and providing a framework that can be systematically tested, extended and, if necessary, falsified.

Keywords: deterministic quantum gravity; spacetime membrane (STM) model; high-order elasticity; emergent gauge symmetry; deterministic decoherence; CKM–PMNS fit; sextic regulator; solitonic black-hole core; tree-level electroweak scattering; hubble-rate tension

Changes from V6 to V7

See ‘Change log.docx’ within Supplementary information.

1. Introduction

Modern physics is built upon two seemingly incompatible foundations: General Relativity (GR) [1–3], which describes gravity through the curvature of spacetime, and Quantum Mechanics (QM) [4–6], whose probabilistic formalism governs microscopic phenomena. Despite remarkable successes within their respective domains, integrating these theories into a coherent framework remains one of contemporary physics' most pressing challenges. Existing approaches—such as String Theory's extra-dimensional constructions and Loop Quantum Gravity's discretised spin-network formalism—provide valuable insights but have yet to deliver a definitive resolution of quantum gravity [7,8]. Meanwhile, enduring puzzles such as the black-hole information paradox and the cosmological-constant problem underline fundamental tensions between GR's smooth geometry and QM's intrinsic randomness [9–11].

The Space–Time Membrane (STM) model proposes spacetime as a four-dimensional elastic membrane interacting with a parallel mirror domain. Every particle excitation on our “face” of the membrane has a corresponding mirror particle on the opposite face, ensuring exact matter–antimatter symmetry at the fundamental level and offering a possible route to addressing the observed baryon asymmetry. The membrane's elastic dynamics simultaneously generate gravitational curvature and quantum-like phenomena: rather than postulating intrinsic randomness, apparent quantum probabilism emerges as a deterministic consequence of chaotic, sub-Planck elastic oscillations.

Gravity in STM. Transverse–traceless spin-2 excitations follow

$$\omega^2 = c_g^2 k^2 + \frac{\alpha_2}{\Lambda_g^2} k^4 + \frac{\alpha_3}{\Lambda_g^4} k^6,$$

with the identification $c_g^2 = T/\rho$, $\alpha_2/\Lambda_g^2 = K_4/\rho$, $\alpha_3/\Lambda_g^4 = \eta/\rho$. Time derivatives remain strictly second order. In the IR, the spin-2 bootstrap reproduces Einstein– Λ ; phenomenologically we obtain broadband inspiral phase shifts (f^2, f^4), ringdown frequency shifts $(\omega/\Lambda_g)^{2,4}$, and short-range Yukawa-type static corrections (symbols catalogued in Appendix Z).

UV control and EFT placement. Alongside these IR phenomenological aims, the refined STM model now has a complete UV control statement. In flat space and with vanishing open-system terms the quadratic Hamiltonian is positive and involves only $\dot{\phi}^2$, so no Ostrogradsky ghosts arise; the dominant sextic elasticity implies Lifshitz scaling with $z = 3$, under which the conservative interacting theory is super-renormalisable by power counting. Euclidean continuation yields reflection positivity for gauge-invariant composites and, via standard OS reconstruction, a positive spectral density. Finally, we state and use a decoupling/matching bound to Yang–Mills below a finite scale, which underwrites the SMEFT mapping employed later in the paper.

Concretely, the displacement field $u(x, t)$ is decomposed into two complementary oscillatory modes that combine into a two-component spinor $\Psi(x, t)$. Mode-by-mode interactions between each spinor component and its mirror antispinor redistribute energy—attractive interactions generate localised curvature (gravity), while repulsive or cancelling interactions reinject energy into the membrane background. Composite photons arise as coherent wave–anti-wave cycles. In the virtual (near-field) regime the energy exchanged in one half-cycle is returned in the next, so the period-averaged flux vanishes. By contrast, when the mode is on-shell—e.g. in an annihilation producing two photons—the same mechanism yields propagating gauge-like waves with positive cycle-averaged flux that carry away energy–momentum; the associated reduction in local strain/curvature equals the radiated energy.

When rapid sub-Planck oscillations in u are coarse-grained, a slowly varying envelope ψ emerges that obeys an effective Schrödinger-like equation. This envelope reproduces interference patterns and apparent wavefunction collapse, recasting standard quantum phenomena (including the Born rule) as manifestations of deterministic chaos.

Determinism and apparent underdetermination. The underlying STM PDE is well-posed and deterministic given complete initial and boundary data; any apparent underdetermination enters only

after coarse-graining, because the envelope/detection map is many-to-one and discards sub-Planck phase detail.

In this interpretation, Feynman's path integral is re-expressed as a stationary-phase approximation to a single underlying wave field, rather than a literal sum over independently realised histories. The familiar kernel $K(x_b, t_b; x_a, t_a)$ then appears as the WKB/multiple-scale approximation to STM's Green function (Appendix D), providing a concrete, if still approximate, link between the membrane PDE and standard quantum-mechanical propagators.

The STM framework further reinterprets key aspects of particle physics. Electroweak symmetry breaking arises from rapid zitterbewegung-like interactions between spinors and mirror antispinors, generating W^\pm and Z^0 masses and yielding CP-violating phases without invoking extra scalar fields. A bimodal spinor decomposition underpins emergent gauge symmetries—U(1), SU(2) and SU(3)—as deterministic elastic connections.

The model incorporates:

- Scale-dependent elastic parameters and higher-order spatial derivatives (notably ∇^6) to regulate ultraviolet divergences.
- Non-Markovian memory kernels to explain deterministic decoherence and effective wavefunction collapse.
- A precise bimodal decomposition of u into a two-component spinor Ψ , yielding emergent gauge bosons.
- A deterministic electroweak symmetry-breaking mechanism via cross-membrane oscillations.
- A multi-loop renormalisation-group analysis and a nonperturbative scalar FRG study, revealing triple-well vacuum structure and elastic basins that can be used to model three fermion generations.

In the gravitational sector, linearised strain fields u_μ link directly to metric perturbations $h_{\mu\nu}$, yielding Einstein-like field equations from the STM action—even when including damping and scale-dependent couplings (Appendix M). A detailed multi-scale derivation (Appendix H) shows that coarse-grained sub-Planck oscillations produce a near-constant vacuum offset acting as dark energy [12,13], and that a mild late-time evolution in stiffness or damping could address the Hubble tension [14].

Crucially, Section 2.10—and the full parameter table in Appendix K.7—now fixes every STM coefficient to physical constants:

$$T = \rho c^2 \simeq 4.82 \times 10^{42} \text{ Pa}, \quad E_{\text{STM}}(\mu) = \frac{c^4}{8\pi G}, \quad g = \sqrt{4\pi\alpha} \simeq 0.3028,$$

$$\lambda_{\text{nd}} = 0.13,$$

together with the vacuum-stiffness offset $\langle \Delta E \rangle \simeq 6.8 \times 10^{-10} \text{ J m}^{-3}$, a macroscopic damping coefficient $\gamma_{\text{nd}} = 0.01$ (corresponding to $\gamma_{\text{phys}} \simeq 1.85 \times 10^{41} \text{ s}^{-1}$), and the sextic regulator $\eta_{\text{nd}} = 0.02$. These calibrations anchor the model quantitatively to the fundamental constants c , G , α and Λ , leaving no free elastic or damping parameters.

Although STM now captures both quantum-field and cosmological-scale phenomena within one PDE, several frontiers remain.

On the thermodynamics front, we have:

- Recovered the leading Bekenstein–Hawking area law via micro-canonical mode counting in the STM solitonic core (Appendix F.4), including estimates of finite-size corrections;
- Calculated grey-body transmission factors and effective horizon temperatures via fluctuation-dissipation (Appendix G.4–G.5);
- Sketched a Euclidean path-integral approach to the evaporation law, matching the leading-order M^3 timescale (Appendix H). Remaining thermodynamic tasks include subleading logarithmic

and power-law corrections to the area law, Page-curve tests of unitarity and detailed first-law verifications (Appendix F.7).

Beyond thermodynamics, our analytic derivations (Appendices C and N) detail mode-by-mode spinor–antispinor couplings, while recent numerics anchored to physically motivated parameters reveal three well-separated mass minima that reproduce the Standard Model’s generational hierarchy, mixing angles and CP-violating phases (Sections 3.1.4 & 4.3). Early tests (Section 3.3; Appendix K.7) suggested the damping coefficient γ might be dispensable, but a full analysis of measurement dynamics and deterministic wavefunction collapse (Section 3.4) confirms that a finite γ is essential. Although γ introduces mild non-conservatism, the model remains stable across a broad range of values.

Appendix T proves global well-posedness, self-adjointness and ghost-freedom, while Appendix U shows anomaly cancellation via mirror doubling. With these foundations in place, Appendix O now completes the spin–statistics link for STM’s bimodal spinor on globally-hyperbolic backgrounds; remaining frontiers are higher-loop renormalisation and the microstate structure of black holes.

Addressing the remaining challenges will be crucial to establishing the STM framework’s consistency across all scales.

Unlike many quantum-gravity schemes, the STM model is rooted in classical continuum elasticity, so it can be tested directly through numerical simulations and laboratory analogues such as metamaterials. By deriving Schrödinger dynamics, the Born rule, gauge symmetries and CP violation from a single deterministic PDE, STM uses far fewer independent postulates than frameworks of comparable scope—for example, the Standard Model plus general relativity, which require separate fundamental fields and symmetry assumptions. Approaches with an even sparser axiomatic core (e.g. asymptotically safe gravity) typically focus on the gravitational sector alone and do not yet reproduce the full gauge and flavour structure that STM aims to encompass.

We therefore encourage further numerical, experimental and theoretical exploration of the STM model as a promising, conceptually transparent route to reconciling quantum phenomena with gravitational curvature.

Organisation of the Paper

- **Section 2 (Methods)** provides a detailed overview of the STM wave equation, including explicit derivations of higher-order elasticity terms, spinor construction, scale-dependent parameters, and the deterministic interpretation of decoherence.
- **Section 3 (Results)** demonstrates how quantum-like dynamics, the Born rule, entanglement analogues, emergent gauge fields ($U(1)$, $SU(2)$, $SU(3)$), deterministic decoherence, fermion generations, and CP violation naturally arise from the deterministic membrane equations.
- **Section 4 (Discussion)** explores the broader implications of these findings, along with possible experimental tests and numerical simulations.
- **Section 5 (Conclusion)** summarises the key theoretical advances, outstanding issues, and potential future directions, including proposals aimed at verifying the STM model’s predictions.

Appendices A–Y (together with Appendix Z) comprehensively present supporting details, derivations, and numerical methods. They address:

- Operator Formalism and Spinor Field Construction (Appendix A)
- Derivation of the STM Elastic-Wave Equation and External Force (Appendix B)
- Gauge symmetry emergence and CP violation (Appendix C)
- Coarse-grained Schrödinger-like dynamics (Appendix D)
- Deterministic entanglement analogues (Appendix E)
- Singularity avoidance (Appendix F)
- Non-Markovian Decoherence and Measurement (Appendix G)
- Vacuum energy dynamics and the cosmological constant (Appendix H)
- Proposed experimental tests (Appendix I)

- Renormalisation Group Analysis and Scale-Dependent Couplings (Appendix J)
- Finite-Element Calibration of STM Coupling Constants (Appendix K)
- Nonperturbative analyses revealing solitonic structures (Appendix L)
- Covariant Generalisation and Derivation of Einstein Field Equations (Appendix M)
- Emergent Scalar Degree of Freedom from Spinor–Mirror Spinor Interactions (Appendix N)
- Spin–Statistics in the STM Framework (Appendix O)
- Reconciling Damping, Environmental Couplings, and Quantum Consistency in the STM Framework (Appendix P)
- Toy Model PDE Simulations (Appendix Q)
- CKM and PMNS flavour fits from an STM elastic template (Appendix R)
- STM Scattering Amplitude Validation (Appendix S)
- Well-Posedness and Ghost-Freedom of the STM PDE (Appendix T)
- Anomaly Cancellation in the STM Model (Appendix U)
- Effective Field Theory and Renormalisation Match for STM (Appendix V)
- Full SM-Gauge EFT (dim-6) and One-Loop RG (Appendix W)

The corresponding SMEFT bridge is completed by one-loop closed-sector running and operator mixing (Appendix W), while Appendix X supplies the causality and Einstein– Λ bootstrap conditions used for the IR spin-2 limit.

- Causality (Non-Markovian) and Einstein– Λ Bootstrap (Appendix X)
- UV structure, scaling and renormalisation, including a $z = 3$ scalar FRG proxy of the STM scalar sector and its robust triple-well region, together with a trimmed basin-field FRG interface that motivates (and specifies a reproducible future data-export format for) the three-basin flavour ansatz used in the CKM/PMNS fits (Appendix Y).

Finally, Appendix Z serves as a Glossary of Symbols, ensuring clarity and consistency of notation throughout.

Scope and level of claim. It is important to stress what the present work does and does not establish. We do not claim a mathematically complete derivation of the Standard Model or a fully rigorous proof that quantum theory is redundant. Rather, starting from a single elasticity PDE we identify a set of consistent emergent structures — propagators, Schrödinger-type envelopes, gauge-like connections and flavour-mixing templates — and check them against selected experimental benchmarks. Within this scope we demonstrate a super-renormalisable elasticity field theory with Osterwalder–Schrader reconstruction and spin–statistics, an Einstein– Λ limit for the transverse–traceless spin-2 modes, a $z = 3$ scalar FRG analysis that identifies a robust triple-well region in effective coupling space and motivates the three-basin elastic template used in our CKM/PMNS fits, and a minimal SMEFT bridge to collider observables, including operator normalisation, one-loop running and operator-mixing blocks in the closed sector (Appendices V–W; conventions in Appendix Z). Several constructions (for example the solitonic FRG models of Appendix L and the non-Markovian master equations) are currently implemented in deliberately simplified truncations and geometries; the $z = 3$ scalar FRG analysis in Appendix Y.10, while still a single-field LPA/LPA' truncation, is directly constrained by the STM elastic coefficients and provides a structurally constrained theoretical motivation for the triple-well basin picture used in our phenomenological flavour template (Appendix R); a full multi-field STM FRG derivation of the absolute mass spectrum is left to future work.

In this manuscript we therefore do not attempt a fully closed, FRG-anchored derivation of all Standard Model parameters; instead we present a single calibrated STM benchmark, test its elastic, gravitational and flavour templates against selected CKM/PMNS and scattering observables, and specify the FRG export contract required to turn this benchmark into a predictive pipeline.

Once the elastic and damping coefficients are fixed in terms of physical constants, the same calibrated parameter set underlies the flavour fits, gravitational sector, EFT matching and proposed experiments, so that the present STM truncation already yields definite, falsifiable predictions rather than a family of arbitrarily tuned models.

While non-trivial and quantitatively constrained in several sectors, the results should only be read as evidence that a deterministic membrane framework can plausibly reproduce key quantum and gravitational features, rather than as a final, fully complete or mathematically unique theory.

Throughout this manuscript the term “toy model” is reserved for specific illustrative constructions, such as the non-calibrated FRG-plus-soliton example of Appendix L; the calibrated STM benchmark itself is intended as a candidate world-theory, not a toy model.

2. Methods

In the Space–Time Membrane (STM) model, spacetime is represented as a four-dimensional elastic membrane governed by a deterministic high-order partial differential equation. This single PDE unifies gravitational-scale curvature with quantum-like oscillations by incorporating higher-order elasticity, scale-dependent stiffness, non-linear terms, and possible non-Markovian effects. Below, we provide the theoretical foundations, outline the operator quantisation that yields quantum-like behaviour, show how gauge fields naturally emerge, discuss renormalisation strategies, and comment on the classical limit.

2.1. Classical Framework and Lagrangian

2.1.1. Displacement Field and Equation of Motion

We model the space–time membrane by a scalar displacement field $u(\mathbf{x}, t)$, which encodes local departures from equilibrium. The canonical equation of motion is constructed so that each term has the dimensions of force per unit volume (N m^{-3}). Higher-order elastic coefficients carry explicit powers of the Planck length $\downarrow_P = \sqrt{\hbar G/c^3}$ where needed to keep SI units manifest. This equation is the master PDE from which all subsequent constructions are derived.

Canonical STM PDE (compact form).

The compact form derived in Appendix B is

$$\rho \frac{d^2 u}{dt^2} + T \nabla^2 u - (E_{STM}(\mu) + \Delta E(\mathbf{x}, t; \mu)) \nabla^4 u + \eta \nabla^6 u - \rho \gamma \partial_t u - \lambda u^3 - g u \bar{\Psi} \Psi + F_{ext}(\mathbf{x}, t) = 0.$$

Here $\Delta E(\mathbf{x}, t; \mu)$ collects the spatially uniform vacuum offset $\Delta E(\mu)$ and any zero-mean local fluctuation $\delta E(\mathbf{x}, t; \mu)$. When fluctuations are not under discussion we write $\Delta E(\mathbf{x}, t; \mu) \rightarrow \Delta E(\mu)$ and may absorb it into $E_{STM}(\mu)$. This expression suppresses explicit \downarrow_P factors, which are absorbed into the higher-order moduli; it is the reference form used throughout the appendices.

The damping parameter γ is the mass-normalised rate (s^{-1}); the corresponding SI force-density coefficient is $\gamma_1 \equiv \rho \gamma$, which appears in the SI PDE as $-\rho \gamma \partial_t u$.

Explicit SI force-density form.

Restoring \downarrow_P to make units manifest gives

$$\rho \frac{d^2 u}{dt^2} + T \nabla^2 u - (E_{STM}(\mu) + \Delta E(\mathbf{x}, t; \mu)) \downarrow_P^2 \nabla^4 u + \eta \downarrow_P^4 \nabla^6 u - \rho \gamma \partial_t u - \lambda u^3 - g u \bar{\Psi} \Psi + F_{ext}(\mathbf{x}, t) = 0.$$

Mass-normalised SI form.

Dividing by ρ yields the form used for non-dimensionalisation and numerics:

$$\frac{d^2 u}{dt^2} + c_T^2 \nabla^2 u - \tilde{E}(\mathbf{x}, t; \mu) \nabla^4 u + \tilde{\eta} \nabla^6 u - \gamma \partial_t u - \tilde{\lambda} u^3 - \tilde{g} u \bar{\Psi} \Psi + \tilde{F}_{ext} = 0,$$

with

$$c_T^2 = \frac{T}{\rho}, \tilde{E}(\mathbf{x}, t; \mu) = \frac{(E_{STM}(\mu) + \Delta E(\mathbf{x}, t; \mu)) \downarrow_P^2}{\rho}, \tilde{\eta} = \frac{\eta \downarrow_P^4}{\rho}, \tilde{\lambda} = \frac{\lambda}{\rho}, \tilde{g} = \frac{g}{\rho}, \tilde{F}_{ext} = \frac{F_{ext}}{\rho},$$

and $\delta\tilde{E}(\mathbf{x}, t) = \delta E(\mathbf{x}, t)\uparrow_P^2/\rho$ when fluctuations are retained. This is the form actually used for non-dimensionalisation and in the numerical solvers (Appendices K.6–K.7).

EFT sign-map.

The spectral identifications

$$\nabla^2 \mapsto -k^2, \nabla^4 \mapsto +k^4, \nabla^6 \mapsto -k^6 \text{ (equivalently } \nabla^{2n} \mapsto (-1)^n k^{2n})$$

are carried into the operator normalisations in Appendix V and Appendix W, so the signs of the derivative operators and the Wilson-coefficient conventions are consistent across the PDE, the emergent Lagrangian and the SMEFT translation (Appendix V §§V.1–V.2, Appendix W §W.2).

In what follows we use the compact (canonical) form; SI units are recovered by reinstating \uparrow_P so that the ∇^4 and ∇^6 terms read $Q_{SI}\uparrow_P^2\nabla^4 u$ and $S_{SI}\uparrow_P^4\nabla^6 u$, with mass-normalised coefficients obtained by dividing by ρ . The UV/renormalisation summary for the conservative sector appears in §2.4, with proofs in Appendix Y.

For gravitational notation we use c_g , $\alpha_{2,3}$ and Λ_g as defined in Appendix Z and mapped to $\{T, \rho, E_{STM}, \eta\}$ using §2.11 (tensor-mode EFT map) together with §2.1.1 (signs/units) and the conversion tables in Appendices K.6–K.7; the IR Einstein– Λ completion is summarised in Appendix M.

Parameter summary.

For ease of reference we collect the main symbols, their roles, and the linear dispersion relation here.

Notation. E_{STM} is the baseline quartic stiffness, ΔE the spatially uniform vacuum offset, and $\delta E(\mathbf{x}, t)$ zero-mean local fluctuations. Tildes ($\tilde{E}, \tilde{\eta}, \tilde{\lambda}, \tilde{g}, \tilde{F}_{ext}$) denote mass-normalised coefficients. We write \uparrow_P for the Planck length, ∇ for spatial gradients, and ∂_t for time derivatives. The emergent wave speed satisfies $c_T^2 = T/\rho$.

Key ingredients.

- ρ : inertial density, fixed by matching dispersion to relativistic propagation.
- T : baseline tension, defines the emergent light-cone speed c_T .
- E_{STM} : quartic stiffness, linked to the Newtonian gravitational sector.
- ΔE : vacuum offset, matched to the observed dark-energy density.
- η : sextic regulator, set by the Planck-scale UV cut-off.
- γ (s^{-1} ; damping rate): in SI force-density form the PDE carries $-\rho\gamma\partial_t u$; after mass-normalisation this is $-\gamma\partial_t u$. We distinguish a Planck-stage calibration rate γ_{UV} (used only to set non-dimensional scales) from the laboratory/environmental rate γ_{env} obtained by ring-down of the same mode (if $A(t) \propto e^{-t/\tau_{amp}}$ then $\gamma_{env} \approx 2/\tau_{amp}$). Unless stated otherwise, γ in figures, fits and phenomenology means γ_{env} . The SI force-density coefficient is $\gamma_1 \equiv \rho\gamma_{env}$.
- λ : cubic nonlinearity, Higgs-like self-interaction.
- g : Yukawa/gauge coupling to spinor bilinears.

While not appearing explicitly in the scalar PDE, the spinor sector carries a milder dephasing γ_f that enters the Dirac-like equations for the spinor and mirror-spinor fields. Coarse-graining ties it to the scalar damping so that flavour decoherence occurs on the same physical timescale as scalar Born-rule collapse:

$$\gamma_f = \frac{1}{2}\gamma_{env}$$

(see §3.4.1 and Appendix K.6).

Dispersion (linear, mass-normalised).

For plane-wave solutions $u \sim e^{i(\mathbf{k}\cdot\mathbf{x}-\omega t)}$ and the rule $\nabla^{2n} \rightarrow (-1)^n k^{2n}$, the linearised dispersion relation is

$$\omega^2 - i\gamma\omega + c_T^2 k^2 + \tilde{E}k^4 + \tilde{\eta}k^6 = 0.$$

For $\gamma > 0$ we have $Im\omega = -\gamma/2 < 0$: exponential decay, as expected for physical damping. With $T, E_{STM}, \eta > 0$, the quadratic energy is positive and the linear operator is sectorial (Appendix T), ensuring well-posedness and the absence of higher-derivative ghosts in the second-order-in-time formulation.

This PDE therefore provides a unified mathematical context in which large-scale curvature emerges as low-frequency deformations and short-scale oscillations mimic quantum phenomena—without extra dimensions or intrinsic randomness.

2.1.2. Lagrangian Density

The elastic dynamics follow from a local Lagrangian density $\mathcal{L}(u, \partial_t u, \nabla u, \dots)$. Damping and external work are handled via a Rayleigh dissipation functional and an explicit forcing term, rather than being placed inside the conservative Lagrangian. As in §2.1.1, the compact (canonical) form suppresses explicit powers of the Planck length \downarrow_P ; the SI-expanded form reinstates them so that every Euler–Lagrange term has units of force per unit volume ($N m^{-3}$).

Canonical (compact) Lagrangian density.

$$\mathcal{L} = \frac{1}{2}\rho(\partial_t u)^2 - \frac{1}{2}T |\nabla u|^2 \frac{1}{2}(E_{STM}(\mu) + \Delta E(\mathbf{x}, t; \mu))(\nabla^2 u)^2 - \frac{1}{2}\eta |\nabla^3 u|^2 - \frac{\lambda}{4}u^4 + \frac{\mathcal{G}}{2}u^2\bar{\Psi}\Psi$$

Variation of the action built from \mathcal{L} reproduces the conservative part of the canonical STM PDE in §2.1.1 (i.e. without damping and F_{ext}).

Explicit SI form (higher-order pieces with \downarrow_P).

Writing the quartic and sextic operators with explicit Planck-length factors,

$$\mathcal{L} = \frac{1}{2}\rho(\partial_t u)^2 - \frac{1}{2}T |\nabla u|^2 \frac{1}{2}(E_{STM} + \Delta E)\downarrow_P^2(\nabla^2 u)^2 - \frac{1}{2}\eta\downarrow_P^4 |\nabla^3 u|^2 - \frac{\lambda}{4}u^4 + \frac{\mathcal{G}}{2}u^2\bar{\Psi}\Psi$$

For covariant discussions we use a 3 + 1 split and write the quartic term in terms of the spatial Laplace–Beltrami operator Δ_Σ ; time evolution remains second order, while higher-order operators act on space alone (Appendix T). With the \downarrow_P^2 and \downarrow_P^4 insertions, each Euler–Lagrange term carries SI units $N m^{-3}$ (Appendix K.6), matching the force-density PDE in §2.1.1.

Rayleigh dissipation (damping).

Damping is introduced via a Rayleigh functional

$$\mathcal{R} = \frac{1}{2}\rho\gamma(\partial_t u)^2.$$

Here γ is the mass-normalised rate (s^{-1}); the corresponding SI force-density coefficient is $\gamma_1 \equiv \rho\gamma$, which appears in the SI PDE as $-\rho\gamma\partial_t u$. In the Euler–Lagrange–Rayleigh equation, $\partial\mathcal{R}/\partial(\partial_t u) = \rho\gamma\partial_t u$ enters on the right-hand side; moving it to the left reproduces precisely the $-\rho\gamma\partial_t u$ term in §2.1.1.

Mass-normalised density (solver form).

Dividing by ρ gives the mass-normalised Lagrangian density

$$\tilde{\mathcal{L}} = \frac{1}{2}(\partial_t u)^2 - \frac{1}{2}c_T^2 |\nabla u|^2 + \frac{1}{2}\tilde{E}(\nabla^2 u)^2 - \frac{1}{2}\tilde{\eta} |\nabla^3 u|^2 - \frac{\tilde{\lambda}}{4}u^4 + \frac{\tilde{\mathcal{G}}}{2}u^2\bar{\Psi}\Psi,$$

with

$$c_T^2 = \frac{T}{\rho}, \quad \tilde{E} = \frac{(E_{STM} + \Delta E)\downarrow_P^2}{\rho}, \quad \tilde{\eta} = \frac{\eta\downarrow_P^4}{\rho}, \quad \tilde{\lambda} = \frac{\lambda}{\rho}, \quad \tilde{\mathcal{G}} = \frac{\mathcal{G}}{\rho}.$$

In this normalisation, damping corresponds to

$$\tilde{\mathcal{R}} = \frac{1}{2}\gamma(\partial_t u)^2,$$

and the Euler–Lagrange–Rayleigh equation yields the mass-normalised PDE $-\gamma\partial_t u$ in place of $-\rho\gamma\partial_t u$, in agreement with §2.1.1 and Appendices B, K, P, Q and T. This mass-normalised form is the starting point for the non-dimensionalisation used in Appendices K.6–K.7.

Assumptions for linear analysis.

For the linear theory we assume

$$u \in H^3(\mathbb{R}^3) \text{ (or } H^3(\Omega) \text{ with Dirichlet/Neumann b.c.)}, \rho, T, E_{STM}, \eta > 0, \gamma \geq 0,$$

and the nonlinear terms are locally Lipschitz on H^3 . These conditions match those used in Appendix T (Thm T.1, Prop. T.2) and underpin the well-posedness and ghost-freedom statements quoted in §2.4.

2.1.3. Hamiltonian Formulation and Poisson Brackets

The Hamiltonian density follows from a Legendre transform of the conservative Lagrangian; Rayleigh damping and external work enter only as generalised forces and do not contribute to the Hamiltonian.

Canonical momentum.

$$\pi(x, t) = \frac{\partial \mathcal{L}}{\partial(\partial_t u)} = \rho \partial_t u.$$

Hamiltonian density.

$$H = \frac{1}{2}\rho(\partial_t u)^2 + \frac{1}{2}T |\nabla u|^2 + \frac{1}{2}(E_{STM}(\mu) + \Delta E(x, t; \mu))(\nabla^2 u)^2 + \frac{1}{2}\eta |\nabla^3 u|^2 + \frac{\lambda}{4}u^4 + \frac{g}{2}u^2\bar{\psi}\psi$$

(In SI-expanded form, insert \uparrow_p^2 and \uparrow_p^4 on the ∇^4 and ∇^6 pieces exactly as in § 2.1.2.)

Canonical brackets.

$$\{u(x), \pi(y)\} = \delta^{(3)}(x - y), \quad \{u(x), u(y)\} = \{\pi(x), \pi(y)\} = 0.$$

Hamilton's equations, $\partial_t u = \delta\mathcal{H}/\delta\pi$, $\partial_t \pi = -\delta\mathcal{H}/\delta u$, reproduce the conservative STM dynamics; energy decay arises solely from the Rayleigh term:

$$\dot{E}(t) = - \int \rho \gamma (\partial_t u)^2 d^3x \leq 0.$$

A rigorous curved-spacetime proof of self-adjointness, ghost-freedom and global well-posedness is given in Appendix T (Thm T.1; Prop. T.2). The BRST-compatible open-system implementation of damping appears in Appendices P and T (Thm T.6).

Energy flux (Noether). For spatial translations, the conservative energy density \mathcal{H} and flux S_i obey the local continuity law

$$\partial_t \mathcal{H} + \partial_i S_i = 0 \quad (\text{no damping/forcing, constant coefficients}).$$

A convenient symmetric representative, consistent with the elastic operator $T\nabla^2 - (E_{STM} + \Delta E)\nabla^4 + \eta\nabla^6$, is

$$\mathcal{H} = \frac{1}{2}\rho u_t^2 + \frac{1}{2}T |\nabla u|^2 + \frac{1}{2}(E_{STM} + \Delta E)(\nabla^2 u)^2 + \frac{1}{2}\eta |\nabla^3 u|^2 + V(u),$$

$$S_i = -Tu_t\partial_i u + (E_{STM} + \Delta E)[u_t\partial_i(\nabla^2 u) - (\nabla^2 u)\partial_i u_t] - \eta[u_t\partial_i(\nabla^4 u) - (\nabla^4 u)\partial_i u_t]$$

This S_i closes the continuity equation for the conservative field equation

$$\rho u_{tt} + T\nabla^2 u - (E_{STM} + \Delta E)\nabla^4 u + \eta\nabla^6 u - V'(u) = 0.$$

Units. $[S_i] = Jm^{-2}s^{-1} = Wm^{-2} = N(ms)^{-1}$.

Mass-normalised flux

Dividing by ρ and writing $c_T^2 := T/\rho$, $\tilde{E} := (E_{STM} + \Delta E)/\rho$, $\tilde{\eta} := \eta/\rho$,

$$\tilde{S}_i = -c_T^2 u_t \partial_i u + \tilde{E}[u_t \partial_i(\nabla^2 u) - (\nabla^2 u) \partial_i u_t] - \tilde{\eta}[u_t \partial_i(\nabla^4 u) - (\nabla^4 u) \partial_i u_t]$$

with $[\tilde{S}_i] = m^3 s^{-3}$ (specific energy flux).

With damping and forcing

Including Rayleigh damping and external work, the local balance becomes

$$\partial_t \mathcal{H} + \partial_i S_i = -\rho\gamma u_t^2 + F_{ext} u_t,$$

and the control-volume balance over V is

$$\frac{d}{dt} \int_V \mathcal{H} d^3x = - \oint_{\partial V} S \cdot dA - \rho\gamma \int_V u_t^2 d^3x + \int_V F_{ext} u_t d^3x.$$

If ΔE varies in space-time

When $\Delta E = \Delta E(x, t)$ is not constant, spatial translation invariance is broken by the explicit coefficient dependence, and the conservative continuity law acquires source/sink terms:

$$\partial_t \mathcal{H} + \partial_i S_i = \frac{1}{2}(\partial_t \Delta E)(\nabla^2 u)^2 + \frac{1}{2} \nabla \Delta E \cdot \nabla [(\nabla^2 u)^2] \quad (\text{no damping/forcing}).$$

Thus strict conservation holds whenever ΔE is constant (or at least $\partial_t \Delta E = 0$ and $\nabla \Delta E = 0$).

2.1.4. Conjugate Momentum and Dispersion

Conjugate momentum. From the conservative Lagrangian density of §2.1.2 the canonical momentum is

$$\pi(\mathbf{x}, t) = \frac{\partial \mathcal{L}}{\partial(\partial_t u)} = \rho \partial_t u.$$

Rayleigh damping contributes only as a generalised force; it is not part of the conservative Hamiltonian.

Mass-normalised linear PDE. Dropping nonlinearity, spinor coupling and forcing, and dividing the SI equation by ρ , the linearised STM equation with physical damping reads

$$\partial_t^2 u + c_T^2 \nabla^2 u - \tilde{E} \nabla^4 u + \tilde{\eta} \nabla^6 u - \gamma \partial_t u = 0,$$

With

$$c_T^2 = \frac{T}{\rho}, \quad \tilde{E} = \frac{(E_{STM} + \Delta E) \downarrow_P^2}{\rho}, \quad \tilde{\eta} = \frac{\eta \downarrow_P^4}{\rho}.$$

Dispersion relation. Insert a plane wave $u(\mathbf{x}, t) = e^{i(\mathbf{k}\cdot\mathbf{x} - \omega t)}$ and use $\nabla^{2n} \mapsto (-1)^n k^{2n}$. One obtains

$$-\omega^2 + i\gamma\omega - c_T^2 k^2 - \tilde{E}k^4 - \tilde{\eta}k^6 = 0,$$

or, equivalently (multiplying by -1),

$$\boxed{\omega^2 - i\gamma\omega + c_T^2 k^2 + \tilde{E}k^4 + \tilde{\eta}k^6 = 0}.$$

Thus $Im\omega = -\gamma/2 < 0$ for $\gamma > 0$: physical exponential decay. The positive signs $+c_T^2 k^2$, $+\tilde{E}k^4$, $+\tilde{\eta}k^6$ correspond to a positive quadratic energy and a sectorial generator (see Appendix T).

Coherence note (envelope visibility).

In interference estimates it is convenient to collect dephasing/inhomogeneous broadening into an effective transverse rate

$$\Gamma_{\perp} = \gamma_f + \gamma_{\phi} + \Gamma_{inh}, \gamma_f \simeq \frac{1}{2}\gamma,$$

so the propagation decoherence length is

$$\downarrow_{dec} \approx \frac{v_g}{\Gamma_{\perp}},$$

with v_g the group speed extracted from the dispersion. For slit–screen time of flight τ , the visibility decays as

$$\mathcal{V}(\tau) \approx \exp[-(\gamma_f + \gamma_{\phi} + \Gamma_{inh})\tau],$$

which reduces to $e^{-\gamma_f \tau}$ in the baseline STM (*no extra spinor dephasing and negligible inhomogeneity*).

Interpretation and limits.

Inertia gives ω^2 ; damping contributes $+i\gamma\omega \Rightarrow Im\omega = -\gamma/2$.

$c_T^2 k^2$ fixes the emergent light-cone speed; $\tilde{E}k^4$ and $\tilde{\eta}k^6$ stiffen the spectrum, with the sextic piece suppressing short-wavelength growth and bounding the UV.

Long-wavelength $k \rightarrow 0$: $\omega^2 \approx c_T^2 k^2$. High- k limit: $\omega^2 \approx \tilde{\eta}k^6$. Weak damping $\gamma \ll |\omega|$: conservative dispersion with a small imaginary correction.

When $\Delta E(\mathbf{x}, t; \mu)$ is non-negligible or periodic, the plane-wave analysis is replaced by the numerical framework of §2.5/Appendix K, or by a Bloch-type treatment for band structure.

Numerics hook (sign/unit fidelity). All solvers (Appendix Q) implement the linear spectral stencil

$$L(k) = -(c_T^2 k^2 + \tilde{E}k^4 + \tilde{\eta}k^6), \text{ and damping as } -\gamma \partial_t u,$$

which exactly matches the mass-normalised PDE above.

When modelling coherence/visibility rather than field amplitude, fold additional channels into $\Gamma_{\perp} = \gamma_f + \gamma_{\phi} + \Gamma_{inh}$ and apply the factor $\exp[-\Gamma_{\perp} \tau]$ (temporal) or $\exp[-\Gamma_{\perp} z/v_g]$ (spatial).

Propagator conventions and boundary prescriptions for L are collected in § 2.1.5. The envelope (Schrödinger) and path-integral structures derived from the same operator appear in § 3, with multi-scale details in Appendix D and open-system/response links in Appendices G and T.

2.1.5. Propagators and Response Functions (Linearised STM)

We fix propagators as inverses of the linearised STM operator L defined in § 2.1.1–2.1.4. We take $u(t, \mathbf{x}) \sim e^{-i\omega t + i\mathbf{k}\cdot\mathbf{x}}$, so $\partial_t \rightarrow -i\omega$ and $\nabla^{2n} \rightarrow (-1)^n k^{2n}$.

Retarded/response. The retarded Green function G_R solves

$$L G_R(\mathbf{x}, t; \mathbf{x}', t') = \delta^{(3)}(\mathbf{x} - \mathbf{x}') \delta(t - t'), G_R = 0 \text{ for } t < t'.$$

For homogeneous coefficients the frequency–wavenumber form reads

$$[-\rho\omega^2 + i\rho\gamma\omega - Tk^2 - (E_{STM} + \Delta E)k^4 - \eta k^6]G_R(k, \omega) = 1.$$

Optionally, in mass-normalised variables (divide by ρ):

$$[-\omega^2 + i\gamma\omega - c_T^2 k^2 - \tilde{E}k^4 - \tilde{\eta}k^6]G_R(k, \omega) = 1,$$

$$\tilde{E} = \frac{(E_{STM} + \Delta E)\downarrow_P^2}{\rho}, \tilde{\eta} = \frac{\eta\downarrow_P^4}{\rho}.$$

For small $\gamma > 0$, each simple pole lies at $\omega \approx \pm\Omega_k - i\gamma/2$ with $\Omega_k^2 = c_T^2 k^2 + \tilde{E}k^4 + \tilde{\eta}k^6$.

Time-ordered, Euclidean, and Keldysh. G_F uses the $+i0^+$ prescription; the Euclidean/Matsubara version follows $\omega \rightarrow i\omega_n$. For non-equilibrium we use the Schwinger–Keldysh triplet (G_R, G_A, G_K) , with G_K fixed by fluctuation–dissipation in equilibrium.

Inhomogeneity and boundaries. If parameters vary spatially, write $L = L_0 + V(\mathbf{x})$; then $G = G_0 + G_0 V G$ with $G_0 = L_0^{-1}$; contour choice fixes causality.

Spectral representation and stability.

$$G_R(\omega, \mathbf{k}) = \int_{-\infty}^{\infty} \frac{d\Omega}{2\pi} \frac{\rho(\Omega, \mathbf{k})}{\omega - \Omega + i0^+}, \rho(\Omega, \mathbf{k}) = -2\text{Im}G_R(\Omega + i0^+, \mathbf{k}).$$

Positivity and sum rules provide checks/identifications.

Envelope connection. The Schrödinger-type envelope kernel A is not the same object as G_R , but shares small- $|k|$ pole data (group velocity, curvature/diffraction, attenuation via γ), bridging membrane waves to envelope dynamics.

Notes on symbols. Here γ is Rayleigh damping (mass-normalised form used in § 3.1.1); g denotes the Yukawa/gauge coupling (not a scalar “gap”); \hbar is the usual reduced Planck constant.

2.2. Operator Quantisation

2.2.1. Canonical Commutation Relations

As in § 2.1.3, quantisation applies to the conservative Hamiltonian sector; the damping and forcing terms arising from the Rayleigh functional are excluded and play no role in the commutator structure.

Building on the Hamiltonian structure just introduced, we promote the displacement field $u(x, t)$ and its conjugate momentum $\pi(x, t)$ to operators $\hat{u}(x, t)$ and $\hat{\pi}(x, t)$ on a suitable Sobolev domain. The classical Poisson bracket

$$\{u(x), \pi(y)\}_{PB} = \delta^3(x - y)$$

is elevated via the Dirac correspondence

$$\{\cdot, \cdot\}_{PB} \longrightarrow \frac{1}{i\hbar}[\cdot, \cdot],$$

which immediately yields

$$[\hat{u}(x, t), \hat{\pi}(y, t)] = i\hbar \delta^3(x - y),$$

with all other commutators vanishing. Thus the non-commutativity of \hat{u} and $\hat{\pi}$ emerges naturally from the membrane’s intrinsic symplectic form, without requiring an extra quantisation postulate.

Spin–statistics. Equal-time CAR for the bimodal spinor Ψ are not assumed here; they are derived in Appendix O from locality, microlocal/Hadamard positivity and normal hyperbolicity, with stability under small, local BRST-compatible GKSL deformations.

2.2.2. Normal Mode Expansion

In nearly uniform regions, one may write

$$\hat{u}(x, t) = \int \frac{d^3k}{(2\pi)^3} e^{ik \cdot x} \hat{u}(k, t), \hat{\pi}(k, t) \text{ similarly.}$$

The associated Hamiltonian sums over the modes, each with a modified dispersion $\omega(k)$. When ΔE varies, a real-space diagonalisation or finite element approach is more suitable. Either way, the operator quantisation ensures a “quantum-like” spectrum of excitations that parallels bosonic fields in standard quantum theory.

2.3. Gauge Symmetries: Emergent Spinors and Path Integral

2.3.1. Bimodal Decomposition and Emergent Gauge Fields

A distinctive aspect of the STM model is constructing a bimodal decomposition of $\hat{u}(x, t)$. Formally, one splits u into two complementary oscillatory components, sometimes referred to as in-phase and out-of-phase fields:

$$u_1(x, t) = \frac{u + u_\perp}{\sqrt{2}}, u_2(x, t) = \frac{u - u_\perp}{\sqrt{2}},$$

and arranges (u_1, u_2) into a two-component spinor $\Psi(x, t)$. Imposing a local phase invariance $\Psi \rightarrow e^{i\alpha(x,t)} \Psi$ necessitates the introduction of gauge fields, e.g. A_μ for $U(1)$. Extending this principle can yield non-Abelian fields W_μ^a ($SU(2)$) and G_μ^a ($SU(3)$), reproducing the main gauge bosons familiar from the electroweak and strong interactions [15,16].

Mechanically, each gauge field arises as a compensating “connection” ensuring that local redefinitions of the spinor field do not alter physical observables. Consequently, photon-like or gluon-like excitations appear as coherent wave modes in the membrane. In standard quantum field theory, “virtual particles” mediate interactions; here, such processes correspond to deterministic wave–anti-wave cycles wherein net energy transfer over a full cycle is zero, aligning with the virtual-exchange picture. By including local phase invariance in the STM action, one automatically generates covariant derivatives $D_\mu = \partial_\mu - i g A_\mu$ (or the non-Abelian analogue), reinforcing how gauge fields naturally emerge from the underlying elasticity.

In the path-integral language, enforcing local spinor symmetries introduces these gauge connections and ghost fields (for gauge fixing) but does not rely on intrinsic randomness. Instead, it unites the deterministic elasticity framework with internal gauge invariance. This places photon-like excitations (for $U(1)$), W^\pm bosons (for $SU(2)$), and gluons (for $SU(3)$) as collective membrane oscillations that preserve local symmetry at each point in spacetime.

2.3.2. Ontology of Non-Abelian Gauge Fields

In STM the familiar non-Abelian gauge symmetries $SU(2)$ and $SU(3)$ arise in exactly the same way as $U(1)$, only now acting on higher-dimensional internal oscillator spaces. Concretely:

SU(2) as Local Doublet Rotations

At each spacetime point the STM spinor is promoted to a two-component doublet $\Psi = (\Psi_1, \Psi_2)$. The internal freedom to rotate

$$\Psi(x) \mapsto U(x) \Psi(x), \quad U(x) \in SU(2),$$

corresponds to choosing a new basis in the two-mode oscillator plane.

To compare $\Psi(x)$ and $\Psi(x + dx)$ without ambiguity we introduce the matrix-valued connection $A_\mu(x) \in \mathfrak{su}(2)$, so that the covariant derivative $D_\mu \Psi = \partial_\mu \Psi - i g A_\mu \Psi$ remains well-defined under local $SU(2)$ rotations.

Physically, each generator of SU(2) is realised as a distinct “twist” or shear of the STM membrane’s two-mode oscillation, and the Yang–Mills field strength $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]$ measures the membrane’s curvature in that internal rotation space.

SU(3) as Local Triplet Rotations

Similarly, for colour we carry a three-component oscillator $\Psi = (\Psi_1, \Psi_2, \Psi_3)$ transforming under local SU(3) rotations $\Psi \mapsto U(x)\Psi$, $U \in SU(3)$.

An eight-component connection $A_\mu^a T^a$ (with T^a the Gell-Mann generators) compensates infinitesimal changes in that three-mode orientation, yielding $D_\mu \Psi = \partial_\mu \Psi - i g_s A_\mu^a T^a \Psi$.

The associated field strength $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f^{abc} A_\mu^b A_\nu^c$ is nothing but the elastic-energy cost of non-commuting shears in the membrane’s colour-triplet oscillation bundle.

Unified Membrane Interpretation

In every case, gauge symmetry is simply the freedom to rotate the internal oscillator basis at each point in a way that costs elastic energy when misaligned.

All familiar Maxwell or Yang–Mills Lagrangians arise from writing down the membrane’s elastic energy as the square of these curvature two-forms.

Thus, U(1), SU(2) and SU(3) gauge fields share a single ontological origin: the tangent-space rotations of the STM membrane’s multimode oscillations.

No mandatory high-energy convergence

Because U(1), SU(2) and SU(3) all arise as different rotational polarisations of the same four-dimensional membrane, their common origin is already encoded in the Lagrangian. The usual grand-unification requirement $g_1 = g_2 = g_3$ at some ultra-high scale is therefore optional, not obligatory. Functional-RG trajectories in Appendix J show that for some stiffness ratios the three couplings *can* approach one another near the sextic fixed point, but nothing in the STM dynamics enforces that coincidence. Hence proton-decay bounds do not constrain STM, and the model accommodates either convergent or non-convergent running without additional fields.

Path-integral viewpoint.

For calculations, the emergent gauge sector can be handled in the usual BRST-fixed path integral,

$$\mathcal{Z} = \int \mathcal{D}u \mathcal{D}\psi \mathcal{D}\bar{\psi} \mathcal{D}A_\mu \mathcal{D}c \mathcal{D}\bar{c} \exp\{i[S_{STM}[u, \psi, A] + S_{gf}[A] + S_{gh}[A, c, \bar{c}]]\}.$$

$$\mathcal{Z}_{CTP} = \int \mathcal{D}\Phi_+ \mathcal{D}\Phi_- \exp\{i[S(\Phi_+) - S(\Phi_-)] + iS_{IF}[\Phi_+, \Phi_-]\},$$

2.3.3. Virtual Bosons as Deterministic Oscillations

In standard quantum field theory, “virtual particles” are ephemeral excitations in Feynman diagrams. Here, such processes are reinterpreted as energy-balanced wave–plus–anti-wave cycles. Over one period the cycle-averaged energy flux through any closed surface vanishes, consistent with a virtual (near-field) exchange:

$$\langle \oint_{\partial V} S \cdot dA \rangle_T = 0,$$

so the membrane’s local elastic/curvature energy is borrowed and returned within each cycle, rather than being dissipated or radiated.

By contrast, on-shell (far-field) emission — e.g. an annihilation on our face producing two photons — carries positive cycle-averaged flux,

$$\langle \oint_{\partial V} S \cdot dA \rangle_T > 0,$$

and the associated decrease of local strain/curvature equals the radiated energy (up to small damping losses). There is therefore no double counting: energy that leaves as real radiation is not simultaneously

assigned to a separate “curvature-relief” reservoir. Bound-state shifts (e.g. Lamb-like) are computed from the coarse-grained Hamiltonian density.

Equivalently, with \mathcal{E} the total STM+gauge energy density and Sits Noether/Poynting-like flux, the balance in a fixed control volume V is

$$\frac{d}{dt} \int_V \mathcal{E} d^3x = - \oint_{\partial V} \mathbf{S} \cdot d\mathbf{A} - \rho\gamma \int_V (\partial_t u)^2 d^3x + \int_V F_{ext} \partial_t u d^3x.$$

Near field: the surface term averages to zero over one period; far field: it does not.

2.4. UV Structure and Power Counting

We summarise the UV properties proven in Appendix Y. In the conservative limit ($\gamma_1 = 0$) the linearised STM Lagrangian yields $\pi = \rho\dot{\phi}$ and a positive quadratic Hamiltonian; higher derivatives are purely spatial, excluding Ostrogradsky modes. The dispersion $\omega^2 = (T/\rho)k^2 + (K_4/\rho)k^4 + (\eta/\rho)k^6$ implies $z = 3$ in $d = 3$. With $[\phi] = 0$ and $[\psi] = 3/2$, local scalar and Yukawa couplings carry positive mass dimension and the superficial degree of divergence obeys $\delta \leq 6 - 2E_\phi - 3E_\psi$, so only the elastic set $\{\rho, T, K_4, \eta\}$ and a finite number of relevant interaction normalisations renormalise. Gauge-invariant composites satisfy reflection positivity and admit a Källén–Lehmann representation; a gauge-covariant gradient flow provides a non-perturbative definition of renormalised local operators. Below a matching scale Λ_{match} the STM gauge-invariant Schwinger functions agree with Yang–Mills up to $\mathcal{O}(p^2/\Lambda_{match}^2)$; this underwrites the SMEFT bridge used in Apps. V–W.

2.5. Renormalisation and Higher-Order Corrections

2.5.1. One-Loop and Multi-Loop Analyses

The sixth-order operator $\eta \nabla^6 u$ ensures strong damping of high-momentum modes, so loop integrals converge more rapidly than in a naive second-order theory. Standard dimensional regularisation and a BPHZ subtraction scheme can be applied to compute self-energy corrections at one-loop or higher orders (see Appendix J). The resulting beta functions typically take the schematic form:

$$\beta(g_{eff}) = ag_{eff}^2 + bg_{eff}^3 + \dots,$$

where a, b are integrals influenced by $|k|^4$ and $|k|^6$ factors in the propagator. Multi-loop diagrams, including “setting sun” or mixed fermion–scalar topologies, refine these flows further. Crucially, running elastic couplings $E_{STM}(\mu)$ and $\Delta E(x, t; \mu)$ can exhibit non-trivial fixed points, opening the door to multiple stable vacua or discrete mass spectra.

2.5.2. Nonperturbative FRG and Solitons

Perturbation theory alone cannot capture phenomena like solitonic black hole cores or multiple vacuum states. Thus we employ Functional Renormalisation Group (FRG) methods in two complementary sectors: a calibrated $z = 3$ scalar analysis for the STM elastic potential (Appendix Y.10) and a deliberately simplified FRG-plus-soliton example model (Appendix L), both tracking an effective action Γ_k as fluctuations are integrated out down to scale k . These flows can reveal topologically stable solutions (e.g. kinks, domain walls) and non-trivial vacuum structure which are crucial for:

- **Fermion generation:** In the calibrated scalar sector (Appendix Y.10) an open set of STM-anchored UV couplings flows to an effective triple-well potential at a finite infrared scale; the three resulting elastic basins define distinct, well-separated mass scales, paralleling the three observed fermion generations and feeding the flavour templates of Appendix R.
- **Black hole regularisation:** Enhanced stiffness from ΔE and ∇^6 stops curvature blow-up, replacing singularities with finite-amplitude standing waves.

In the ultraviolet this $z = 3$ behaviour reflects a Lifshitz-type anisotropic scaling inherited from the STM ∇^6 regulator acting on the elastic/envelope field, and in Appendix Y.10 we analyse this sector in the standard way by freezing to a simple background (a local inertial/flat limit) in order to run

the FRG. This should be distinguished from Hořava–Lifshitz gravity: in STM the Lifshitz structure appears first in the elastic sector (second order in time, higher order in space), while the metric and Einstein– Λ gravity emerge in the infrared spin-2 channel (Appendix X.2 / Appendix M), rather than being postulated as a fundamentally anisotropic metric theory.

2.6. Classical Limit and Stationary-Phase Approximation

In a classical or macroscopic regime, one sets $\hbar \rightarrow 0$ or assumes heavy damping. The path integral

$$\int Du \exp\left\{\frac{i}{\hbar} S_{STM}[u]\right\}$$

is dominated by stationary-phase solutions of the PDE. Thus, the membrane behaves as a purely classical object with fourth- and sixth-order elasticity. Conversely, at sub-Planck scales—where the chaotic interplay of ΔE and ∇^6 acts—coarse-graining these rapid oscillations yields interference, Born-rule-like probability patterns, and gauge bosons as emergent wave modes (Appendix D).

Thus the familiar Schrödinger equation and its path-integral form are simply calculational devices—valid envelope approximations to our single, deterministic STM wave equation—rather than fundamental postulates of nature.

2.7. Non-Markovian Decoherence and Wavefunction Collapse

Although the STM PDE is fully deterministic, real-world observations exhibit effective wavefunction collapse. In STM this emerges from non-Markovian decoherence. We split the field into slow (“system”) and fast (“environment”) components, integrate out the fast sector on a closed-time-path (Schwinger–Keldysh) contour, and obtain a Feynman–Vernon influence functional $S_{IF}[u_s^+, u_s^-]$ [19]. The reduced state of the slow component then satisfies an integro-differential memory-kernel master equation. Off-diagonal elements in the pointer basis decay deterministically due to the finite correlation time of the eliminated modes, reproducing apparent measurement collapse without introducing fundamental randomness. In the Markov limit (short memory) the kernel reduces to a GKSL form with dephasing rate $\gamma_f \simeq \frac{1}{2}\gamma$ (as in §§2.1.1–2.1.4), and the construction is BRST-compatible (Appendix T.6).

This same non-Markovian structure supports deterministic entanglement analogues (Appendix E): at the coarse-grained level, CHSH-type correlations arise from a non-factorisable global standing wave and deterministic routing under spinor-based measurement operators, with decoherence and attractor dynamics supplied by the same finite- γ, γ_f open-system framework. Any apparent measurement dependence here reflects contextual boundary conditions and shared preparation (see §3.4.4), not a long-lived bath-mediated communication between distant wings. The rate and mechanism of decoherence are, in principle, testable in laboratory analogues and metamaterial platforms (Section 4.1; Appendix I).

2.8. Persistent Waves, Dark Energy, and the Cosmological Constant

In the long-wavelength, low-frequency limit $k \ll k_*$, the full SI STM equation

$$\rho \partial_t^2 u + T \nabla^2 u - (E_{STM} + \Delta E(\mathbf{x}, t)) \downarrow_P^2 \nabla^4 u + \eta \uparrow_P^4 \nabla^6 u = 0$$

reduces because the ∇^4 and ∇^6 operators are suppressed by factors k^2 and k^4 . The surviving SI balance is

$$\rho \partial_t^2 u + T \nabla^2 u \approx 0,$$

and, dividing by ρ ,

$$\partial_t^2 u - c_T^2 \nabla^2 u \approx 0, c_T^2 = \frac{T}{\rho}.$$

Both terms in the SI form carry units of force per unit volume Nm^{-3} (since $[\rho u_{tt}] = Nm^{-3}$ and $[T\nabla^2 u] = Nm^{-3}$). After division by ρ , every term has units of acceleration ms^{-2} .

The slow, mirror-induced modulation of the quartic stiffness,

$$E_{eff}(\mathbf{x}, t; \mu) = E_{STM}(\mu) + \Delta E(\mathbf{x}, t; \mu),$$

does not appear as a zero-derivative “mass term” $+\langle \Delta E \rangle u$ in the wave equation. Rather, it enters the energy balance through the $E_{eff}(\nabla^2 u)^2/2$ contribution to the energy density \mathcal{H} . When ΔE varies slowly, the local balance (§2.1.3) reads

$$\partial_t \mathcal{H} + \partial_i S_i = \frac{1}{2}(\partial_t \Delta E)(\nabla^2 u)^2 + \frac{1}{2} \nabla \Delta E \cdot \nabla [(\nabla^2 u)^2] - \rho \gamma u_t^2,$$

so phase-locked persistence of the standing/interference pattern comes from the parametric input $\frac{1}{2}(\partial_t \Delta E)(\nabla^2 u)^2$ balancing Rayleigh loss $\rho \gamma u_t^2$ in the near field (where the surface flux averages to zero over a cycle). The spatially uniform component $\langle \Delta E \rangle(t) = V^{-1} \int \Delta E d^3 x$ that survives coarse-graining acts as a tiny, slowly varying stiffness offset on large scales (H.4) and maps onto Λ_{STM} in the IR Einstein- Λ sector (see T.2); there is no double counting with Λ (§2.8/T.2 note).

Units note.

- In SI equations of motion: Nm^{-3} (force density).
- In energy density \mathcal{H} : $Jm^{-3} = Pa$ (pressure).
- Planck factors appear only with higher-order terms in SI: \uparrow_P^2 multiplies $\nabla^4 u$ and \uparrow_P^4 multiplies $\nabla^6 u$; the $\nabla^2 u$ term has no \uparrow_P factor.

A “Eureka” reinterpretation of the double-slit

Persistent waves via modulated stiffness

Historically, STM grew out of a simple but stubborn puzzle. Starting from a mirror-domain model in which particle–antiparticle pairs exchange energy with a spacetime membrane, the aim was to write down an elastic wave equation that reduces, at long wavelengths, to an Einstein-like field equation, yet still supports quantum-like interference. The eureka moment was to reinterpret the double-slit experiment not as an abstract statement about amplitudes, but as indirect evidence for persistent elastic standing waves on the membrane itself. In such a model those fringes would generically disperse or decay, so one is immediately forced to ask: how can long-lived, non-radiating interference patterns exist in an Einstein-analogue elastic spacetime that is continually exchanging energy with a mirror domain?

The critical insight was that mirror exchange should not appear as a crude body force, but as a small modulation of the quartic stiffness in the curvature-carrying part of the elastic energy. In STM this is written as

$$E_{eff}(x, t; \mu) = E_{STM}(\mu) + \Delta E(x, t; \mu),$$

with ΔE driven by mirror-sector dynamics. Even before the refined quantum machinery is introduced, a simple energy-balance analysis shows that a time-dependent ΔE naturally provides parametric energy exchange between the mirror sector and the membrane: persistent near-field waves can be sustained without explicit forcing, while purely ephemeral vacuum fluctuations average to zero and leave no lasting imprint. Time-averaging over the oscillations then reveals a small non-zero offset $\langle \Delta E \rangle$ in the effective stiffness. This offset behaves as a tiny vacuum-stiffness bias: it does not represent net stored energy for transient “vacuum ephemerals”, but the cumulative effect of long-lived, persistent waves.

In the full STM formulation this picture is made explicit. Treating the double-slit fringes as persistent elastic standing/interference waves on a Rayleigh-damped membrane immediately poses a sharper puzzle: with Rayleigh damping present, such waves cannot survive without a gentle, phase-locked trickle of energy. In STM this persistence arises because rapid mirror exchange (App. P)

induces a small, slowly varying modulation of the effective quartic stiffness $E_{\text{eff}}(x, t; \mu)$ as above. This modulated stiffness acts as a parametric feedback, phase-locking the oscillations and compensating the Rayleigh loss in the near field, so that the interference pattern survives without net radiation. When we coarse-grain, the spatially uniform component $\langle \Delta E \rangle$ appears as a tiny, slowly varying vacuum-stiffness offset in the Einstein-like sector, playing the rôle of an effective cosmological constant. In this way, the same stiffness modulation that keeps “quantum” interference patterns alive at laboratory scales also generates dark-energy-like behaviour on cosmological scales, providing a direct link between the quantum and cosmic regimes in the STM PDE.

At the level of energy balance (§2.1.3), a space–time dependent ΔE adds a source term. Locally,

$$\partial_t \mathcal{H} + \partial_i S_i = \frac{1}{2} (\partial_t \Delta E) (\nabla^2 u)^2 + \frac{1}{2} \nabla \Delta E \cdot \nabla [(\nabla^2 u)^2] - \rho \gamma (\partial_t u)^2,$$

so in a steady, near-field periodic state (no net surface flux, $\langle \oint \mathbf{S} \cdot d\mathbf{A} \rangle_T = 0$) the time-averaged parametric input balances damping,

$$\left\langle \frac{1}{2} (\partial_t \Delta E) (\nabla^2 u)^2 \right\rangle_T \approx \langle \rho \gamma (\partial_t u)^2 \rangle_T.$$

The uniform component $\langle \Delta E \rangle$ that survives coarse-graining then acts as a tiny, slowly varying stiffness offset on large scales, providing the same Λ_{STM} discussed in T.2 (with no double counting of Λ ; see §2.8/T.2).

B Emergent cosmological constant

The strain-to-curvature map of Appendix M.6 identifies the constant offset with a vacuum-energy term

$$\Lambda = \frac{8\pi G}{c^4} \langle \Delta E \rangle.$$

Because $E_{\text{STM}} \simeq c^4/8\pi G \approx 4.82 \times 10^{42} \text{Pa}$, an imperceptible fractional shift

$$\frac{\langle \Delta E \rangle}{E_{\text{STM}}} \sim 10^{-53}$$

reproduces the observed dark-energy density $\rho_\Lambda \approx 10^{-9} \text{Pa}$. Thus STM links quantum interference and cosmic acceleration without introducing extra fields or stochastic postulates.

A full spin-2 bootstrap showing that these assumptions yield Einstein’s equations with a cosmological constant appears in Appendix X §X.2.

For the detailed derivation of the vacuum-energy offset and its transition into the cosmological constant, see Appendix H (microphysical stiffness offsets) together with Appendices M.6–M.9 (emergent Λ in the 4D gravitational sector).

C Ultraviolet safety and solitonic cores

At large strain the sextic regulator $\eta \nabla^6 u$ dominates, raising the effective stiffness and preventing curvature blow-up. Appendices F.2–F.6 and L show that in collapse-like configurations this replaces general-relativistic singularities with finite-energy solitonic cores whose microcanonical entropy reproduces the *leading* Bekenstein–Hawking area law $S_{\text{BH}} = A/4G\hbar$ up to $\mathcal{O}(\lambda_c/R^*)$ corrections (Appendix F.4).

In the conservative $z = 3$ regime this UV softening is not only power-counting: Appendix Y.11.5 exhibits representative TT one-loop integrals built from the sextic propagator that are explicitly UV convergent, consistent with the finite local counterterm basis and UV-safety definition of Appendix Y.9.

D Numerical window

Even with $\langle \Delta E \rangle / E_{\text{STM}} \sim 10^{-53}$, quartic and sextic terms re-enter at LIGO-band strains ($\sim 10^{31} \text{Pa}$), far above laboratory scales yet well below the Planck modulus—ensuring consistency from tabletop interferometers to gravitational-wave astronomy.

In essence. A single eight-parameter elasticity law explains persistent quantum fringes, an effective cosmological constant, and singularity avoidance. The *eureka* insight—that double-slit coherence demands a mirror-induced, additive stiffness modulation—turns out to be the same ingredient that could drive the Universe’s late-time acceleration.

2.9. Action Principle in Curved Spacetime

2.9.1. Action Principle

We embed the STM framework on a four-dimensional Lorentzian manifold $(M, g_{\mu\nu})$ by writing the long-wavelength covariant completion as a single action

$$S = \int_M d^4x \sqrt{-g} \left[\frac{c^4}{16\pi G} R + \mathcal{L}_\phi + \mathcal{L}_\Psi + \mathcal{L}_{int} \right],$$

where R is the Ricci scalar of the metric $g_{\mu\nu}$.

Interpretation. The Einstein–Hilbert term is not introduced here as an independent “fundamental postulate”, but as the IR effective geometric wrapper implied by the STM strain–metric correspondence developed in Appendix M (linear mapping and spin-2 bootstrap). In the present manuscript we adopt the covariant action as a compact packaging of that correspondence, so that standard variational machinery yields the same long-wavelength equations. The logically prior claim is therefore: given the STM micro-PDE and the strain-built metric perturbation $h_{\mu\nu}[u]$, the unique local, diffeomorphism-invariant spin-2 dynamics at lowest derivative order is Einstein– Λ (Appendix X §X.2; Appendix M.8–M.9).

The STM Lagrangian splits into three parts:

- the scalar “membrane” sector \mathcal{L}_ϕ ,
- the two-component spinor sector \mathcal{L}_Ψ , and
- their elastic interaction \mathcal{L}_{int} .

All ordinary derivatives are replaced by Levi–Civita covariant derivatives, and each elastic constant enters as a diffeomorphism-invariant scalar. In particular, we define

$$\square := \nabla^\mu \nabla_\mu, \quad \{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}, \quad \bar{\Psi} = \Psi^\dagger \gamma^0.$$

2.9.2. Field Equations

Varying S with respect to $g^{\mu\nu}$ yields the Einstein equations with an STM stress–energy tensor:

$$\frac{c^4}{8\pi G} (R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}) = T_{\mu\nu}^{(\phi)} + T_{\mu\nu}^{(\Psi)} + T_{\mu\nu}^{(int)}.$$

In this paper the covariant variation is used as a consistency check and compact derivation of the IR equations; the substantive STM claim is that the Einstein– Λ form is recovered in the long-wavelength limit of the strain-metric map (Appendix M.8–M.9) and the spin-2 bootstrap assumptions stated in Appendix X §X.2.

Variation with respect to the scalar field ϕ gives a covariant sixth-order membrane equation:

$$\rho_0 \square \phi - E_{STM} \square^2 \phi - g \bar{\Psi} \Psi = 0.$$

Variation with respect to $\bar{\Psi}$ produces the curved-space Dirac equation with nonlinear coupling:

$$i \hbar \gamma^\mu \nabla_\mu \Psi - m \hbar \Psi - g \phi \Psi = 0.$$

2.9.3. Flat-Space and WKB Limits

By specialising $g_{\mu\nu} \rightarrow \eta_{\mu\nu}$, replacing $\nabla_\mu \rightarrow \partial_\mu$ and taking the semi-classical (WKB) limit, one recovers:

- the sixth-order scalar membrane PDE;

- the nonlinear Schrödinger-like envelope equation with STM coefficients;
- the elastic spinor–scalar coupling driving unseeded spinor emergence.

Thus the covariant formulation reduces exactly to the flat-space STM model under the appropriate limits.

2.10. Physical Calibration of STM Elastic Parameters

Even though the STM equation is written in dimensionless form, its coefficients must reproduce familiar physical constants when reinstated with units. The table below summarises each STM symbol, its calibrated SI value, and the physical anchor (derivation given in Appendix K.7):

STM symbol	Value (SI)	Anchor
ρ	$5.36 \times 10^{25} \text{ kg m}^{-3}$	κ/c^2
T	$4.82 \times 10^{42} \text{ Pa}$	ρc^2
Quartic stiffness $K_4 \equiv E_{STM\downarrow P}^{\uparrow 2}$ (coefficient of $\nabla^4 u$ in SI force-density form)	$K_4 = \frac{c^4}{8\pi G} \simeq 4.82 \times 10^{42} \text{ N}$	GR matching for the quartic operator.
ΔE	$6.8 \times 10^{-10} \text{ J m}^{-3}$	observed ρ_Λ
η	$9.3 \times 10^{11} \text{ Pa}$	UV cut off
g	0.3028	$\sqrt{4\pi\alpha}$
λ	0.13	Higgs quartic *
γ (coefficient of the term $\rho\gamma\partial_t u$)	$1.85 \times 10^{41} \text{ s}^{-1}$	Planck-time decoherence $\tau_c \approx L_P/c$

*Calibrated to the magnitude of the SM Higgs quartic to set the symmetry-breaking scale in the effective low-energy sector. This is not a claim of a fundamental Higgs field in STM: electroweak masses arise from deterministic face–mirror zitterbewegung and gauge-sector dressing (see §2.3 and App. C.3.1), with $V(u)$ acting as an emergent elastic proxy that fixes the infrared scale.

These eight calibrated coefficients — $\rho, T, E_{STM}, \Delta E, \eta, g, \lambda$ and γ — anchor the STM model quantitatively to c, G, α, Λ and the Planck scales, yielding a fully testable system of dimensionless parameters for use in Sections 3 and 4. The same sign choices ensure positivity/sectoriality in the conservative UV analysis (§2.4; App. Y). In explicit SI force-density form the higher-order terms are $K_4 \nabla^4 u$ and $\eta \nabla^6 u$, with $K_4 \equiv E_{STM\downarrow P}^{\uparrow 2}$. When reporting SI coefficients, quote K_4 in newtons rather than E_{STM} in pascals, and provide A_4 or K_4 as appropriate for spatial-propagation fits.

2.11. Tensor-Mode EFT Mapping (STM \rightarrow GR with Soft UV Corrections)

In this subsection we isolate the transverse–traceless tensor sector of STM and present a minimal effective-field-theory (EFT) mapping to a GR-like graviton with soft UV corrections. The goal is to provide a clean dictionary between the STM elastic coefficients and the usual graviton EFT parameters, not to claim a full derivation of general relativity from STM.

Sign choices and parameter map.

We work with the same sign conventions as in §2.1.1 and §2.4 and assume

$$\rho > 0, \quad T > 0, \quad E_{STM} \geq 0, \quad \eta > 0.$$

In the tensor sector we identify

$$c_g^2 = \frac{T}{\rho}, \quad \frac{\alpha_2}{\Lambda_g^2} = \frac{E_{STM}}{\rho}, \quad \frac{\alpha_3}{\Lambda_g^4} = \frac{\eta}{\rho},$$

with $c_g \rightarrow 1$ under IR Lorentz restoration (solver/SI conversions as in App. K.6–K.7).

In the explicit-SI PDE of §2.1.1 the quartic piece is $E_{STM\downarrow P}^{\uparrow 2} (\nabla^2 u)^2$, so the mass-normalised SI coefficient is

$$\tilde{E} = \frac{E_{STM}}{\rho} \downarrow p,$$

and one may equivalently write $\alpha_2/\Lambda_g^2 = \tilde{E}$. In what follows α_2, α_3 are dimensionless EFT parameters expected to be $\mathcal{O}(1)$, while Λ_g encodes the effective UV scale of the tensor sector.

Minimal graviton EFT (closed/unitary sector).

Restricting to transverse–traceless (TT) modes on a GR background, the linearised STM tensor sector is second order in time and higher order in space. A convenient EFT parametrisation of the TT dispersion relation is

$$\omega^2 = c_g^2 k^2 + \frac{\alpha_2}{\Lambda_g^2} k^4 + \frac{\alpha_3}{\Lambda_g^4} k^6 \quad (c_g \rightarrow 1 \text{ in the IR}),$$

which carries no extra polarisations and no higher-time derivatives (so no Ostrogradsky ghosts).

In this language:

- the $c_g^2 k^2$ term coincides with GR's massless graviton in the IR;
- the k^4 and k^6 pieces encode STM's quartic and sextic stiffness as soft UV corrections;
- observable consequences appear as small modifications to gravitational-wave dispersion/phasing, static weak-field tails and black-hole ringdown spectra.

Observational constraint: gravitational-wave dispersion.

If the emergent spin-2 sector inherits residual higher-spatial-derivative corrections at long wavelengths, gravitational-wave propagation acquires weak frequency dependence. Writing the group velocity as

$$v_g(q) = c[1 - \beta(q/\Lambda_{LR})^n + \dots],$$

a wavepacket of central wavenumber q and bandwidth Δq propagated over distance D develops an arrival-time broadening

$$\Delta t \sim D \frac{\partial}{\partial q} \left(\frac{1}{v_g} \right) \Delta q.$$

The observed absence of appreciable dispersion across the detected gravitational-wave bandwidth therefore bounds β/Λ_{LR}^n and, equivalently, places a lower bound on Λ_{LR} (for $n > 0$, Λ_{LR} must lie well above the GW band). This supplies a direct, falsifiable constraint on the Lorentz-restoration scale implied by the STM UV regulator.

Here q denotes the physical wavenumber of the propagating GW mode (not the FRG scale k used in Y.10–Y.12).

Appendix M gives the Einstein-like mapping in more detail and §M.11 quotes the corresponding phenomenological scalings.

Static weak-field potential.

In the static, weak-field limit the tensor-mode EFT reduces to a modified Poisson equation for the potential Φ :

$$\left(-T\nabla^2 + E_{STM}\nabla^4 + \eta\nabla^6 \right) \Phi(\mathbf{r}) = 4\pi GM\delta^{(3)}(\mathbf{r}).$$

Solving this with the sign choices above yields the standard Newtonian term

$$\Phi_N(r) = -\frac{GM}{r}$$

plus short-range Yukawa and/or oscillatory corrections, whose ranges and relative weights are fixed by the roots of the characteristic polynomial

$$-Tq^2 + E_{STM}q^4 + \eta q^6 = 0.$$

These corrections are controlled by the same (E_{STM}, η) that enter the graviton dispersion; they therefore provide an independent way to constrain α_2/Λ_g^2 and α_3/Λ_g^4 via laboratory and Solar-System tests of gravity.

Scope of the mapping. The dictionary above is restricted to the closed, linear transverse–traceless (TT) sector, which captures the radiative tensor content in the weak-field regime (Appendix M.8), and assumes that scalar and vector perturbations are either non-propagating or heavy/screened over the regimes relevant for gravitational-wave dispersion/phasing, near-field static tests, and ringdown phenomenology. Non-linear back-reaction and strong-field effects are treated separately in Appendix M; here we keep only the minimal tensor-mode EFT needed to connect STM’s calibrated elastic coefficients to observable TT two-point structures. The conservative UV logic used to interpret this mapping is the same as in Appendix Y: time derivatives remain strictly second order while UV control is provided by higher-order spatial stiffness (Appendix Y.2), leading to $d = 3, z = 3$ scaling (Appendix Y.3) and a finite counterterm/renormalisation set in the conservative truncation (Appendix Y.4 and Appendix Y.9).

UV scaling (Lifshitz-type, as used in Appendix Y). In the regime where sextic stiffness dominates, the TT two-point structure behaves as $G_{TT}(\omega, k) \sim (\omega^2 + k^6)^{-1}$, matching the same $z = 3$ Lifshitz-type scaling [22] defined in Appendix Y.3 (sextic dominance $\Rightarrow z = 3$ in $d = 3$). The present section uses this as a kinematic EFT dictionary for tensor propagation; Appendix Y provides the underlying conservative scaling and power-counting framework.

We use ‘Lifshitz-type’ purely to denote the anisotropic $z = 3$ UV scaling implied by STM’s sextic spatial stiffness in the conservative sector; unlike Hořava–Lifshitz gravity [23], STM does not postulate a fundamentally foliated metric theory as the UV starting point. Instead, the metric description is an IR/emergent covariant EFT fixed at two-derivative order by the spin-2 bootstrap, while the UV softening resides in the underlying elastic degrees of freedom.

Why IR Einstein– Λ does not imply GR’s UV problem. The spin-2 bootstrap (Appendix X.2) fixes the unique two-derivative IR completion to Einstein– Λ , but it does not assert that the Einstein–Hilbert truncation is the correct description at arbitrarily high momenta. In STM, UV behaviour is governed by the higher-spatial-derivative stiffness (the same physics encoded here by the k^4 and k^6 terms), and the conservative $z = 3$ power counting implies that (when divergences occur) they renormalise only a finite set of structures already present in the calibrated elastic sector, consistent with the divergence bound of Appendix Y.4 and the finite renormalisation set of Appendix Y.9. This is consistent with the EFT statement in Appendix X.2 that deviations from GR enter through higher-derivative operators suppressed by Λ_{STM}^{-2} (cf. Appendix V/W).

Physical interpretation and alternative pipeline (STM versus geometry-first GR)

The standard derivation of the Einstein field equations is rightly admired for its internal closure: starting from general covariance and local stress–energy conservation, the contracted Bianchi identity $\nabla^\mu G_{\mu\nu} = 0$ singles out the Einstein tensor $G_{\mu\nu}$ as the natural divergence-free curvature object, enabling a universal coupling to a conserved stress–energy tensor $T_{\mu\nu}$. In the usual geometry-first presentation, curvature is introduced as the primitive notion and its operational meaning—free fall, tidal response and focusing—is recovered a posteriori.

STM proposes a complementary ordering of explanation: mechanics first, geometry second. One begins with a physical medium (the STM elastic field and its stress–energy functionals), then performs a coarse-graining in which rapidly varying sub-Planck structure is integrated out, leaving a low-frequency effective description of what observers can operationally measure: the propagation of long-wavelength excitations, clock–rod comparisons, and the relative acceleration of nearby inertial trajectories. In this pipeline, a metric and its curvature are not fundamental objects but macroscopic bookkeeping variables introduced precisely to summarise those operational effects in a coordinate-free form.

The primitive gravitational observable in this mechanics-first pipeline is congruence evolution. For a timelike geodesic congruence with tangent u^μ , the Raychaudhuri equation shows that the Ricci

contraction $R_{\mu\nu}u^\mu u^\nu$ controls the local volume focusing or defocusing of freely falling bundles. STM adopts this same operational content, but with a different ontology: Ricci curvature is not taken as fundamental structure, but as the coarse-grained focusing tensor that encodes the medium's effective compressive or dilational response once microscopic elastic structure is averaged over. Geometry, in this sense, records focusing behaviour rather than causing it.

Once focusing is taken as primary, the role of the Einstein tensor becomes transparent. Any macroscopic description intended to represent a closed balance law must be compatible with local conservation. In geometric language this requirement is expressed by $\nabla^\mu G_{\mu\nu} = 0$, guaranteed by the contracted Bianchi identities. From the STM viewpoint this appears not as an unexpected mathematical miracle but as the covariant packaging required for a consistent effective stress–balance law. The celebrated geometric structure is therefore retained, but its interpretation is inverted: geometry is the universal, coordinate-free representation of a conservation-respecting mechanical response.

Matter sourcing is likewise re-read in mechanics-first terms. In the geometry-first pipeline one specifies matter fields on spacetime and computes $T_{\mu\nu}$. In STM, the stress–energy entering the operative field equations is understood as the coarse-grained energy–momentum carried by excitations of the same underlying medium. In the projected low-energy description adopted here, this is represented by the spinor sector $T_{\mu\nu}^{(\Psi)}$, while uniform offsets in the medium's energy or stiffness budget are absorbed into an effective vacuum term. The resulting operative field equation takes the familiar Einstein– Λ form,

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}^{(\Psi)},$$

but with a distinct ontological reading: the right-hand side encodes the medium's coarse-grained excitation content, while the left-hand side encodes the medium's coarse-grained focusing response, written in the unique conservation-compatible covariant form.

Finally, the cosmological term acquires a straightforward interpretation in this pipeline. A spatially uniform coarse-grained offset in the medium's energy or stiffness budget biases focusing even in the absence of localised excitations. In a generally covariant geometric description, the only such universal contribution is a $\Lambda g_{\mu\nu}$ term. The cosmological constant is thus interpreted not as an arbitrary addition but as the macroscopic signature of a background offset of the underlying medium.

This interpretation is intentionally ontological rather than algebraic. It is fully compatible with the standard geometric reading of General Relativity and does not alter its field equations; rather, it provides a mechanics-first explanatory route whose distinctiveness becomes testable only when ultraviolet corrections, dispersion effects, or departures from the Einstein– Λ limit are considered.

2.12. Summary of Methods

We start from a single high-order elastic wave equation for the membrane displacement u , incorporating scale-dependent stiffness, fourth- and sixth-order spatial derivatives, linear damping, cubic non-linearity, Yukawa-like coupling to emergent spinors and external forcing.

Canonical quantisation promotes u and its conjugate momentum to operators in a suitable Sobolev space, with self-adjoint Hamiltonian terms up to ∇^6 . All eight coefficients are *calibrated once* to (c, G, α, Λ) and the Planck scales (table in §2.10; App. K.7); thereafter, no additional elastic or damping parameters are tuned in the predictions reported here. In the IR, the spin-2 bootstrap (App. X.2) shows that STM's two-derivative gravitational sector closes uniquely to Einstein– Λ with matter stress–energy supplied by the emergent spinors.

A bimodal decomposition of u yields a two-component spinor field; imposing local phase invariance generates U(1), SU(2) and SU(3) gauge fields.

A multiple-scale (WKB) expansion separates fast sub-Planck oscillations from a slow envelope, giving an effective Schrödinger-like equation whose interference, Born-rule density and decoherence follow deterministically once $\gamma > 0$ is included (see Section 3.4).

Functional and perturbative renormalisation analyses exploit the ∇^6 term to tame UV divergences, reveal non-trivial fixed points (fermion generations) and support solitonic cores (singularity avoidance). Quantitatively, this conservative UV control is captured by the $d = 3, z = 3$ scaling and canonical dimensions in Appendix Y.3 and by the resulting divergence bound and finite counterterm set in Appendix Y.4 and Appendix Y.9, while Appendix Y.10 gives the corresponding scalar FRG implementation; the spin-2 bootstrap fixes only the two-derivative IR closure to Einstein- Λ (Appendix X.2).

The model is immediately testable through a Mylar-membrane interferometer whose geometry fixes the quartic stiffness/propagation coefficient A_4 by design, leaving the mass-normalised damping γ as the sole exported laboratory parameter (see App. I.2). This single number calibrates flavour dephasing ($\gamma_f = \gamma/2$) and informs gravitational-wave amplitude analyses ($\Gamma \simeq \gamma/(2c)$, with c the speed of light); consequently, a Mylar measurement or bound on γ propagates directly to tighter cross-sector constraints (see App. I.6)

3. Results

This section presents the principal findings of the Space-Time Membrane (STM) model.

Throughout this section we work in the near-paraxial, small-amplitude regime of the linearised operator L , with quasi-monochromatic carriers (ω_0, \mathbf{k}_0) , slowly varying coefficients $\{\rho, T, E_{STM}, \eta, \gamma, \Delta E\}$, and a clear carrier-envelope scale separation; all envelope and path-integral statements are at leading non-trivial order in these parameters. We use the operator, Fourier and damping conventions fixed in § 2.1.5 throughout Section 3.

Gravitational-wave and static predictions use the tensor-mode parameterisation introduced in §2.11, with IR Einstein- Λ behaviour supplied by Appendix X/M.

We begin by examining perturbative results, illustrating how quantum-like dynamics, gauge symmetries, and deterministic decoherence arise from a high-order elasticity framework. We then turn to nonperturbative effects, whose full derivation—via the Functional Renormalisation Group (FRG)—appears in Appendix L.

3.1. Perturbative Results

3.1.0. Emergent Schrödinger-Like Dynamics and the Born Rule

We start from the mass-normalised STM equation of motion (see §2.1.1):

$$\partial_{tt}u - \gamma \partial_t u + c_T^2 \nabla^2 u - \tilde{E} \nabla^4 u + \tilde{\eta} \nabla^6 u - \tilde{\lambda} u^3 - \tilde{g} u \bar{\psi} \psi = 0$$

where $c_T^2 = T/\rho$, $\tilde{E} = (E_{STM} + \Delta E) \downarrow_p^2 / \rho$, $\tilde{\eta} = \eta \downarrow_p^4 / \rho$, and $\gamma > 0$ is the calibrated damping (Appendix K.7). (SI force-density form has $-\rho \gamma \partial_t u$; after dividing by ρ this becomes $-\gamma \partial_t u$.)

With the conventions of § 2.1.5, the linearised dispersion is

$$\omega^2 - i\gamma\omega + c_T^2 k^2 + \tilde{E} k^4 + \tilde{\eta} k^6 = 0 \quad \Rightarrow \quad \text{Im } \omega = -\gamma/2 < 0,$$

so damping is strictly stabilising (Appendix T).

Envelope/WKB reduction. By coarse-graining the rapid, sub-Planck oscillations in $u(x, t)$, one obtains a slowly varying “envelope” $\Psi(x, t)$. Specifically, one applies a smoothing kernel (often Gaussian) and adopts a WKB-type ansatz,

$$\Psi(\mathbf{x}, t) = A(\mathbf{x}, t) e^{\frac{i}{\hbar} S(\mathbf{x}, t)},$$

and expanding (3.1) order-by-order separates into a Hamilton-Jacobi equation for S and a continuity equation for A . Using the operator, Fourier and damping conventions of § 2.1.5, the near-paraxial coarse-graining yields the envelope law stated below. At leading order the slow envelope obeys an effective Schrödinger equation,

$$i\hbar\partial_t\Psi = -\frac{\hbar^2}{2m_{eff}}\nabla^2\Psi + V_{eff}(\mathbf{x})\Psi - i\hbar\gamma\Psi, \quad (3.3)$$

where m_{eff} and V_{eff} are fixed combinations of c_T^2 , \tilde{E} , $\tilde{\eta}$ and the slow background; the ∇^2 term controls the UV through its contribution to those coefficients (details in Appendix D). The small positive γ is the calibrated, mass-normalised damping ($\gamma_{nd} = 0.010$), entering here as a linear attenuation of the envelope.

Born rule and determinism. With $\gamma > 0$, the envelope dynamics contract onto measurement-defined attractors, and $|\Psi|^2$ realises the Born weights via deterministic sub-Planck chaos (Section 3.4; Appendix P). Thus STM reproduces interference and outcome statistics without intrinsic randomness [20].

Coarse-graining versus indeterminism. It is important to distinguish (i) the deterministic microdynamics of the full field u from (ii) the effective envelope-level statistics of ψ . The Born weights arise because coarse-graining and thresholded readout define a non-invertible projection from microstate to macroscopic outcome: many distinct sub-Planck phase histories of u map to the same observed pointer value. This multiplicity is a feature of the observational V_{eff} projection, not a failure of determinism of the underlying PDE.

While this deterministic approach now reproduces the Born rule and (for non-factorisable joint preparations, Appendix E) approaches the Tsirelson bound, it still departs conceptually from the mainstream view that quantum indeterminism is fundamental. Rigorous loop-level checks (e.g. chiral anomalies) and targeted experiments — such as ultra-long interferometry or fast-switch Bell tests that probe the finite memory time predicted by the STM kernel — are needed to confirm whether the model matches standard quantum mechanics at all scales.

3.1.1. Uncertainty Relations from STM

Fourier/kinematic route. Any STM wave-packet obeys $\Delta x \Delta k \geq \frac{1}{2}$. With $p = \hbar k$ for the envelope's canonical pair (Appendix D), this gives

$$\Delta x \Delta p \geq \frac{\hbar}{2}.$$

Canonical/commutator route. The envelope action carries the symplectic term $i\hbar A^* \partial_t A$, hence $[x, p] = i\hbar$ and the same bound $\Delta x \Delta p \geq \hbar/2$.

Time-energy. With small mass-normalised damping $\gamma > 0$, the retarded pole has width $Im\omega \simeq -\gamma/2$, so $\Delta\omega\Delta t \gtrsim \frac{1}{2}$ and therefore $\Delta E\Delta t \gtrsim \hbar/2$. (See the dispersion and damping sign already established in § 2.1/§ 2.1.5 and used in § 3.1.1.)

Pipeline Rosetta Flow

STM PDE (§ 2.1) → invert L (choose contour) → **propagators** G_R, G_A, G_F /Keldysh (G_R, G_A, G_K);
 STM PDE → multi-scale/WKB → envelope (**Schrödinger**);
 envelope → integrate out fast modes → effective action S_{eff} → **path-integral** kernel K .

3.1.2. Emergent Gauge Symmetries

A hallmark of STM is that gauge structure emerges from the bimodal decomposition of the displacement field $u(x, t)$. Writing the two slow envelopes as (u_1, u_2) and assembling them into a two-component spinor $\Psi(x, t)$, we impose local phase covariance on Ψ , which forces gauge connections via minimal coupling.

Abelian case (U(1)). Under a local re-phasing

$$\Psi(x) \rightarrow e^{i\theta(x)}\Psi(x),$$

covariance of the envelope dynamics requires

$$\partial_\mu \rightarrow D_\mu = \partial_\mu - iq A_\mu(x),$$

with a Maxwell term $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ in the emergent Lagrangian and conserved current $J^\mu = \bar{\Psi}\gamma^\mu\Psi$.

Non-Abelian extension. Demanding local symmetry in the internal mode space and colour space yields the Yang–Mills structure

$$D_\mu = \partial_\mu - i g_1 Y B_\mu - i g_2 \tau^a W_\mu^a - i g_3 T^a G_\mu^a,$$

with the standard field strengths. In this reading, photon-, W^\pm/Z^0 - and gluon-like excitations are coherent membrane modes (collective waves) rather than stochastic quanta [16]. Processes usually described as “virtual boson exchange” are reinterpreted as wave–plus–anti-wave cycles that exchange phase/momentum while averaging zero net energy over a cycle (cf. § 2.1.2, Appendix P).

Analogies (intuition building).

- *Electromagnetism as phase connection.* The U(1) potential plays the role of a geometric “connection” that keeps the local phase of the two-mode envelope aligned across the membrane—exactly as a classical bundle connection aligns phases along a fibre [16].
- *Strong force as a lattice of coupled oscillators.* Visualising the membrane as a lattice of linked oscillators, each site carrying an effective “colour,” the elastic coupling stiffens with separation, producing a linearly rising energy cost and hence confinement; gluon-like modes are the coherent waves on those links [15].
- *Virtuals as counter-oscillations.* Internal lines in Feynman graphs map to counter-propagating wave pairs whose instantaneous energy budget is balanced over a period, preserving exact energy conservation while reproducing the same effective interactions.

Electroweak breaking (deterministic mechanism). When a uniform background displacement is present and face–mirror couplings are switched on, rapid zitterbewegung between Ψ and its mirror partner $\tilde{\Psi}_\perp$ (§ 3.7) generates effective mass terms for the weak gauge fields, mimicking electroweak symmetry breaking without positing a fundamental scalar at the microscopic level. The constructive details and parameter map are given in Appendix C.3.1.

Formal consistency: anomalies and damping.

- **Anomaly cancellation.** Appendix U proves that mirror doubling renders the chiral spectrum vector-like, so all perturbative gauge, mixed and gravitational anomalies cancel on any globally-hyperbolic background; BRST nilpotency is preserved.
- **BRST-compatible dissipation.** The small Lindblad terms used for damping/dephasing commute with the BRST charge (or are BRST-exact), so the physical cohomology is preserved under open evolution (Appendix T, Thm. T.6), consistent with the dissipator choices in § 3.4 and Appendix P.

Empirical touchstone. The emergent gauge Lagrangian reproduces the tree-level $e^+e^- \rightarrow \mu^+\mu^-$ differential cross-section, including γ – Z interference and one-loop leptonic running $\alpha(s)$ (Appendix S; Scattering_amplitude.py). This anchors the correspondence beyond formal symmetry.

Data-facing bridge. For comparisons beyond tree-level $e^+e^- \rightarrow \mu^+\mu^-$, Appendix V provides a compact STM→SMEFT dictionary (including $O_{2B}, O_{2W}, \Psi^2F^2, \Psi^4$) and Appendix W extends this to the full $SU(3) \times SU(2) \times U(1)$ set with one-loop running to m_Z and the W, Y oblique mapping. These let global-fit bounds be translated directly into STM parameter inequalities.

Scope and next steps. With $U(1) \times SU(2) \times SU(3)$ realised and anomaly freedom secured, full equivalence with the Standard Model requires extending tests to non-Abelian multi-leg processes and loop corrections; those (together with the effective-scalar channel of Appendix C) are the next targets.

Result (Appendix O). In the conservative limit, scalar/membrane observables commute at space-like separation (CCR), while the bimodal spinor obeys CAR and anticommutes at spacelike separation; the graded locality persists for $0 \leq \gamma < \gamma_*$ under local BRST-compatible GKSL deformations.

Computational note. A practical realisation of the elastic–spinor coupling, constraint preservation, and the GKSL step used in our solvers is given in §3.1.4 (see also Appendix P; Appendix T, Thm. T.6).

3.1.3. Deterministic Decoherence and Bell-Inequality Violations

Splitting the displacement field into slow “system” and fast “environment” components, $u = u_S + u_E$, and integrating out u_E with a Feynman–Vernon influence functional (Appendix G) yields a non-Markovian master equation for the reduced state $\rho_S(t)$:

$$\frac{d\rho_S}{dt} = -\frac{i}{\hbar} [H_S, \rho_S] - \int_0^t d\tau K(t - \tau) \mathcal{D}[\rho_S(\tau)],$$

where the memory kernel K encodes finite bath correlation times and $\mathcal{D}[\cdot]$ is a quadratic superoperator fixed by the STM couplings and the coarse-grained bath statistics. In the short-memory limit (Appendix P), the kernel reduces to a GKSL/Lindblad form that is BRST-compatible on curved backgrounds (Appendix T, Thm. T.6), ensuring positivity and preservation of the physical subspace.

Appendix X extends this analysis to genuinely non-Markovian kernels, showing that under the STM sign and regularity assumptions the Schwinger–Keldysh reduced dynamics remain retarded and completely positive/trace-preserving

Measurement and collapse (deterministic). The small but fixed Rayleigh damping $-\gamma \partial_t u$ in the scalar sector (mass-normalised form) induces envelope-level dephasing of the emergent spinor channels at rate $\gamma_f \simeq \frac{1}{2}\gamma$ (Appendix K.6–K.7, § 3.4). With spinor-based measurement operators (e.g. $\hat{M}(\theta) = \cos\theta \sigma_x + \sin\theta \sigma_z$), the ensuing dynamics deterministically drive the state to one of the measurement basins (Appendix E), without stochastic postulates.

Different analyser settings correspond to different boundary conditions (i.e. different coarse-grained measurement operators acting on the same underlying field), so comparing runs across settings is comparing different boundary-value problems, not diagnosing indeterminism of the micro-PDE.

Bell correlations from internal phase. Each spin packet carries a fixed internal phase between its two elastic modes; a Stern–Gerlach analyser at angle θ projects that phase onto its eigen-basis. The coincidence curve

$$P_{\uparrow}(\theta) = \cos^2 \frac{\theta}{2}, \quad P_{\downarrow}(\theta) = \sin^2 \frac{\theta}{2}$$

follows as squared overlaps (derivation in Appendix E.3). For two analysers at angles a, b , the correlation is $E(a, b) = \cos(a - b)$, and the CHSH parameter reaches $2\sqrt{2}$, [20,21] violating the classical Bell bound while arising from a deterministic PDE with finite-memory dissipation.

Caveat and targets. Laboratory environments are often near-Markovian; quantitative comparison of STM-predicted decoherence times and memory kernels with experiment (including environment-induced superselection) remains an open programme. The curved-space, BRST-compatible dissipator used here guarantees complete positivity and gauge compatibility (Appendices P, T), anchoring those comparisons on a self-consistent foundation.

3.1.4. Fermion Generations, Flavour Dynamics, and Confinement

Motivation for three basins. Combining the perturbative running of the elastic couplings (Appendix J) with a $z = 3$ scalar FRG proxy analysis (Appendix Y.10) suggests that STM-compatible ultraviolet elastic couplings can, for suitable effective parameters, support a triple-well potential along a dominant scalar direction at a finite coarse-graining scale. In line with the triple-well region \mathcal{R}_3 identified there, we adopt as a working hypothesis that the full STM scalar–fermion–gauge system can generate three robust elastic basins in configuration space, with well-separated minima.

Basis of the CKM/PMNS results and what Appendix Y adds. Importantly, the flavour-mixing scans reported in Appendix R do not use FRG outputs from Appendix Y. Appendix R is a self-contained phenomenological construction based solely on the three-mode elastic ansatz and derivations developed in Appendix R itself: a trivalent envelope localised near three wells x_1, x_2, x_3 , a Hermitian 3×3 elastic operator Ω built from inter-well couplings and phases, and a scan/fit procedure (implemented in `STM_flavour_scan_chi2_r1.py` – see Supplementary Information) in which basin/well positions and overlap-controlled couplings are treated as inputs scanned within STM-calibrated ranges.

Appendix Y is included for theoretical support and to define a future predictive route from STM to basin data: Appendix Y.10 provides a conservative single-direction proxy showing how triple-well structure arises naturally in a stabilised sextic potential over a well-defined region \mathcal{R}_3 , and clarifies why additional matter/gauge contributions can plausibly realise the mildly negative effective quartics needed for multi-basin structure in STM-consistent calibrations. Appendix Y.12 then specifies a three-component basin-field FRG truncation together with an explicit flavour-interface export contract (basin amplitudes/curvatures/overlaps at an extraction scale k^*) which, in a later matter-complete implementation, could replace phenomenological basin inputs by FRG-derived basin observables—turning the Appendix R template from a scan-based fit into a predictive pipeline. A full multi-field FRG derivation of a unique STM calibration point within \mathcal{R}_3 is left to future work.

CP phases from deterministic zitterbewegung. The bimodal spinor Ψ on the physical face and its mirror partner Ψ_\perp on the opposite face undergo rapid, antisymmetric zitterbewegung. Coarse-graining these face–mirror oscillations imprints complex phases into effective Yukawa terms, generating CP violation. Weak gauge fields and electroweak mixing emerge from the same elastic coupling structure; the constructive map is given in Appendix C.3.1.

Quantitative flavour fits (no flavour tuning). With the calibrated non-dimensional set ($E_{nd}, \eta_{nd}, \gamma_{nd} = 0.010, \gamma_{f,nd} = 0.005$) and an STM prior ensemble that is flat in the CKM elastic parameters and broad STM-scale Gaussian in a minimal seesaw block, the scans (Appendix R), evaluated in the unitary ($\gamma = \gamma_f = 0$) limit for mixing extraction, reproduce:

- **CKM:** all nine moduli at PDG-2024 precision; $\chi_{CKM}^2 = 0.21$ (primary band) and 0.33(sensitivity band). A short CP-phase polish then aligns the Jarlskog to $J_{PDG} \approx 3.08 \times 10^{-5}$ under a stiff $|U|$ penalty, without materially altering $|U|$; unitarity is preserved to $\|U^\dagger U - \mathbb{1}\|_F \sim 10^{-15}$. Acceptance fractions and residuals are tabulated in Appendix R.
- **PMNS:** a parameter-space fit (normal ordering) to $\{\sin^2\theta_{12}, \sin^2\theta_{23}, \sin^2\theta_{13}, \delta_{CP}, \Delta m_{21}^2, \Delta m_{32}^2\}$, yielding $\chi_{PMNS}^2 = 3.30$ (primary band) and 6.40(sensitivity band); the displayed $|U|$ is reconstructed from the parameter best-fit.

Treating quark and lepton sectors as independent scans, the joint acceptance is their product (reported in Appendix R), underscoring how non-generic it is to match both sectors without extra tuning.

Generation flow and stability. The discrete vacuum structure also explains observed decay patterns. Quarks, subject to strong colour interactions, sit on a confining elastic background; higher-generation quarks (associated with the higher fixed points) carry excess elastic energy and deterministically relax downward to lower-generation vacua via gauge-mediated channels. Leptons, by contrast, are not colour-confined; the electron, anchored at the lowest fixed point, is therefore stable.

Confinement as elastic stiffening. In a lattice-of-oscillators analogy, each site carries an effective “colour”. The inter-site elastic energy rises with separation, producing a linearly increasing cost that prevents colour isolation, a classical analogue of QCD confinement. Gluon-like excitations are coherent wave–plus–anti-wave cycles on the links; their exact energy cancellation over a full cycle provides the deterministic counterpart to “virtual-gluon exchange”. A natural corollary is that pure-gluon (glueball) states should be extremely elusive—no unambiguous experimental candidate has yet been confirmed—consistent with this picture.

Anomaly freedom is built-in. Mirror doubling renders the full chiral spectrum vector-like; all perturbative gauge, mixed and gravitational anomalies cancel on any globally-hyperbolic background (Appendix U). With consistency secured (Appendices T–U), the remaining programme focuses on absolute mass scales, higher-loop renormalisation, and extending scattering tests beyond the $e^+e^- \rightarrow \mu^+\mu^-$ benchmark (Appendix S).

3.1.5. Computational Implementation (Summary)

The elastic–spinor coupling enters the discrete momentum update via

$$\dot{\pi} = \rho \partial_t^2 u = T \nabla^2 u - (E_{STM} + \Delta E) \nabla^4 u + \eta \nabla^6 u - \rho \gamma \partial_t u - \lambda u^3 \underbrace{-g \bar{\Psi} \Psi}_{\text{elastic-spinor}},$$

with $\pi = \rho \partial_t u$. We discretise u on a staggered Cartesian grid with a fourth-order central stencil; Ψ is collocated, and gauge fields are stored as link variables $U_\mu(x) = \exp[-i a g A_\mu(x)]$ so that covariant differences are exact on the lattice. The conservative part is advanced by leapfrog; damping/dephasing is included by a local GKSL step (Strang splitting), with jump-operator densities $L_i(\xi)$ chosen to commute with the discrete Gauss operator (or, equivalently, a short orthogonal projection onto the Gauss-law kernel each step).

Stability is enforced by a CFL bound $c \Delta t / \Delta x \leq 0.4$ (empirical margin for the sixth-order term), and we observe second-order temporal and fourth-order spatial convergence under Richardson tests. Monitors include: trace preservation $\text{tr} \rho(t)$, minimum eigenvalue drift (positivity), discrete Gauss residual $\|G\|_2$, and energy decay $\dot{\mathcal{E}} \approx -\int \rho \gamma (\partial_t u)^2$ matching the Rayleigh term in mass-normalised form $-\gamma \partial_t u$. The dissipative step respects graded locality and preserves the CCR/CAR structure (Appendix O, O.H2), consistent with the open-system framework in Appendix T (Thm. T.6).

Rosetta — one physics, three dialects. The results of § 3.1 can be read equivalently as: properties of the pole structure of $G_R(\omega, \mathbf{k})$ (response); statements about the envelope Hamiltonian inferred from the STM dispersion (evolution); or the stationary-phase structure of an effective action S_{eff} for the slow envelope (path sum). The same single-medium origin also fixes $\Delta x \Delta p$ and $\Delta E \Delta t$: they follow from Fourier duality and the envelope's canonical term, with numerical factors set by the STM dispersion data used in § 3.1.1 and Appendix D.

3.2. Nonperturbative Effects

To probe dynamics beyond perturbation theory we use Functional Renormalisation Group (FRG) treatments in two sectors: a $z = 3$ scalar flow for an effective STM-compatible elastic potential (Appendix Y.10), and a deliberately simplified FRG-plus-soliton model (Appendix L).

In the Local Potential Approximation (LPA/LPA') the effective potential $V_k(\phi)$ evolves with RG scale k under a gapped regulator consistent with the STM dispersion (the ∇^6 term provides a natural UV cutoff; cf. Appendix T for positivity/sectoriality). This analysis supports three robust features:

- **Solitons (kinks) and domain walls.** For double- or multi-well V_k , the one-dimensional static Euler-Lagrange equation (derived from the STM energy density with $+\eta |\nabla^3 u|^2$) admits finite-energy kink/domain-wall solutions that interpolate between vacua. These persist under LPA' (field-dependent wave-function renormalisation), with tensions renormalised but signs unchanged.
- **Discrete vacua and generation pattern.** Multiple minima of the coarse-grained potential yield discrete vacua and associated elastic scales. In the conservative single-direction proxy, the triple-well region \mathcal{R}_3 identified in the $z = 3$ scalar truncation (Appendix Y.10) provides a template for three elastic basins with well-separated minima at a finite extraction scale k_* . In the calibrated setting discussed there, additional sectors beyond the scalar-only truncation can materially affect the effective quartic structure at intermediate scales; Yukawa and gauge interactions provide a direct STM-natural route to negative contributions to the quartic running (see §Y.10.7), while multicomponent anisotropy can generate discrete axis vacua in a minimal three-component basin-field truncation (Appendix Y.12). The resulting basin amplitudes, curvatures and inter-basin overlap proxies define the strict flavour-interface export contract specified in Appendix Y.12 for a future predictive pipeline. In the present paper, these FRG-exported basin observables are not used as inputs to the Appendix R mixing scans: Appendix R treats the basin/well data phenomenologically (within STM-calibrated ranges) and constructs the mixing operator Ω from the Appendix R derivations alone. Phase defects associated with kink or domain-wall backgrounds provide deterministic sources for CP phases (see §3.1.4).

- **Core regularisation in collapse analogues.** In spherically symmetric toy models, the short-distance stiffening from the $+\eta\nabla^6 u$ term halts gradient blow-ups and replaces singular cores with finite-amplitude, solitonic/standing-wave interiors. The quadratic energy is positive (Appendix T), so these cores are dynamically stable in the model PDE. A full GR-coupled analysis is deferred to Appendix M's coarse-grained (Einstein-like) limit.

A detailed derivation is in Appendix L, where FRG flows, defect energetics, and vacuum selection are computed consistently with the STM sign conventions of § 2.1.1 (i.e. $+T\nabla^2 u - \tilde{E}\nabla^4 u + \tilde{\eta}\nabla^6 u$ in the mass-normalised PDE). Two caveats keep the discussion conservative:

- **Black-hole thermodynamics.** While core regularisation appears generically in the PDE, a complete derivation of black-hole thermodynamics from those cores is not yet provided. Our covariant, long-wavelength thermodynamic treatment does recover the area law (Appendix F.4):

$$S_{BH} = \frac{A}{4G} + \mathcal{O}\left(\frac{\lambda_c}{R_*}\right),$$

with corrections suppressed by the ratio of a microscopic Compton scale λ_c to the horizon radius R_* . Computing the full Hawking spectrum and a microscopic entropy count for the solitonic cores remains open.

- **Back-reaction & anomalies.** The FRG flows used here preserve the positivity/sectoriality of the linear operator (Appendix T). Coupling to the emergent gauge–spinor sector respects anomaly cancellation by mirror doubling (Appendix U). A full multi-loop FRG with dynamical gauge fields is future work; present conclusions are based on LPA/LPA' with calibrated STM couplings.

Perspective. This non-perturbative picture—stable defects, discrete vacua, UV-tamed flows—provides theoretical motivation and consistency support for the three-basin hypothesis used in the flavour construction of §3.1.4/Appendix R and for the core-regularisation claim used in §4. It stays fully consistent with the damping/open-system framework (Appendix P) and the well-posedness/ghost-freedom proofs (Appendix T). The CKM/PMNS fits themselves are produced entirely within Appendix R (three-mode operator Ω plus the scan/fit code) and do not use FRG outputs from Appendix Y. The same FRG machinery also suggests a route to an absolute STM mass scale: in principle, a fully coupled multi-field FRG flow, anchored to the eight PDE parameters of Appendix K.7, would map the microscopic elastic coefficients into an infrared effective triple-well potential, whose curvatures then fix both the fermion mass scales and the lightest non-zero excitation (the STM mass gap). In this paper we do not carry out that full calculation. Instead we treat the three elastic basins and their associated scales as phenomenological inputs, constrained by the STM calibration window and by the existence of a triple-well region in the $z = 3$ scalar truncation (Appendix Y.10). Appendix Y shows that a multi-basin structure is natural in an STM-consistent FRG proxy and specifies a basin-data export contract (Appendix Y.12) that could, in a later matter-complete implementation, replace these phenomenological inputs by FRG-derived basin observables; a detailed multi-field STM FRG computation of the absolute mass gap is left to future work.

Our treatment here focuses on solitonic structures in the membrane's displacement field. For a complementary perspective showing how these solitons manifest as curvature regularisation in an emergent spacetime geometry, see Appendix M for the Einstein-like derivation.

3.3. Toy Model PDE Simulations

We use numerical experiments to illustrate the core STM dynamics and the emergent spinor structure, working entirely in the nondimensional (ND) formulation consistent with Appendix K.7. Unless stated otherwise the ND couplings are

$$A_2 = 0.10, \quad A_4 = 1.0, \quad A_6 = 0.02, \quad \lambda = 0.13, \quad g_{nd} = 0.3028.$$

Two runs are considered:

- **Undamped:** $\gamma = 0$ with the stiffness reservoir switched off, $\kappa_E = \Gamma_E = 0$.

- **Damped:** $\gamma = 0.01$ with $\kappa_E = \Gamma_E = 0.05$.

Both simulations are implemented in the Python codes listed in Appendix Q and use periodic FFT domains with an optional thin radial PML to suppress wrap-around.

3.3.1. Scalar \rightarrow Spinor Simulation

We solve the 2-D STM PDE on a periodic box of side L (ND units). Time integration uses a stiff-stable split:

- Crank–Nicolson for the ∇^6 term,
- a staggered (leap-frog/Verlet-like) update for ∇^4 , the nonlinear gauge coupling and forcing,
- a **raised-cosine ramp** for the coupling $g(t) = g_{nd} \frac{1}{2} [1 - \cos(\pi \min(1, t/t_{ramp}))]$ (rather than a linear ramp) to avoid exciting high- k modes at start-up.

Initial data seed only the scalar field:

$$u_{prev}(x, y) = \tan h \left(\frac{\sqrt{(x-0.5)^2 + (y-0.5)^2} - R_0}{\sqrt{2}} \right), \psi_1 = \psi_2 = 0,$$

so no spinor is present at $t = 0$. As the evolution proceeds, the nonlinear term $-g_{nd}u |\psi|^2$ pumps the spinor channel. After coarse-graining $u \mapsto P$ and extracting $\partial_t P$ we identify the envelope-level spinor components

$$\Psi_1 \propto P \text{ and } \Psi_2 \propto \partial_t P e^{i\pi/2}$$

(with mirror partners $\bar{\Psi}_i = -\Psi_i$), matching the plots in Figure 1.

Key observations.

Unimodal u (a single “bubble”) generates bimodal $|\Psi_1|$ and $|\Psi_2|$: P is smooth but $\partial_t P$ has two signed lobes, giving two peaks in $|\Psi_i|$. These are envelope features, not separate “particles”. The relative phase $\pi/2$ between Ψ_1 and Ψ_2 is retained in the mirror sectors, evidencing an emergent $U(1)$ phase structure despite seeding only u . Damping $\gamma > 0$ suppresses high-frequency noise; with the implicit ∇^6 step and a sufficiently fine grid/timestep, the conservative limit $\gamma = 0$ is numerically stable. (These $\gamma = 0$ tests are cross-checks; deterministic collapse requires $\gamma > 0$ —see § 3.4.)

3.3.2. STM Schrödinger-Like Envelope

Using the multiple-scale derivation of Appendix D, the slowly varying envelope $A(X, T)$ of the STM membrane displacement satisfies, to next order in the small parameter ϵ ,

$$2i\rho\omega_0\partial_T A - \rho\gamma\partial_T A = k_0^4 \Delta E A + (6E_0 k_0^2 + 15\eta k_0^4) \partial_X^2 A + \dots,$$

where ω_0 and k_0 are fixed by the $\mathcal{O}(1)$ and $\mathcal{O}(\epsilon)$ carrier-dispersion conditions (D.5.1)–(D.5.2). In the conservative limit $\gamma \rightarrow 0$ one recovers the free-particle form

$$i\partial_T A = -\frac{\hbar_{eff}}{2m_{eff}} \partial_X^2 A + V_{eff}(X)A,$$

with explicit STM formulae for \hbar_{eff} , m_{eff} , V_{eff} given in (D.6.2).

Implementation details (Fourier propagation used in figures)

We simulate a standard double-slit aperture $A(x)$, pad by N_{pad} for FFT resolution, compute

$$E(k) = \mathcal{F}\{A\}, I_{ref}(k) = |E(k)|^2,$$

then apply the STM higher-order spatial-propagation phase and (optional) damping:

$$E_{stm}(k) = E(k) \exp \left[-i \left(K_4^{prop} k^4 + K_6^{prop} k^6 \right) z \right] \exp \left[-\frac{\gamma z}{2v_g} \right].$$

See Figure 2 [undamped], Figure 3 [damped].

Plotting/normalisation: for Figures 2–3 we use a common normalisation to the Fraunhofer peak,

$$I_{ref} \leftarrow \frac{I_{ref}}{\max(I_{ref})}, I_{stm} \leftarrow \frac{I_{stm}}{\max(I_{ref})},$$

so that damping remains visible as a reduced STM throughput on the same γ -scale.

Coefficient mapping (propagation picture). Use

$$K_4^{prop} = A_4 L_*^3 (m^3), K_6^{prop} = A_6 L_*^5 (m^5),$$

with the non-dimensional variables $k_{nd} = kL_*$, $z_{nd} = z/L_*$ giving the equivalent phase

$$\exp\left[-i\left(A_4 k_{nd}^4 + A_6 k_{nd}^6\right)z_{nd}\right].$$

The amplitude law follows the envelope result

$$\frac{A_{out}}{A_{in}} \simeq \exp\left(-\frac{\gamma t_{flight}}{2}\right), t_{flight} = \frac{z}{v_g},$$

hence the spatial factor $\exp[-\gamma z/(2v_g)]$.

Key observations

The STM quartic–sextic propagation enters as a pure phase factor in the far-field wavenumber k_x . Consequently, in the undamped case the far-field intensity satisfies

$$I_{stm}(k_x) = |E(k_x)|^2 = I_{ref}(k_x)$$

(up to numerical/FFT discretisation), and Fraunhofer peak positions are retained to better than 10^{-4} . With damping included via the envelope factor $\exp[-\gamma z/(2v_g)]$, the STM intensity acquires an overall throughput reduction

$$I_{stm} \propto \exp\left(-\frac{\gamma z}{v_g}\right);$$

under the common normalisation used for Figures 2–3 this appears as a uniform lowering of the STM curve while fringe geometry remains unchanged. Any residual “jaggedness” in the undamped plots is a finite-grid artefact removable by mild padding increase.

Initial simulations suggested stable spinor configurations without explicit damping; the refined deterministic analysis in §3.4 shows non-zero damping is required for deterministic collapse and proper measurement outcomes.

Symbols. These are propagation coefficients (K_4^{prop}, K_6^{prop}) (length powers) and must not be confused with the SI force-density PDE coefficients $K_4 = E_{STM} L_p^2$ (N) and $K_6 = \eta L_p^4$ (N·m²) used in Appendix K.

Consistency note (dimensions).

Sections 3.3.1–3.3.2, the attached simulations, and the Appendix-K tables are mutually consistent: the PDE solvers and spinor runs are strictly ND; the Schrödinger-like demos use $K_4^{prop} = A_4 L_*^3$, $K_6^{prop} = A_6 L_*^5$ so that $K_4^{prop} k^4 z$ and $K_6^{prop} k^6 z$ are dimensionless), and damping is mapped from time to space via $t = z/v_g$. The Python references for these sections are listed in Appendix Q.

3.4. Measurement Problem and Dynamical Filtering

One of the longstanding puzzles in quantum foundations is the measurement problem: how a linear, deterministic dynamics yields definite outcomes. In STM, “collapse” is reinterpreted as dynamical filtering into basin-of-attraction minima: the analyser sets a conservative bias; the readout supplies dissipation; trajectories flow to stable fixed points. No ad hoc postulate is required.

Scope note (Bell/CHSH): what STM does and does not claim. In §3.4 we explain local outcome formation (basis selection + dissipative filtering) within a deterministic PDE. In Appendix E we

show an entanglement analogue whose CHSH parameter can reach $2\sqrt{2}$. This is not presented as a counterexample to Bell's theorem under its standard assumptions. Rather, STM's Bell phenomenology is framed as relaxing measurement independence at the coarse-grained effective-variable level via contextual boundary conditions and shared preparation (see §3.4.4 and Appendix E.3.5 "Clarification").

Synopsis (one-pixel wins). For single-quantum input, the STM field couples locally to pixels via jump operators $L_i \propto \sqrt{\gamma(\mathbf{x}_i)}\Pi(\mathbf{x}_i)$. The open evolution yields exactly one jump: once any pixel absorbs, the conditional field is vacuum in all channels ("winner-take-all"). Click probabilities obey $p_i \propto \int_{\text{pixel } i} |A(\mathbf{x})|^2 d^2x$, so the two-slit pattern builds up shot-by-shot while each shot yields one click. Multi-quantum inputs (e.g. coherent states) permit multiple clicks with standard bosonic statistics. (See § 3.4.3 for the $|A|^2$ mapping and CHSH.)

3.4.1. Envelope Equation and Elastic Damping

Analysers-local stiffness anisotropy. To represent basis selection by an analyser in the conservative sector, we introduce a small, local anisotropy of the quadratic elastic energy in a thin slab \mathcal{A} :

$$\mathcal{E}_{\text{tens}} \rightarrow \frac{1}{2}T |\nabla u|^2 + \frac{1}{2}\delta T \left[(\mathbf{n} \cdot \nabla u)^2 - (\mathbf{n}_{\perp} \cdot \nabla u)^2 \right] \chi_{\mathcal{A}}(\mathbf{x}), \quad |\delta T| \ll T.$$

Varying this term adds a conservative force

$$\delta \mathcal{F} = \nabla \cdot [\delta T (\mathbf{n} \cdot \nabla u) \mathbf{n} - \delta T (\mathbf{n}_{\perp} \cdot \nabla u) \mathbf{n}_{\perp}] \chi_{\mathcal{A}}(\mathbf{x})$$

alongside the baseline STM terms. Dissipative screen effects are treated separately, via position-selective Rayleigh damping $\gamma(\mathbf{x})$ in a thin absorbing layer. The anisotropy is chosen small enough that it selects a measurement basis without appreciably distorting the free-propagation pattern.

We separate basis selection from irreversibility: the analyser's weak conservative anisotropy selects the local measurement axis (Appendix D, Eq. D.6.12), whereas the position-dependent Rayleigh layer implements the absorbing readout. In this split, the sinusoidal (Born-type) weights are fixed by geometric overlap under the analyser's mode mixing, and the screen converts those weights into click frequencies (Appendix E.3).

Damping calibration. Coarse-graining Planck-time kicks yields a physical damping

$$\gamma_{\text{phys}} \simeq \alpha_d \frac{\hbar}{c^2} \omega_p^3, \quad \omega_p = \sqrt{\frac{c^5}{\hbar G}},$$

with $\alpha_d \sim 10^{-2}$ geometric. Using the reference density ρ and time scale T_0 gives the non-dimensional value used in all scalar simulations,

$$\gamma_{\text{nd}} = \frac{\gamma_{\text{phys}} T_0}{\rho} \simeq 0.010.$$

For position readout we take $\gamma \rightarrow \gamma(\mathbf{x})$ only in a thin high- γ layer at the screen; the local power deposition

$$\dot{W} = \rho \gamma(\mathbf{x}) (\partial_t u)^2$$

provides the detector signal.

Envelope equation. The slow envelope $A(\mathbf{x}, t)$ obeys a complex Ginzburg–Landau / nonlinear Schrödinger-type evolution

$$i\partial_t A = -\frac{1}{2m_{\text{eff}}} \nabla^2 A + (\beta_R + i\beta_I) |A|^2 A - i\gamma_{\text{nd}} A,$$

where $m_{\text{eff}}, \beta_R, \beta_I$ are fixed combinations of $(E_{\text{STM}}, \eta, \lambda)$. A small positive $\beta_I > 0$ balances linear loss and drives $|A|$ towards

$$|A|_{ss} = \sqrt{\frac{\gamma_{nd}}{\beta_I}},$$

preventing secular growth in the reduced description. Spinor fields couple via the Yukawa term, so we include a milder envelope-level dephasing $\partial_t \Psi \rightarrow \partial_t \Psi - \gamma_f \Psi$ with $\gamma_f = \frac{1}{2}\gamma$, giving $\gamma_{f,nd} = 0.005$. Varying α_d by $\pm 20\%$ shifts both rates proportionally; CKM/PMNS/seesaw predictions move by $< 10^{-4}$. With $\gamma_{nd} \simeq 0.010$ and $\beta_I > 0$, the envelope filters any initial superposition into stable attractors, setting the stage for the deterministic measurement mechanism described below.

3.4.2. Phase-Space Picture and Basins of Attraction

Let a two-mode local excitation have components u_1, u_2 at fixed radius r . Writing $u_1 = r \cos \theta$, $u_2 = r \sin \theta$, the analyser induces an effective conservative potential

$$V_{meas}(\theta) = V_0 \cos^2(\theta - \theta_{meas}), V_0 \sim \frac{1}{2} \delta T k^2 |U|^2 \times (\text{slab factor}),$$

derived in Appendix D.6 from δT in \mathcal{A} . With mild angular damping $\gamma_\theta > 0$, the phase dynamics reduce to overdamped gradient flow on the unit circle,

$$\dot{\theta} = -\partial_\theta V_{meas}(\theta) = 2V_0 \sin(\theta - \theta_{meas}) \cos(\theta - \theta_{meas}),$$

so θ relaxes to one of the two minima $\theta = \theta_{meas}$ or $\theta_{meas} + \pi$ (the analyser's two outcomes). Figure 4 illustrates trajectories spiralling into the nearest minimum under γ_{nd} and the nonlinear feedback encoded by V_{meas} . In this separation of roles, the analyser sets the basis via V_{meas} ; the screen/readout supplies $\gamma(\mathbf{x})$ and finalises a single macroscopic click.

Specialisations.

- *Stern–Gerlach*. The inhomogeneous magnetic field fixes θ_{meas} ; dissipation is confined to the readout regions.
- *Polariser / PBS*. A polariser sets θ_{meas} in the (H, V) plane; a polarising beam splitter is conservative (unitary splitter) with damping in the two detectors. Malus' law $P(H | \theta_0) = \cos^2(\theta_{meas} - \theta_0)$ follows from the same potential.
- *Screen (position)*. For a position-sensitive screen we use a thin absorbing layer

$$\gamma(\mathbf{x}) = \gamma + \gamma_{scr} \chi_\delta(x - x_{scr}), \gamma_{scr} > 0,$$

and

$$\dot{W} = \rho \gamma(\mathbf{x}) (\partial_t u)^2$$

integrated over pixels. The first pixel to reach threshold flags the event.

Footnote: Here $L(x) \propto \sqrt{\gamma(x)} \Pi(x)$ is the local Lindblad choice (ghost-number zero, BRST-safe); Π is the local pixel/voxel projector, distinct from $\pi = \rho \partial_t u$ (Appendix P, §P.6).

3.4.3. From Deterministic Filtering to Born-Rule Statistics

For a two-mode degree of freedom whose internal ellipse angle is θ_0 , an analyser set to θ_{meas} defines the measurement basis and the small- γ filtering dynamics selects one of two stable outcomes. In the coarse-grained description this yields the familiar overlap weights $P(+1 | \theta_{meas}, \theta_0) = \cos^2\left(\frac{\theta_{meas} - \theta_0}{2}\right)$ and $P(-1 | \theta_{meas}, \theta_0) = \sin^2\left(\frac{\theta_{meas} - \theta_0}{2}\right)$. If θ_0 is effectively uniform over runs, $\theta_0 \in [0, 2\pi)$, then averaging over θ_0 gives unbiased single-wing marginals $P(+1 | \theta_{meas}) = \int_0^{2\pi} \frac{d\theta_0}{2\pi} P(+1 | \theta_{meas}, \theta_0) = \frac{1}{2}$ and $P(-1 | \theta_{meas}) = \frac{1}{2}$. Importantly, this uniform micro-phase sampling is not responsible for the Bell-type correlation itself. The latter arises because the prepared two-wing excitation is a single non-separable global mode, rather than two independent subsystems with separable ontic states.

Two-wing correlations (what is being averaged). For a correlated pair with shared internal phase information arising from the joint preparation, the coarse-grained correlation can be written as an average over unresolved microstate variables (here parameterised by θ_0). In the simplified overlap model, for two analysers at a and b acting on a correlated pair with shared θ_0 , one obtains a sinusoidal dependence on the relative setting,

$$E(a, b) \propto \int_0^{2\pi} \frac{d\theta_0}{2\pi} f(a, \theta_0) f(b, \theta_0) \sim \cos(a - b),$$

while for a singlet-like preparation the correlation is $E(a, b) = -\cos(a - b)$. (See Figure 5 for the sinusoidal dependence, and Appendix E.5 for the corresponding CHSH construction.)

Within STM, the apparent randomness is epistemic at the coarse-grained level: the ensemble over θ_0 reproduces Born-rule statistics, while each individual run is governed by deterministic filtering into one of the two attractor basins defined by the local measurement setting. Any GKSL/Lindblad representation used for the reduced dynamics is therefore interpretive rather than foundational, i.e. an effective coarse-grained description of the same deterministic filtering and thresholded absorption mechanism.

Scope note (what the uniform θ_0 assumption does — and does not do). The assumption that θ_0 is effectively uniform over runs is used here to recover unbiased single-wing marginals and the overlap-model expectations above. It is not, by itself, a mechanism for CHSH/Tsirelson saturation. The CHSH result in Appendix E relies on (i) a non-factorisable shared preparation (Appendix E.3.1) and (ii) contextual branch-selection/routing under boundary conditions (Appendix E.3.4), with the associated Bell-assumption scope discussed in §3.4.4.

STM's Bell phenomenology should therefore be read as contextual at the coarse-grained level: the effective hidden variables reflect the shared preparation and the local boundary conditions implemented by the analysers (including finite-memory decoherence in each wing), so the strict factorisation premises used in the standard Bell derivation do not apply in the same way. The underlying PDE remains local and no-signalling; there is no superluminal influence at readout and no conspiratorial coordination between human setting choices and microscopic elastic variables.

3.4.4. Assumptions and Local Constraints (Measurement Sector)

We collect here the key assumptions underlying the measurement-sector construction and indicate how they may be constrained or falsified.

- **Tiny analyser anisotropy (basis selection only).**

$$|\delta T| \ll T, V_0 \sim \frac{1}{2} \delta T k^2 |U|^2 \times (\text{slab factor}).$$

Potential falsifier: if no measurable pre-screen drift or port bias ε is observed, then $|V_0| < \varepsilon$, implying

$$|\delta T| < \frac{2\varepsilon}{k^2 |U|^2} \times (\text{slab factor})^{-1}.$$

- **Thin analyser relative to diffraction length.**

The analyser thickness L_A must be small compared with the diffraction length,

$$L_A \ll L_{\text{diff}} = \frac{2m_{\text{eff}}}{k\Delta k},$$

so that it selects a basis without significantly altering the free-space interference pattern.

- **Conservative analyser (no outcome forcing).**

Angular damping within the analyser satisfies

$$\gamma_\theta \tau_A \ll 1,$$

where τ_A is the transit time through A . Dissipation is then confined to the readout/screen region, ensuring that the analyser's role is basis selection rather than direct outcome biasing.

These conditions make the measurement sector experimentally constrainable: deviations from the predicted drift, fringe pattern, or click statistics bound the admissible parameter ranges. Broader discriminating tests are summarised in §3.4.5.

Interpretive scope (separability). Accordingly, the present construction should not be read as a separable hidden-variable model with pre-assigned local ellipse orientations; the “opposite ellipse” relation is a constraint on a single joint excitation, not on two independent local states. Within STM, particle localisation refers to the spatial concentration of energy at readout, not to factorisation of the underlying excitation. Localised oscillations may therefore represent spatially separated lobes of a single constrained membrane mode rather than independent subsystems. This is particularly natural in entanglement experiments, where the pair is created by a single interaction region (e.g. a common decay or scattering event), which in STM generically prepares a joint excitation with a fixed internal relation (such as the opposite-ellipse constraint). Subsequent spatial separation corresponds to the unfolding of this joint mode into two localised readouts, not to dynamical separation into independent states. On this view, separability is not an axiom but a dynamical question of whether the shared constraint survives propagation and coarse-graining. An explicit construction of the corresponding non-separable effective state is given in Appendix E.2.

3.4.5. Falsifiability and Discriminating Predictions

The STM account of Born statistics and Bell-type correlations is not purely interpretive: it makes parameterised, falsifiable predictions tied to the readout sector, even though the absolute magnitude of the Rayleigh damping coefficient γ is not yet fixed (and is expected to be constrained by forthcoming interferometric measurements). First, STM predicts strict no-signalling at the operational level: since clicks arise from local basis selection followed by local irreversible absorption, single-wing statistics must be independent of the remote analyser setting; any reproducible dependence $P(A | a, b) \neq P(A | a)$ would falsify the construction, and can be assessed within modern loophole-reduced Bell-test protocols [24]. Second, STM's Born mechanism is explicitly detector-energetic: with absorbed power density $p_{abs}(\mathbf{x}) = \rho\gamma(\mathbf{x})(\partial_t u)^2$, STM predicts that controlled changes to the effective absorption profile reweight detection counts in a definite way in the linear absorption regime,

$$\text{counts}(\mathbf{x}) \propto \int dt \rho\gamma(\mathbf{x})(\partial_t u)^2 \propto \gamma(\mathbf{x}) |\Psi(\mathbf{x})|^2.$$

Accordingly, observation of invariant $|\Psi|^2$ histograms under deliberately patterned changes to $\gamma(\mathbf{x})$ (in a regime where absorption is demonstrably rate-limiting) would falsify the proposed readout mechanism; closely related experimental traditions already treat detector nonlinearity/response calibration as a primary systematic when bounding deviations from Born structure via higher-order interference tests [25]. Finally, since Bell correlations are attributed here to non-separable joint preparation rather than separable local states, sufficiently strong phase scrambling or mode-mixing applied to one wing is expected to degrade the Bell correlation while leaving single-wing statistics intact; establishing a quantitative coherence budget remains an open task, but the qualitative dependence provides a clear discriminator. (For comparison, threshold/energy-accumulation detector models have also been analysed as routes to Born-like click statistics, underscoring that detector physics can be made experimentally discriminating rather than merely interpretive [26].)

3.4.6. Summary

STM recovers Born-rule statistics from deterministic filtering plus local absorption: analyser-defined basis rotation yields overlap weights $P(\pm 1 | \theta_{meas}, \theta_0)$, with unbiased single-wing marginals obtained when θ_0 is effectively uniform over runs. Bell/CHSH behaviour is not attributed to θ_0 -averaging but to non-separable joint preparation (a single constrained global excitation), with outcome definiteness realised by local irreversible absorption and the $|\Psi|^2$ weighting grounded in deposited

power $\int dt \rho \gamma(x) (\partial_t u)^2$. §3.4.4 discusses Bell-assumption scope; §3.4.5 gives discriminating tests (no-signalling, absorber-profile reweighting, and controlled decoherence sensitivity).

3.5. Parameter Constraints and Stability Observations

In exploring the STM PDE numerically—both in the full 2 D scalar + spinor runs and in our 1D double-slit far-field test we use the non-dimensional constants fixed by §2.10/App. K.7, which appear to provide stable, well-behaved solutions:

All non-dimensional constants ($E_{4,nd}$, η_{nd} , β , γ_{nd} , g_{nd} , λ_{nd}) are fixed by the Planck-anchored calibration in Appendix K.7.

3.5.1. Envelope Locking

In the reduced, multiple-scale (“envelope”) approximation (Appendix D), the slowly varying amplitude $A(x, t)$ of a carrier wave satisfies

$$\frac{\partial A}{\partial t} + v_g \frac{\partial A}{\partial x} = \beta |A|^2 A - \gamma_{nd} A,$$

where $v_g = \partial\omega/\partial k$ is the group velocity (see D.5.1). Under homogeneous boundary conditions ($\partial_t A = \partial_x A = 0$), the steady-state amplitude is

$$|A|_{ss} = \sqrt{\frac{\gamma_{nd}}{\beta}}.$$

Hence, for $\beta > 0$, a small positive γ_{nd} is required to balance nonlinear growth and lock the envelope to a finite amplitude:

$$\beta > 0 \implies \gamma_{nd} > 0.$$

While this condition arises within the multiple-scale (envelope) approximation, recent theoretical developments (Section 3.4, Appendix P) establish that a small but non-zero damping term is also physically necessary in the full STM framework to realise deterministic decoherence and recover the Born rule. Although numerical integrations of the undamped STM wave equation ($\gamma = 0$) remain formally stable and self-adjoint under modern schemes (e.g., Crank–Nicolson, BDF), such conservative dynamics do not reproduce collapse or measurement outcomes. Therefore, while envelope-level damping offers a simplified model of amplitude locking, the complete physical theory now supports the presence of a small $\gamma > 0$ as essential for matching phenomenology.

(For units and mappings between γ and γ_1 see App. K.6; for BRST-safe open-system realisations see App. T/P.)

3.5.2. Spinor Stability (Conservative Solver)

Toy-model simulations using the conservative spinor–lump solver of Appendix O (no Rayleigh damping, no pump/relax term; cf. STM_spinor_undamped.py) show that the non-dimensional quartic self-coupling λ_{nd} (LAM) and gauge-like coupling g_{nd} (G_ND) do not need to lie in a very narrow window in order to avoid runaway behaviour. On 2D grids with $L = 8$, $N = 32$ –64 and evolution times up to $t_{nd} \sim 5$ –10, solutions remain well behaved for at least

$$0 \leq \lambda_{nd} \lesssim 1, \quad 0 \leq g_{nd} \lesssim 1,$$

with the calibrated choice $(\lambda_{nd}, g_{nd}) = (0.13, 0.3028)$ lying comfortably inside this region. Across this range the membrane field settles to $\max_{x,y} |u| \sim 0.8$ and rms amplitudes $A_{rms} \sim 10^{-2}$, with no evidence of numerical blow-up or unbounded spinor growth. For reproducible simulations it is therefore sufficient to restrict to a moderate coupling regime, e.g.

$$0 \leq \lambda_{nd} \lesssim 0.5, \quad |g_{nd}| \lesssim 0.5,$$

which leaves a generous safety margin around the calibrated values while avoiding unnecessarily extreme choices.

3.5.3. Double-Slit Interference Constraints

Let $k_s = 2\pi/\lambda_{light}$ be the central diffraction wavenumber for light of wavelength λ_{light} . Two conditions guarantee high-contrast Fraunhofer fringes:

- **UV regulator:**

$$E_{4,nd}k_s^4 + \eta_{nd}k_s^6 \ll \frac{\hbar_{eff}k_s^2}{2m_{eff}}.$$

- **Damping over flight time:** With time-of-flight $T_{TOF} \approx \frac{Zm_{eff}}{\hbar_{eff}k_s}$, one requires

$$\gamma_{nd}T_{TOF} \ll 1,$$

so that fringe contrast is not visibly degraded even for metre-scale propagation distances Z . A weak analyser slab placed *before* the screen may be taken in the $V_0 \rightarrow 0^+$ limit to avoid pre-detection distortion; basis selection then occurs via unitary mixing, with outcome set by the screen's $\gamma(\mathbf{x})$.

3.5.4. Practical Takeaways

For robust, high-contrast STM-PDE simulations, ensure that:

- **Envelope lock:** Choose β and γ_{nd} of the same sign so that $|A|_{ss} = \sqrt{\gamma_{nd}/\beta}$ is well defined.
- **Gauge/self-coupling window (conservative solver):** Parameter sweeps of the undamped spinor-lump code show stable evolution for moderate non-dimensional couplings $0 \leq \lambda_{nd} \lesssim 1$ and $0 \leq g_{nd} \lesssim 1$. As a conservative default we recommend $|g_{nd}| \lesssim 0.5$ and $0.01 \lesssim \lambda_{nd} \lesssim 0.5$, with the calibrated choice $g_{nd} \simeq 0.303$, $\lambda_{nd} = 0.13$ well inside this window.
- **UV regulator check:** Verify $E_{4,nd}k_s^4 + \eta_{nd}k_s^6 \ll \hbar_{eff}k_s^2/(2m_{eff})$.
- **Damping constraint:** Keep $\gamma_{nd}T_{TOF} \ll 1$.

Adherence to these guidelines reproduces stable envelopes, bounded spinor amplitudes and pristine interference patterns across all toy-model tests.

Benchmark inequalities. The solver set from Appendix K ($g \simeq 0.303, \lambda = 0.13$) refers to the calibrated physical couplings and satisfies the Appendix V.9 bounds obtained from representative global-fit scales, namely $M_{mir} \gtrsim 0.062TeV$, $|\lambda\xi| \lesssim 0.444$, $\lambda \lesssim 0.141$. Appendix W.3 shows that one-loop running to m_Z shifts the pure-gauge Wilsons by only $\mathcal{O}(10\%)$, leaving these inequalities intact.

3.6. Validation of Emergent Electroweak Amplitudes

To demonstrate that STM's emergent gauge structure reproduces Standard-Model results, we compute the tree-level cross-section for $e^+e^- \rightarrow \mu^+\mu^-$, including photon exchange and γ -Z interference, with the fine-structure constant run via leptonic vacuum polarisation:

- **Running coupling:** $\alpha(s) = \alpha_0/[1 - (\alpha_0/3\pi)\ln(s/m_e^2)]$ (leptonic VP; adequate at the 10^{-3} level for our benchmarks).
- **Pure-QED benchmark:** the differential cross-section $\alpha(s)^2/(4s)(1 + \cos^2\theta)$ is reproduced by the STM code.
- **Electroweak interference:** adding Z-exchange in the s -channel with $g_V^f = -\frac{1}{2} + 2\sin^2\theta_W$, $g_A^f = -\frac{1}{2}$, and $\sin^2\theta_W = 0.23126$ yields the familiar γ -Z pattern.

Numerics (differential, at $\theta = 90^\circ$) $\sqrt{s} = 10 GeV$: $\sigma_{\gamma+Z}/\sigma_\gamma \approx 0.999$ $\sqrt{s} = 43 GeV$: $\sigma_{\gamma+Z}/\sigma_\gamma \approx 0.992$

These are consistent with PETRA/PEP phenomenology (e.g. CELLO 0.98 ± 0.04). This provides a stringent check that STM's single PDE, when coarse-grained to its emergent Lagrangian, reproduces classic electroweak amplitudes within experimental uncertainties. See Supplementary `Scattering_amplitude.py`.

Using the Appendix Y bound, the STM \rightarrow YM matching below Λ_{match} guarantees that our tree-level $e^+e^- \rightarrow \mu^+\mu^-$ calculation (App. S) is the correct low-energy limit up to $\mathcal{O}(p^2/\Lambda_{match}^2)$ effects.

Note. The ratios above refer to the differential cross-section at 90° . The integrated (total) cross-section shows slightly different per-cent-level shifts because the $\cos\theta$ interference term averages differently.

3.7. Zitterbewegung from Face–Mirror Antisymmetry (and Why Its Frequency Matches Dirac)

Summary. The antisymmetric mirror face enforces two normal modes (symmetric/antisymmetric). Any face-localised state necessarily mixes them, producing a fast “trembling” with frequency set by the spectral gap. The result is the same as Dirac's $2E/\hbar$ prediction, with c replaced by the membrane light-cone speed c_T ; damping only attenuates the envelope.

Conservative two-face Hamiltonian and normal modes

Place the spinors on the two faces of the membrane: ψ (top), χ (mirror). The scalar displacement is odd across the mid-plane, $u^{(+)} = -u^{(-)}$, so the inter-face coupling is antisymmetric. At quadratic order (suppressing gauge terms for clarity) and for a uniform background u_0 ,

$$\mathcal{H}_{cons} = \psi^\dagger \left(-i c_T \mathbf{ff} \cdot \nabla + m_{eff} \beta \right) \psi + \chi^\dagger \left(-i c_T \mathbf{ff} \cdot \nabla + m_{eff} \beta \right) \chi + \kappa u_0 (\bar{\psi} \chi + \bar{\chi} \psi),$$

with $E(\mathbf{k}) \equiv \sqrt{c_T^2 k^2 + m_{eff}^2}$. For plane waves $e^{i(\mathbf{k}\cdot\mathbf{x} - \omega t)}$,

$$i\partial_t \begin{pmatrix} \psi \\ \chi \end{pmatrix} = \begin{pmatrix} h_D(\mathbf{k}) & \kappa u_0 \\ \kappa u_0 & h_D(\mathbf{k}) \end{pmatrix} \begin{pmatrix} \psi \\ \chi \end{pmatrix}, \quad h_D(\mathbf{k}) = c_T \mathbf{ff} \cdot \mathbf{k} + m_{eff} \beta.$$

Diagonalising with $\psi_\pm = (\psi \pm \chi)/\sqrt{2}$ gives two normal modes with

$$\omega_\pm(\mathbf{k}) = \pm E(\mathbf{k}).$$

Interference, trembling, and the Dirac frequency

A face-localised initial state,

$$\Psi(0) = a_+ \psi_+ + a_- \psi_-, \quad |a_+|^2 + |a_-|^2 = 1,$$

evolves as

$$\Psi(t) = a_+ \psi_+ e^{-iEt/\hbar} + a_- \psi_- e^{+iEt/\hbar}.$$

Any one-face observable contains a cross term $\propto e^{-i(E-(-E))t/\hbar} = e^{-i2Et/\hbar}$. The zitterbewegung frequency is therefore

$$\omega_{ZB}(\mathbf{k}) = \frac{2E(\mathbf{k})}{\hbar} = \frac{2}{\hbar} \sqrt{c_T^2 k^2 + m_{eff}^2},$$

so at rest $\omega_{ZB} = 2m_{eff}/\hbar$, and ultra-relativistically $\omega_{ZB} \approx 2c_T |k|/\hbar$.

Interpretation. Dirac attributed ZB to interference of positive/negative energy components of a single spinor. Here, the same spectral gap $2E$ arises from symmetric vs antisymmetric face–mirror modes enforced by the antisymmetric domain. The conservative spectrum remains bounded below; the algebra of the interference—and hence the frequency—is identical (with $c \rightarrow c_T$).

Amplitude vs frequency

The amplitude on one face scales with the inter-face admixture (set at rest by $|\kappa u_0|$):

$$\mathbf{A}_{ZB} \propto \frac{|\langle \psi_- | \hat{\mathcal{O}} | \psi_+ \rangle|}{E(\mathbf{k})}$$

By contrast, the frequency depends only on the gap $2E(\mathbf{k})$ and is independent of the mixing strength once $\mathbf{k} \neq 0$.

Heisenberg check (velocity operator)

With $H = c_T \mathbf{ff} \cdot \mathbf{k} + m_{eff} \beta$ on one face,

$$\frac{d\hat{\mathbf{x}}}{dt} = c_T \mathbf{ff}, \quad \frac{d\mathbf{ff}}{dt} = \frac{i}{\hbar} [H, \mathbf{ff}] = -\frac{2i}{\hbar} (H \mathbf{ff} - c_T \mathbf{k}).$$

Solving,

$$\mathbf{ff}(t) = \frac{c_T \mathbf{k}}{E} + \left(\mathbf{ff}(0) - \frac{c_T \mathbf{k}}{E} \right) e^{-2iEt/\hbar},$$

and integrating once gives

$$\langle \mathbf{x}(t) \rangle = \mathbf{x}_0 + \mathbf{v}_g t + \underbrace{\frac{\hbar}{2E} \left(\mathbf{ff}(0) - \frac{c_T \mathbf{k}}{E} \right)}_{\mathbf{A}_{ZB}} \sin\left(\frac{2Et}{\hbar} + \phi_0\right),$$

confirming $\omega_{ZB} = 2E/\hbar$ (with $c \rightarrow c_T$) from the operator route.

Damping and coarse-graining

Open-system effects enter as scalar Rayleigh damping (mass-normalised) $-\gamma \partial_t u$ and spinor dephasing $\gamma_f \simeq \frac{1}{2}\gamma$. To leading order in γ_f , the ZB envelope decays while the frequency is unchanged:.

$$\text{ZB}(t) \sim \mathbf{A}_{ZB} e^{-\gamma_f t} \sin\left(\frac{2Et}{\hbar} + \phi_0\right), \quad \omega_{ZB} = \frac{2E}{\hbar} \text{ (no first-order shift)}$$

Coarse-graining over $t \gg \gamma_f^{-1}$ leaves a residual microscopic phase in cross-face bilinears; this seeds the complex phases used later in the Yukawa/CP structure.

In particular, the zitterbewegung frequency $\omega_{ZB} = 2E/\hbar$ should be understood as the beat frequency between coarse-grained symmetric and antisymmetric face-mirror modes, not as the fundamental oscillation rate of the underlying STM membrane, whose sub-Planck carrier oscillations are much faster and have already been integrated out.

Cross-references. Definitions of c_T , m_{eff} , and units: §2.1.1. Damping conventions: §3.4.1. Solver form and mappings: Appendices K.6–K.7. Positivity/sectoriality: Appendix T.

3.8. Summary

- **Effective Schrödinger-like dynamics** By coarse-graining the rapid, sub-Planck oscillations in $u(\mathbf{x}, t)$, we obtain a slowly varying envelope $A(\mathbf{x}, t)$ that obeys an effective Schrödinger equation. This reproduces interference phenomena and a deterministic Born-rule interpretation without invoking intrinsic randomness.
- **Emergent gauge symmetries** A bimodal decomposition of the displacement field produces a two-component spinor $\Psi(\mathbf{x}, t)$. Enforcing local phase invariance on Ψ yields $U(1)$, $SU(2)$ and $SU(3)$ gauge fields as collective elastic modes, giving deterministic analogues of photons, W/Z bosons and gluons.
- **Direct PDE validation** Section 3.3 showed that the full STM PDE—with higher-order dispersion terms and no explicit damping ($\gamma = 0$)—remains self-adjoint and numerically stable under modern implicit schemes (e.g. Crank–Nicolson). Toy-model simulations reproduce emergent spinor wave packets and Fraunhofer fringes, confirming the core STM dynamics in a fully

conservative setting. Electroweak benchmark (differential at $\theta = 90^\circ$): $\sigma_{\gamma+Z}/\sigma_\gamma \approx 0.999$ at $\sqrt{s} = 10 \text{ GeV}$, ≈ 0.992 at $\sqrt{s} = 43 \text{ GeV}$; consistent with PETRA/PEP phenomenology.

- **Zitterbewegung from face–mirror antisymmetry (Dirac frequency)** The antisymmetric mirror face enforces symmetric/antisymmetric normal modes. Any face-localised state necessarily mixes them and exhibits trembling at

$$\omega_{ZB} = \frac{2E(\mathbf{k})}{\hbar}, E(\mathbf{k}) = \sqrt{c_T^2 k^2 + m_{eff}^2},$$

- i.e. Dirac’s result with $c \rightarrow c_T$. The amplitude scales with inter-face mixing $\propto |\kappa u_0|/E$; the envelope decays as $e^{-\gamma_f t}$ with $\gamma_f \simeq \frac{1}{2}\gamma$. Damping attenuates ZB but does not shift ω_{ZB} at leading order.
- **Stability and interference constraints** In the envelope approximation (Section 3.5) we derived practical parameter windows: envelope locking requires $\gamma > 0$ only to arrest secular growth in the reduced model; spinor stability demands $g_{nd} \lesssim 0.1$ and $\lambda_{nd} \gtrsim 10^{-2}$; high-fidelity interference requires $E_{4,nd} k_s^4 + \eta_{nd} k_s^6 \ll \hbar_{eff} k_s^2 / (2m_{eff})$ and $\gamma T_{TOF} \ll 1$.
- **Non-Markovian decoherence and Bell violations** Integrating out fast modes via a Feynman–Vernon influence functional yields a non-Markovian master equation whose memory kernel produces deterministic wavefunction collapse. Spinor-based measurements recover Bell-inequality violations (up to $2\sqrt{2}$) without any stochastic postulates.
- **Fixed points and solitonic cores** Perturbative RG and FRG analyses, supported by the sextic regulator, reveal discrete renormalisation-group fixed points that naturally account for three fermion generations. Non-perturbative solutions include stable, finite-amplitude solitonic cores that avert curvature singularities in black-hole analogues.

4. Discussion

With these central results established, we now explore their broader significance. In particular, we examine how deterministic elasticity underpins quantum-like behaviour and gauge interactions, reassess the interpretation of spacetime singularities and dark energy, and outline concrete avenues for experimental validation and further theoretical development.

Incorporating this Hamiltonian-to-commutator derivation into the STM framework anchors the quantum postulate firmly in the same continuum elasticity that gives rise to gravity and gauge fields. By showing that the canonical commutation relations follow directly from the membrane’s classical symplectic structure—rather than being an auxiliary assumption—we close the conceptual loop: the familiar non-commutativity of \hat{u} and $\hat{\pi}$ is a direct consequence of deterministic elasticity, and no separate “quantisation machinery” is required.

The STM model illustrates how deterministic, classical chaos in membrane oscillations, in the idealised set-ups studied here, can reproduce key quantum-like phenomena such as interference patterns, apparent wavefunction collapse and Born-rule statistics. Within this scope, the deterministic elasticity provides a concrete physical reinterpretation of quantum randomness and suggests that explicit stochastic postulates may be dispensable.

One operator, three dialects. The main claims of § 4 can be viewed equivalently as statements about $G_R(\omega, \mathbf{k})$ ’s poles (response), about the envelope Hamiltonian inferred from STM’s dispersion (evolution), or about the stationary-phase structure of an effective action S_{eff} (path sum). In this sense, Schrödinger dynamics, propagators, the Feynman sum and the standard uncertainty bounds are complementary emergent descriptions of the same single-medium physics.

Measurement synopsis. The double-slit story reads naturally in STM: a weakly open readout layer implements local damping; the envelope obeys Schrödinger transport between analyser and screen; a single jump (pixel) terminates each one-quantum run, and the Born-rule pattern arises from many runs. This reproduces familiar phenomenology while keeping all ingredients (transport, damping, click statistics) inside STM’s deterministic-plus-open framework.

The model represents a bold attempt to unify gravitational curvature with quantum-like phenomena within a single deterministic framework based on high-order elasticity. By incorporating second-, fourth-, and sixth-order spatial derivatives, scale-dependent parameters, and non-Markovian effects, we find that many hallmark features of quantum field theory can emerge naturally from the membrane's classical dynamics.

Falsifiability of elastic gravity. Three complementary probes constrain $(c_g, \alpha_2, \alpha_3, \Lambda_g)$: (i) multimessenger bounds on $c_g = \sqrt{T/\rho}$; (ii) broadband inspiral phasing with distinct f^2 and f^4 trends from the dispersion series; and (iii) black-hole ringdown shifts scaling as $(\omega/\Lambda_g)^{2.4}$. Solar-System and laboratory tests limit the short-range Yukawa components set by $\uparrow_1 \sim \sqrt{K_4/T}$ and $\uparrow_2 \sim (\eta/T)^{1/2}$, complementing the GW constraints.

Below, we examine the implications of these findings, compare them with standard quantum field theory, and consider practical routes toward experimental validation.

4.1. Emergent Quantum Dynamics and Decoherence

Building on the deterministic sub-Planck filtering mechanism of Section 3.4, we now turn to the broader phenomenology of the STM framework.

Throughout this section the calculations should be read as illustrative templates—worked out explicitly in simple laboratory geometries (planar screens, homogeneous baths) with parametrised kernels—rather than as exhaustive or unique derivations of all quantum-mechanical decoherence phenomena. The underlying STM evolution and causality structure have been established on generic globally hyperbolic backgrounds (Appendices O, T, X); here we specialise to idealised set-ups to keep the decoherence story transparent.

A key aspect of our perturbative analysis is that by coarse-graining the rapid, sub-Planck oscillations of the membrane's displacement field $u(x, t)$, one obtains a slowly varying envelope $\Psi(x, t)$. This envelope obeys an effective Schrödinger-like equation,

$$i\hbar \frac{\partial \Psi}{\partial t} = - \frac{\hbar^2}{2m_{eff}} \nabla^2 \Psi + V_{eff}(x) \Psi,$$

mimicking the familiar quantum mechanical form. Crucially, the sixth-order spatial derivative $\nabla^6 u$ in the STM wave equation dampens short-wavelength modes, ensuring that ultraviolet divergences do not arise. Moreover, the Born rule emerges through deterministic sub-Planck chaos in the presence of the finite damping, fixed in Section 3.4, replacing the postulated randomness of conventional quantum theory.

The refined STM now couples IR explanatory power to UV discipline: $z = 3$ scaling with super-renormalisable interactions, reflection positivity and positive spectral measures for gauge-invariant composites, and a rigorous YM decoupling bound that legitimises our SMEFT usage. This places STM closer to asymptotically safe/Lifshitz-type programmes in spirit, while retaining a single classical elastic field with emergent gauge/spinor content rather than fundamental quantum fields.

By splitting $u(x, t)$ into a system component u_S and an environment u_E , we further showed that non-Markovian decoherence follows from integrating out the fast modes u_E .

This framework reproduces the suppression of off-diagonal density-matrix elements through its finite memory kernel; collapse to a definite outcome follows only when the small Planck-time-scale damping term γ is included, as shown in Section 3.4, all within a deterministic PDE context. Notably, as soon as we implement spinor-based measurement operators and allow for correlated sub-Planck modes, the model achieves Bell-inequality violations (CHSH up to $2\sqrt{2}$) via the same small- γ attractor mechanism in a purely classical wave setting.

Although the STM framework now reproduces the Born rule, approaches the Tsirelson bound and predicts laboratory decoherence rates within current uncertainties, mainstream interpretations still regard quantum randomness as fundamental. Future work must verify that the deterministic, small- γ damping mechanism remains consistent with all phenomena—including kilogram-scale

macroscopic superpositions, loop-level anomalies and ultra-long-baseline phase coherence—before STM can plausibly be regarded as a viable deterministic alternative to indeterministic quantum theory.

4.2. Emergence of Gauge Symmetries and Virtual Boson Reinterpretation

Through a bimodal decomposition of the displacement field, the STM model constructs a spinor $\Psi(x, t)$. Requiring local phase invariance on Ψ naturally introduces gauge fields corresponding to $U(1)$, $SU(2)$, or $SU(3)$ [16]. Consequently, photon-like and gluon-like excitations arise as deterministic wave modes rather than quantum fluctuations. Meanwhile, the usual concept of virtual bosons—pertinent to standard quantum field exchanges—is replaced by wave-plus-anti-wave oscillations that transfer no net energy over a full cycle [15]. This classical reinterpretation preserves energy conservation at every instant and bypasses the notion of “transient particle creation,” typical of conventional perturbation theory.

This reinterpretation also clarifies how force mediation, in particular electromagnetism and the strong interaction, can be understood as elastic “connections” in a high-order continuum. The STM PDE itself underlies these gauge fields once spinor local symmetries are introduced. Thus, standard gauge bosons like photons, W^\pm , or gluons appear as coherent membrane oscillations, illustrating how quantum-like gauge interactions might emerge from deterministic elasticity.

For the strong force specifically, visualising the membrane as a chain or lattice of linked oscillators clarifies how confinement arises deterministically from classical elasticity. Each lattice site can be regarded as carrying a colour charge, and the coupling between these sites stiffens with separation to approximate a linear potential. This property prevents the separation of colour charges into free isolated states, directly mimicking the linear potential and confinement behaviour central to QCD. Deterministic gluon-like excitations, represented by coherent waves propagating along oscillator links, thereby mediate the strong interaction without requiring intrinsic randomness or virtual particle fluctuations.

While this approach elegantly reinterprets gauge fields, verifying quantitative equivalence with the Standard Model’s scattering amplitudes and loop processes is crucial; tree-level electroweak amplitudes are now covered, (see § 3.6; Appendix S for differential cross-sections at $\theta = 90^\circ$); loop-level and purely gluonic channels remain to be verified. Detailed calculations would need to show that these “wave-anti-wave” cycles match Feynman diagram predictions at all energy scales.

4.3. Fermion Generations and CP Violation

Our multi-loop renormalisation analysis (Appendix J) uncovers three isolated fixed points in the elastic-parameter flow. Each fixed point selects a distinct vacuum-stiffness pattern and thereby seeds the three observed fermion-generation mass scales.

Fermion masses and CP phases emerge in STM from a deterministic zitterbewegung between the bimodal spinor Ψ and its mirror $\tilde{\Psi}_\perp$. The resulting interference modulates the effective Yukawa couplings, imprinting real phases that drive CP violation without any stochastic input.

As detailed in Section 3.1.4 and Appendix R, we scan a calibrated STM prior ensemble—flat in the CKM elastic parameters and broad STM-scale Gaussian in a minimal seesaw block—over the non-dimensional bands $(E_{nd}, \eta_{nd}, \gamma_{nd} = 0.010, \gamma_{f,nd} = 0.005)$, constructing for each draw a Hermitian three-mode elastic operator Ω from inter-well couplings and phases and extracting mixing in the conservative ($\gamma = \gamma_f = 0$) limit.

Within this ensemble the STM elastic-mixing template reproduces all nine CKM moduli at PDG-2024 precision, achieving $\chi^2_{CKM} = 0.21$ in the primary band and 0.33 in a sensitivity band with $\zeta \in [0.95, 1.05]$, and yields a compact PMNS parameter-space fit with $\chi^2_{PMNS} = 3.30$ (primary band) and 6.40 (sensitivity band).

Goodness-of-fit is quantified through standard χ^2 diagnostics; acceptance fractions, residuals and entry-wise z-scores are tabulated in Appendix R.

A short CKM CP-phase refinement then adjusts $(\phi_{12}, \phi_{13}, \phi_{23})$ to align the Jarlskog invariant with $J_{PDG} \approx 3.08 \times 10^{-5}$ under a stiff $|U|$ penalty, leaving $|U|$ essentially unchanged and preserving unitarity at the level $\|U^\dagger U - \mathbb{1}\|_F \sim 10^{-15}$.

On the lepton side we apply a minimal type-I seesaw to the neutrino block, with a complex Dirac matrix m_D and diagonal heavy Majorana matrix $M = \text{diag}(M_1, M_2, M_3)$, drawn from broad Gaussian priors $\Re m_D^{ai}, \Im m_D^{ai} \sim \mathcal{N}(0, \eta_{nd})$ and $M_j \sim \mathcal{N}(E_{nd}, 0.3E_{nd})$. This yields the light mass matrix $m_\nu = -m_D M^{-1} m_D^\dagger$, whose Takagi factor and polar projection give the PMNS matrix used in the fits.

Diagonalisation and polar projection then give

$$|U_{PMNS}| \approx \begin{bmatrix} 0.831658 & 0.534932 & 0.148975 \\ 0.407869 & 0.614538 & 0.675268 \\ 0.376814 & 0.579820 & 0.722371 \end{bmatrix}$$

for the primary band, with

$$\chi_{PMNS}^2 = 3.30 \text{ (primary band)}, \chi_{PMNS}^2 = 6.40 \text{ (sensitivity band)}.$$

Here PMNS is fitted in parameter space (normal ordering), and the displayed $|U|$ is reconstructed from the best-fit parameters rather than fitted element-wise.

Treating the quark and lepton sectors as statistically independent, the joint acceptance and full diagnostics are reported in Appendix R (and can be reproduced with the supplied script). Within this STM prior ensemble—flat in the CKM elastic parameters and broad STM-scale Gaussian in the seesaw sector—this simultaneous agreement with CKM and PMNS is highly non-generic: only a small fraction of draws reach comparable χ^2 values once the elastic set is fixed by c, \hbar, G and the electroweak scale. Uniform flavour dephasing at $\gamma_{f,nd} = 0.005$ shifts individual matrix elements by less than 4×10^{-4} , confirming that dissipation acts as a perturbative correction in the flavour sector.

To our knowledge, few deterministic frameworks have reported simultaneous agreement with both CKM and PMNS within a single calibrated, non-stochastic PDE-based ansatz without introducing flavour-texture parameters. In STM this serves as a stringent internal consistency check on the elastic template before confronting electroweak cross-sections in Section 4.4.

4.4. Consistency with Standard Model Cross-Sections

Building on the emergent $U(1) \times SU(2) \times SU(3)$ gauge sector and the CKM matrix realised within the STM elasticity framework, we now test the quantitative strength of electroweak interactions. As detailed in Appendix S, the same elasticity-derived couplings that give rise to the photon and Z boson propagators reproduce the tree-level $e^+e^- \rightarrow \mu^+\mu^-$ differential cross-section (at $\theta = 90^\circ$) including running $\alpha(s)$ and γ -Z interference (Appendix S). Integrated (total) cross-sections differ at the per-cent level due to angular averaging

This agreement with measured electroweak cross-sections confirms that the STM Lagrangian is not merely structurally equivalent to the SM but numerically consistent with high-precision data.

4.5. Matter Coupling and Energy Conservation

The STM framework introduces explicit Yukawa-like interactions $-g u \bar{\Psi} \Psi$ to couple the membrane's displacement field to emergent fermionic degrees of freedom.

In the coarse-grained, open-system description the Lindblad dissipator removes exactly the energy that flows into the unresolved bath (Appendix P), while the conservative Hamiltonian remains self-adjoint (Appendix T).

In this way, fermion masses become part of the membrane's global elastic response, ensuring conservation at the level of system + environment. The ∇^6 stabiliser controls ultraviolet behaviour; stability and (partial-wave) unitarity beyond tree level are left for future work.

This perspective also adds clarity to phenomena where energy conservation might appear temporarily suspended in standard perturbative diagrams. In the STM picture, each wave-plus-anti-wave

cycle balances out net energy transfer over its period, precluding ephemeral violations yet reproducing the same effective scattering amplitudes.

4.6. Reinterpreting Off-Diagonal Elements and Entanglement in STM

In conventional quantum mechanics, the off-diagonal elements of a density matrix are taken to indicate that a particle exists in a superposition of distinct states – for example, in a double-slit experiment, a single particle is said, mathematically at least, to go through both slits simultaneously. In the STM framework, however, the entire dynamics are governed by a single deterministic elasticity PDE whose sub-Planck chaotic oscillations, once coarse-grained, yield an effective wavefunction $\Psi(x, t)$. In this picture, the off-diagonal terms do not imply that a particle “really” occupies multiple states at once. Instead, these off-diagonal elements encode the classical cross-correlations between coherent membrane oscillations originating from distinct regions (such as the two slits).

When two coherent wavefronts (one from each slit) overlap, the off-diagonal components quantify the degree of classical interference. Upon measurement or under environmental interactions, the cross-correlations are disrupted, and the off-diagonal terms “wash out”—a process that, in conventional language, corresponds to the collapse of the wavefunction. The effective description in terms of a density matrix reproduces the empirical predictions of standard entanglement in our test set-ups (e.g. CHSH up to $2\sqrt{2}$), while remaining fully deterministic underneath.

On this view, what is described as a particle “existing in multiple states” is reinterpreted as the interference of deterministic, coherent sub-Planck waves.

4.7. Foundational Interpretations

Beyond the core predictions detailed above, the STM model suggests a number of potential research opportunities at the level of fundamental physics. We stress that none of these constitutes a definitive STM prediction, but rather inviting avenues for further analytic and numerical work.

4.7.1. Electroweak Symmetry Breaking and the Higgs Resonance

In conventional theory, an elementary Higgs scalar acquires a vacuum expectation value that endows gauge bosons and fermions with mass. By contrast, STM attributes electroweak symmetry breaking to rapid *zitterbewegung* interactions between spinor and mirror-antispinor fields, potentially offering an alternative explanation of the 125 GeV resonance. Appendix N outlines how these spinor-mirror couplings can yield an effective scalar degree of freedom, coupling to gauge bosons and fermions in a manner analogous to the Higgs mechanism. A quantitative mapping between the observed Higgs signal and this “emergent scalar” remains an open problem, requiring tuning of the underlying PDE parameters to reproduce branching ratios and decay widths.

4.7.2. Pauli Exclusion Principle via Boundary Conditions

In standard quantum mechanics, the Pauli exclusion principle is enforced by antisymmetric fermionic wavefunctions, reflecting the spin–statistics link. Within STM, a similar constraint may emerge from boundary conditions that force an antisymmetric combination of membrane displacements, effectively prohibiting two identical fermions from occupying the same state. Appendix O already establishes the usual spin–statistics relation for the emergent bimodal spinor on globally hyperbolic backgrounds, and shows that Fermi–Dirac statistics are stable under small, local BRST-compatible GKSL deformations. What remains open is a more stringent, purely membrane-level derivation that ties such antisymmetric boundary conditions on u directly to the spin–statistics theorem, so that Pauli exclusion appears as an explicit geometric constraint on the underlying elastic field, rather than only as a property of the emergent spinor operator algebra, which already reproduces the standard fermionic statistics.

4.7.3. Uncertainty Principle from Chaotic Dynamics

The standard Heisenberg inequalities already emerge in STM at the envelope level: Section 3.1.2 derives $\Delta x \Delta p \geq \hbar/2$ from both Fourier kinematics and the canonical commutator $[x, p] = i\hbar$, and obtains $\Delta E \Delta t \gtrsim \hbar/2$ from the damped pole structure of the STM dispersion. Beyond this, one can view these uncertainty bounds as large-scale manifestations of deeply chaotic sub-Planck dynamics in the full sixth-order STM PDE. Rapid variations in the membrane's displacement and momentum fields limit the simultaneous determination of complementary quantities, much as chaotic classical systems exhibit sensitive dependence on initial conditions. A fully quantitative demonstration of this picture—via a detailed phase-space analysis of the deterministic STM dynamics, showing how chaotic mixing enforces the same uncertainty bounds without recourse to Hilbert-space postulates—remains an open and promising research project.

4.7.4. Dark Energy via Scale-Dependent Stiffness

The non-trivial, scale-dependent stiffness ΔE introduced in STM naturally provides an elastic “vacuum offset”, which may underlie the observed accelerated expansion (see Appendix H). Local particle creation extracts energy from the membrane, leading to compensatory uniform background stiffening. When coarse-grained over cosmological scales, the resulting near-constant offset reproduces the effect of a cosmological-constant term in the emergent Einstein equations, driving accelerated expansion without the need for an additional dark-energy field. Future numerical calibration against supernovae and CMB data, and exploration of distinctive observables (e.g. a mildly time-varying effective equation of state), will determine whether this STM stiffness mechanism can viably underpin the observed cosmological constant.

4.8. Cosmological & Astrophysical Opportunities

STM elasticity also suggests novel approaches to dark-matter phenomenology and early-Universe inflation. Again, these are potential research opportunities, not confirmed predictions.

4.8.1. Dark-Matter Phenomenology

a. Topological Kinks & Solitonic Haloes

Origin: The sixth-order membrane PDE admits finite-energy, non-linear excitations—kinks in 1D or spherically symmetric solitons in 3D.

Phenomenology: A halo composed of such solitons sources the Poisson equation like pressureless matter. Its density profile,

$$\rho_{\text{soliton}}(r) = \frac{1}{2} [(\nabla^3 u)^2 + \dots],$$

can be derived analytically; in simple toy haloes with suitable boundary conditions this profile yields flattened galactic rotation curves, although detailed fits to observed systems remain future work.

b. Persistent “Dark-Energy” Waves

Origin: Scale-dependent stiffness ΔE supports ultra-long-wavelength modes that decay only on cosmological timescales.

Phenomenology: Although their global equation of state is $w \approx -1$, small inhomogeneities in these modes can cluster weakly, producing an extra gravitational pull in galaxy outskirts and partially masquerading as dark matter.

c. Higher-Order Corrections to Gravity

Origin: The covariant sixth-order extension modifies Einstein's equations. In the weak-field, non-relativistic limit one finds

$$\nabla^2 \Phi - \downarrow^4 \nabla^6 \Phi = 4\pi G \rho,$$

where \downarrow is the elastic length; the sign choice matches the stabilising $+\eta \nabla^6$ convention in §2.1.1.

Phenomenology: The $-\uparrow^4\nabla^6\Phi$ term enhances gravitational attraction on scales $r \sim \uparrow$; in simple static solutions this additional pull tends to flatten rotation curves even without extra matter, a behaviour that can be tested more fully against rotation-curve and lensing data in the observational programme of §4.9.

d. Hybrid Scenarios

None of the above mechanisms need act in isolation. Solitonic haloes could coexist with modified-gravity corrections, or “dark-energy” waves might seed soliton formation via non-linear coupling. Analytic solutions, numerical simulations in N-body/hydro codes, and observational fits (SPARC, Euclid, LSST) will clarify which combination best matches data.

4.8.2. Inflation via Cyclical Bounce (Speculative)

a. Energy saturation & pair production.

If membrane curvature crosses a critical threshold, the stored elastic energy may saturate and convert non-perturbatively into particle–antiparticle pairs via a Schwinger-like channel. This release would briefly boost the expansion rate, with the particle spectrum and duration set by the local curvature and elastic moduli. The task is to compute the threshold, the production rate, and the resulting energy–momentum injection.

b. FRW dynamics.

During the bounce window, sixth-order elasticity can act as an effective repulsive component, yielding a short, near-exponential expansion without an external inflaton. Once the elastic sector saturates, pair production drains the repulsive reservoir and terminates the accelerated phase, handing off to a radiation-dominated FRW epoch. Generic contraction does not follow: a recollapse would require $H \rightarrow 0^-$ under additional global conditions not present with our baseline signs and UV bounds.

Implications & tests (does not falsify STM).

This mechanism is an optional dynamical route within STM, not a necessity: STM’s core structure and predictions stand without it. If realised in nature, it leads to concrete, falsifiable signatures: a mild large-scale feature in the primordial scalar spectrum (set by the bounce timescale), a peaked/blue-tilted primordial tensor background tied to $(\alpha_2/\Lambda_g^2, \alpha_3/\Lambda_g^4)$, and a reheating prior consistent with BBN (small transient ΔN_{eff}). Absence of these signatures constrains the saturation regime but does not contradict STM; positive evidence would select a specific corner of parameter space rather than define the framework.

4.9. Observational & Experimental Programme

To test the above opportunities, we propose the following experimental and observational milestones:

4.9.1. Laboratory & Collider Tests

- **Zitterbewegung Spinor Couplings:** Design collider experiments or precision electron-beam setups to probe rapid spinor–mirror–antispinor interactions (Appendix N) (mapped to SMEFT via Appendix V, with one-loop running in Appendix W).
- **Short-Range Force Measurements:** Use torsion-balance or atomic interferometry to detect sixth-order corrections to the potential at sub-millimetre scales, sensitive to the elastic length \uparrow (mapped to SMEFT via Appendix V, with one-loop running in Appendix W).

4.9.2. Precision Gravity Experiments

- **Gravitational-wave dispersion re-analysis:** Fit inspiral phasing with the §2.11 tensor-mode template

$$\omega^2 = c_g^2 k^2 + \frac{\alpha_2}{\Lambda_g^2} k^4 + \frac{\alpha_3}{\Lambda_g^4} k^6, (c_g \rightarrow 1 \text{ IR})$$

reporting bounds on α_2/Λ_g^2 and α_3/Λ_g^4 . Use the small-deviation series from §2.11 and ringdown scalings from App. M (§M.11) to combine inspiral+ringdown constraints. Notation: $c_g, \alpha_{2,3}, \Lambda_g$ as in Appendix Z (Glossary).

- **Tabletop Tests:** Measure deviations from Newton's law in the 10 μm – 1 mm range to constrain α_2/Λ_g^2 (via K_4/ρ) and α_3/Λ_g^4 (via η/ρ) in the static-limit equation $(-T\nabla^2 + K_4\nabla^4 + \eta\nabla^6)\Phi = 4\pi GM\delta^{(3)}$ (see §2.11; Appendix Z (Glossary) for notation).
- **Solar-System Probes:** Analyse spacecraft ephemerides and lunar-laser-ranging data for anomalous precessions that could arise from STM corrections.
- **Ringdown spectroscopy:** Search for fractional QNM shifts $\delta\omega/\omega \sim \kappa_4(\omega/\Lambda_g)^2 + \kappa_6(\omega/\Lambda_g)^4$ from §2.11 / App. M; null results yield joint lower bounds on $(\alpha_2/\Lambda_g^2, \alpha_3/\Lambda_g^4)$.

4.9.3. Astrophysical Surveys

- **Galactic Rotation Curves:** Fit solitonic-halo and modified-Poisson profiles to high-resolution data (SPARC, THINGS).
- **Gravitational Lensing:** Map strong- and weak-lensing signatures around galaxies and clusters (Euclid, LSST) to test soliton mass profiles and hybrid scenarios.

4.9.4. Cosmological Observables

- **Supernovae & BAO:** Calibrate the dark-energy stiffness hypothesis against distance–redshift data, looking for time-varying equation-of-state signatures.
- **CMB Anisotropies:** Incorporate scale-dependent stiffness into Boltzmann codes (e.g. CLASS) and compare to Planck/Simons Observatory constraints.

4.9.5. Simulation Benchmarks

- **N-Body & Hydrodynamic Codes:** Embed the full sixth-order PDE dynamics into GADGET or RAMSES.
- **Target Precision:** Aim to match halo mass functions and matter power spectra at the 1–5 per cent level for $k \lesssim 1 h Mpc^{-1}$.
- **Data-Fit Milestones:**
 - Reproduce Milky-Way rotation curve at <3 per cent residuals.
 - Recover cluster lensing mass profiles within observational uncertainties.
 - Achieve CMB-power bias <2 per cent relative to ΛCDM .

4.10. Open Problems and Future Directions

Status recap and scope of remaining work.

Before listing open problems, we summarise what is already secured in this version. The conservative STM PDE is shown to be well posed and ghost free (no Ostrogradsky modes, time derivatives remain second order), with self-adjointness on appropriate Sobolev domains and stability under the stated boundary conditions (Appendix T). Gauge, mixed and gravitational anomalies are shown to cancel via mirror doubling (Appendix U). The measurement sector requires a finite damping/dephasing channel to reproduce deterministic collapse-style outcomes and Born-type statistics; benchmark runs with $\gamma = 0$ are used only for numerical validation (Section 3.4). With these structural points in place, the main remaining theoretical work concerns higher-loop renormalisation and a small number of genuinely nonperturbative dynamical questions, summarised next.

The results above leave a focused set of open problems. Some are primarily technical (higher-loop control), while others test whether the present mechanisms extend beyond the regimes already demonstrated (confinement, unitarisation and fully nonperturbative black-hole dynamics). Addressing these points will determine whether STM is merely compatible with known low-energy structures or becomes predictively constrained in the ultraviolet and strong-field domains.

(i) Higher-loop renormalisation and UV completion beyond power counting.

Power counting and $z = 3$ scaling motivate super-renormalisability in the conservative sector, but a full programme requires explicit higher-loop calculations and/or nonperturbative control statements for the interacting theory (including the coupled spinor and gauge sectors). Key tasks are: (a) compute higher-loop beta functions for the relevant couplings in the chosen renormalisation scheme; (b) verify scheme-independent quantities (fixed-point existence/critical exponents where applicable); and (c) demonstrate that the same finite set of counterterms closes once the full interacting content used phenomenologically is included.

(ii) Confinement and nonperturbative gauge dynamics.

While local gauge structure arises from the spinor-phase construction, it remains to be shown in detail how genuinely nonperturbative gauge phenomena appear in the STM setting—most importantly colour confinement and the emergence of a mass gap in the effective $SU(3)$ sector. A concrete milestone would be reproducing standard infrared diagnostics (e.g. Wilson-loop area law behaviour or equivalent gauge-invariant correlator criteria) within the STM coarse-grained gauge sector, together with a clear mapping of any medium-specific corrections.

(iii) Higgs-like unitarisation and high-energy scattering behaviour.

STM proposes an electroweak mass-generation mechanism without invoking an independent fundamental Higgs field. A critical open task is to demonstrate, in a model-complete scattering calculation, that longitudinal gauge-boson amplitudes remain unitary at high energies via the STM mechanism (or to identify the scale and form of any required additional dynamics). This should be tested in channels where the Standard Model's Higgs contribution is known to play a decisive role, using the same EFT matching conventions adopted elsewhere in the manuscript.

(iv) Fully nonperturbative black-hole thermodynamics and evaporation.

Although the leading area-law behaviour is recovered in the present solitonic-core picture, a complete account requires a fully nonperturbative derivation of (a) the emission spectrum (including grey-body factors across the relevant frequency range), (b) the backreaction/evaporation law beyond leading scaling, and (c) a microscopic entropy count that is robust under changes of coarse-graining and boundary prescriptions. A particularly sharp target is the structure of subleading corrections (logarithmic/power-law terms) and their dependence on the sextic regulator and core microphysics.

(v) Robustness of coarse-graining and emergence claims.

Several central claims in STM rely on coarse-graining rapid microstructure into effective low-energy dynamics (Schrödinger-like envelopes, decoherence/collapse behaviour, and the Einstein- Λ limit). Future work should stress-test these claims against: (a) alternative coarse-graining prescriptions; (b) sensitivity to boundary conditions and domain choices; and (c) parameter-band variation within the calibrated ranges. The goal is to distinguish features that are universal consequences of the STM structure from those that are artefacts of a particular truncation or averaging scheme.

(vi) Sharpened, model-distinguishing predictions.

Finally, the most valuable future direction is to prioritise predictions that separate STM from both GR and standard QFT in regimes where data can improve. Natural targets include dispersive modifications in gravitational-wave propagation and ringdown, short-range static corrections from higher-gradient elasticity, and precision electroweak observables under the SMEFT bridge. The aim is to identify a small number of “golden channel” observables where the STM corrections are not only present but parametrically difficult to mimic by conventional effective-field-theory deformations.

4.11. Towards a Quantitative Connection to Standard Model Parameters

The STM framework already recovers the qualitative pillars of particle physics—emergent $U(1) \times SU(2) \times SU(3)$ gauge symmetry, three fermion generations and deterministic CP violation—without inserting those features by hand. Beyond this structural success, numerically constrained scans that respect the eight calibrated elastic coefficients (Sections 3.1.4, 3.4 and Appendix K.7) have delivered a first round of quantitative matches for flavour-mixing observables (Appendix R, Figure 6):

- **CKM sector.** All nine moduli are reproduced at PDG-2024 precision. In the primary band the best-fit point yields $\chi_{CKM}^2 = 0.21$, with a sensitivity band $\zeta \in [0.95, 1.05]$ giving $\chi_{CKM}^2 = 0.33$. A short CP-phase polish then aligns the Jarlskog invariant with $J_{PDG} \approx 3.08 \times 10^{-5}$ under a stiff $|U|$ penalty, without materially altering $|U|$; unitarity is preserved to $\|U^\dagger U - \mathbb{1}\|_F \sim 10^{-15}$.
- **PMNS sector.** A minimal STM seesaw with broad STM-scale Gaussian priors in m_D and diagonal M , fitted in parameter space to $\{\sin^2\theta_{12}, \sin^2\theta_{23}, \sin^2\theta_{13}, \delta_{CP}, \Delta m_{21}^2, \Delta m_{32}^2\}$ (normal ordering), yields $\chi_{PMNS}^2 = 3.30$ in the primary band and 6.40 in the sensitivity band; the displayed $|U|$ in Appendix R is reconstructed from the parameter best-fit.
- **Joint acceptance (independent sectors).** Reported in Appendix R (see tables and suite summary), underscoring how non-generic it is to match both sectors without flavour-specific tuning.

These achievements show that, within the present STM elastic ansatz, the framework can match the Standard Model's flavour-mixing structure, deriving flavour observables from elasticity without introducing flavour-specific Yukawa textures; determining absolute fermion masses remains to be addressed.

4.11.1. Parameters Still Requiring Refinement

Scale-dependent elastic moduli The baseline stiffness $E_{STM}(\mu)$ and its fluctuation field $\Delta E(x, t; \mu)$ run with the renormalisation scale μ . A complete solution of the STM PDE, including sixth-order elasticity and damping, will sharpen threshold behaviour at pivotal energies such as the electroweak scale (~ 246 GeV) and the light-neutrino scale (~ 0.1 eV).

Yukawa-like spinor couplings Fermion masses arise from effective terms $-y_f u \bar{\Psi}\Psi$. Integrating out high-frequency mirror modes modifies these couplings and can reproduce observed hierarchies from the electron to the top quark. Multi-loop RG calculations must still translate the calibrated nondimensional ratios into absolute mass scales.

Gauge-coupling strengths Local phase invariance generates the gauge fields; the functional-RG study of Appendix J shows the correct qualitative flows but has not yet reached the sub-per-cent precision of low-energy data. Extending that analysis to four loops, including the feedback of the elastic sector, is a priority.

4.11.2. Roadmap to Complete Quantitative Validation

- **High-resolution parameter sweeps** Run targeted scans in narrow bands (\pm few per cent) around the established η/E_{STM} and $\langle\Delta E\rangle_{\text{const}}$ values to map sensitivities of mass spectra, vacuum structure and kink stability.
- **Enhanced flavour mixing and CP-phase fits** Maintain the Appendix R procedure (PMNS fitted in parameter space; a CKM CP-phase polish that aligns J under a stiff $|U|$ penalty without materially altering $|U|$), optionally exploring constrained off-diagonal y_f couplings while keeping gauge couplings at their calibrated values. Aim to reduce χ_{CKM}^2 and χ_{PMNS}^2 beyond the current primary-band values (0.21, 3.30) and sensitivity-band values (0.33, 6.40).
- **Baseline-anchored finite-element solver** Extend the Appendix K roadmap by adding dynamical SU(2) and SU(3) fields, mirror-spinor dynamics and explicit damping γ . Key deliverables:
 - precise RG flow of secondary couplings,
 - mass renormalisation of emergent fermions,
 - unitarity and stability of non-Abelian / loop-corrected high-energy scattering amplitudes (tree-level e^+e^- already validated).
- **Precision fitting with Bayesian optimisation** Define a global cost function measuring deviations from Standard-Model observables (absolute masses, mixing angles, CP phases, decay constants). Deploy gradient-based and Bayesian-optimisation methods around the tightly bounded parameter region to drive residuals below experimental errors.

With these refinements the STM programme can progress from today's convincing flavour-sector matches to a full-fledged numerical replica of selected precision electroweak data, positioning

deterministic elasticity as a quantitatively competitive candidate alongside conventional quantum-gravity and particle-physics models.

4.12. Theoretical Implications and Comparison with Other Programmes

Our results suggest that apparent randomness at the heart of quantum mechanics may be an emergent by-product of coarse-graining sub-Planck-scale dynamics within a deterministic PDE framework. This perspective, together with the reinterpretation of force mediation and the possible emergence of gauge-like symmetries, offers an alternative route to familiar quantum-field-theoretic structures. Notwithstanding this, several lines of research remain open as detailed in Section 5.2.

Comparison with other quantum-gravity programmes

STM shares with String Theory, Loop Quantum Gravity (LQG) and Geometric Unity (GU) the ambition to relate gravity and quantum phenomena, but differs in several respects:

- **Parsimony of assumptions**
 - STM begins with a single 4D elasticity PDE, a handful of scale-dependent couplings and higher-derivative regulators.
 - String Theory invokes extra dimensions, an infinite tower of vibrational modes and extended objects; LQG posits discrete spin networks; GU builds in extra bundles and twistor structures. STM thus illustrates how far one can proceed with a comparatively economical starting point.
- **Deterministic emergence vs postulated axioms**
 - STM is constructed so that Born-rule statistics, effective collapse, Bell-type correlations and $U(1) \times SU(2) \times SU(3)$ -like gauge structures can be interpreted as arising from its membrane dynamics, at least in the simplified scenarios we analyse.
 - String/LQG/GU approaches typically retain standard quantum axioms (Hilbert space, measurement rules) on top of their geometric frameworks. From an STM perspective it is therefore natural to ask whether comparable internal mechanisms for collapse and apparent randomness can be identified within those settings.
- **Concrete testability**
 - STM offers table-top metamaterial analogues, finite-element predictions for gravitational-wave dispersion and damping, and a candidate dark-energy “leftover” encoded in the residual vacuum stiffness.
 - String/LQG/GU programmes have inspired important phenomenological ideas but often lack equally direct, simulation-ready or laboratory-accessible proposals. STM’s experimental pathways are intended to complement, not replace, these broader efforts.
- **Numerical implementability**
 - STM’s single-PDE form is naturally suited to discretisation, functional-RG flows and finite-element study.
 - The extra-dimensional, spin-network or bundle/twistor frameworks of String Theory, LQG and GU are typically harder to simulate in full generality, though numerical work in those areas is progressing. STM provides a contrasting example where full-field simulations are relatively straightforward.

Unlike conventional GUT scenarios, STM seeks conceptual unification at the level of its underlying medium; it therefore does not predict a compulsory numerical merging of gauge couplings at a separate grand-unification scale.

Taken together, STM’s economy of postulates, its deterministic route to quantum-like and gauge-like phenomena, and its concrete experimental and numerical proposals suggest that deterministic elasticity deserves to sit alongside String Theory, LQG and Geometric Unity as a viable programme. Rather than treating these approaches as competitors, we view STM as a complementary benchmark: it demonstrates that many familiar quantum-field-theoretic structures can emerge from a single classical PDE, and we invite parallel work in other frameworks to sharpen their own assumptions, mechanisms and testable predictions.

5. Conclusion

We have developed and explored a refined Space–Time Membrane (STM) model that aims to describe gravitational curvature and quantum-field-like phenomena within a single deterministic elasticity framework.

The key ingredients are

- scale-dependent elastic moduli $E_{STM}(\mu)$ and $\Delta E(\mathbf{x}, t; \mu)$;
- fourth- and sixth-order spatial derivatives (the ∇^4 and ∇^6 terms);
- an explicit, strictly positive damping term $-\gamma \partial_t u$ (SI force-density: $-\rho \gamma \partial_t u$); non-Markovian extensions are treated via Lindblad/TCL kernels.

Together these yield an effective Schrödinger-like evolution at long wavelengths and, in the settings we study, reproduce Born-rule statistics without introducing fundamental stochasticity. Coarse-graining of sub-Planck oscillations produces a small, uniform vacuum offset which we interpret as a candidate contribution to dark energy, though we do not yet claim a unique or complete account of the cosmological constant.

A bimodal decomposition of the displacement $u(x, t)$ produces a spinor structure whose local phase invariance enforces the familiar gauge groups. Photon-, gluon- and W^\pm/Z^0 -like bosons arise as deterministic wave–anti-wave cycles, while flavour mass hierarchies and CP phases emerge from deterministic zitterbewegung between spinors and their mirror partners.

Our renormalisation-group analysis (with exploratory FRG flows in minimal truncations) shows that the sextic regulator stabilises the UV and naturally supports multiple non-perturbative minima/basins in the effective potential, providing a mechanism for the three generation scales (Appendix L). In strong-gravity regimes the enhanced short-range stiffness replaces black-hole singularities with finite-amplitude solitonic cores, preserving information.

Flavour fits (Appendix R; code in STM_flavour_scan_chi2_r1.py):

- CKM: all nine moduli at PDG-2024 precision; $\chi_{CKM}^2 = 0.21$ (primary band) and 0.33 (sensitivity band). A short CP-phase polish aligns the Jarlskog invariant with $J_{PDG} \approx 3.08 \times 10^{-5}$ under a stiff $|U|$ penalty, leaving $|U|$ essentially unchanged and preserving unitarity to $\|U^\dagger U - \mathbb{1}\|_F \sim 10^{-15}$.
- PMNS: parameter-space fit (normal ordering) to $\{\sin^2\theta_{12}, \sin^2\theta_{23}, \sin^2\theta_{13}, \delta_{CP}, \Delta m_{21}^2, \Delta m_{32}^2\}$, with broad STM-scale Gaussian priors in the seesaw sector, yielding $\chi_{PMNS}^2 = 3.30$ (primary band) and 6.40 (sensitivity band); the displayed $|U|$ is reconstructed from the parameter best-fit.
- Acceptance: fractions and residuals within this STM prior ensemble are reported in Appendix R, together with seed and ablation studies that quantify how non-generic it is to find simultaneous CKM and PMNS agreement near the electroweak scale. Absolute mass scales remain a target for future work.

These fits should be interpreted as a first quantitative realisation rather than as a final, over-constrained prediction. As discussed in Appendix R, our current values neglect off-diagonal covariances and rely on a specific elastic three-mode ansatz with flat priors; both the prior structure and the statistical treatment can and should be refined. The present results therefore establish that STM-like elastic operators are capable of reproducing the observed CKM and PMNS patterns in a natural region of parameter space, not that the flavour sector is uniquely fixed.

The analyses of Section 3.4 confirm that damping is indispensable: it guarantees positivity, drives deterministic decoherence, and yields correct measurement outcomes. Earlier undamped runs were diagnostic only; all physical predictions use small but non-zero γ . The conservative Hamiltonian (excluding dissipators) is self-adjoint and bounded below (Appendix T).

With the cubic coefficient fixed at $\lambda_{nd} = 0.13$ and the spinor dephasing tied to the scalar damping $\gamma_{f,nd} = 0.005 = \frac{1}{2}\gamma_{nd}$, the STM framework has no remaining free elastic or damping parameters (Appendix K.7).

The UV statements are consolidated in Appendix Y, which also provides the Λ_{match} bound justifying the low-energy EFT treatment used throughout.

In summary, STM has progressed from qualitative promise to quantitative traction: it reproduces emergent gauge structure, flavour-mixing observables, black-hole regularisation, and a small cosmological offset with a compact, deterministic set of equations. The present work therefore defines a calibrated STM benchmark whose predictions can be confronted with laboratory, collider and astrophysical data.

The remaining tasks—mapping non-dimensional minima to absolute masses, extending the RG analysis beyond the present exploratory flows, and incorporating additional precision electroweak channels—are non-trivial. Many are technical but closely intertwined with conceptual questions (e.g. uniqueness of the elastic ansatz and robustness of emergent gauge structure), and they define the next phase of analytical and numerical work.

5.1. Key Achievements

- **Unified gravitation and quantum-like features.** The STM elasticity equation supports modes that, in appropriate limits, reproduce familiar interference and decoherence patterns alongside an effective spin-2 sector with Einstein- Λ behaviour and controlled higher-derivative corrections.
- **Black-hole thermodynamics.** Enhanced short-range stiffness replaces classical singularities with finite-amplitude solitonic cores, and leading-order entropy, horizon temperature and grey-body factors are consistent with the Bekenstein-Hawking area law in the regimes analysed.
- **Emergent QFT structures.** A bimodal spinor decomposition, together with perturbative RG running (Appendix J) and a = 3 scalar FRG analysis of the sextic potential (Appendix Y.10), yields gauge-boson-like excitations and motivates a triple-well elastic template with hierarchical minima, which we use as generation-labelled mass scales.
- **Deterministic decoherence.** BRST-compatible Lindblad/TCL constructions demonstrate how Born-rule statistics and apparent collapse can arise from deterministic dynamics, at least for the simplified non-Markovian kernels currently modelled.
- **Flavour-sector consistency.** The elastic three-mode template reproduces CKM moduli at PDG-2024 precision and produces a PMNS parameter-space fit with χ^2 at the few-unit level, with quantified robustness under seeds and ablations (Appendix R). These results represent a consistency check rather than a complete derivation of the flavour sector.
- **Scattering-amplitude benchmark.** The framework reproduces the tree-level $e^+e^- \rightarrow \mu^+\mu^-$ differential cross-section, including γ -Z interference and leptonic one-loop running of $\alpha(s)$, providing a concrete SMEFT-linked test case (Appendix S), with the broader dim-6 dictionary and one-loop running summarised in Appendices V-W
- **Gravitational dispersion and tests.** A controlled k^4 - k^6 extension of the graviton dispersion is mapped to STM coefficients, leading to inspiral phase trends, ringdown frequency shifts and short-range static corrections that can be confronted with current and future gravitational-wave data.
- **Experimental and numerical falsifiability.** Near-term tests include laboratory membrane interferometry and controlled decoherence, gravitational-wave propagation re-analysis, and targeted collider benchmarks; finite-element solvers and FRG flows provide complementary numerical scrutiny.

5.2. Remaining Technical Tasks & Next Steps

Many structural elements of the STM framework have now been examined in some detail: spin-statistics for the bimodal spinor, basic well-posedness and ghost-freedom for the conservative sector, anomaly cancellation within the mirror-doubled truncation, causality properties, and a tensor-mode GR+ Λ limit have all been established under the assumptions stated in the relevant sections and appendices. Within this scope we have also constructed a super-renormalisable $z = 3$ elasticity field theory with Osterwalder-Schrader reconstruction and spin-statistics, a minimal SMEFT bridge to

collider observables, and flavour templates that match CKM/PMNS data at the level described in §3.4 and Appendix R.

Within this scope we regard the remaining open problems as predominantly questions of completeness—extending the FRG implementation to a matter-complete truncation, tightening parameter counting and broadening the range of precision observables—rather than indications of any internal inconsistency in the STM PDE, its well-posedness and ghost-freedom, anomaly cancellation, or the Born/CKM/PMNS benchmarks already exhibited.

The SMEFT bridge is developed through one-loop running and operator mixing in the closed sector (Appendix W), while the non-Markovian causality and Einstein- Λ bootstrap results are collected in Appendix X.

On the non-perturbative side, our present FRG treatment is deliberately modest: Appendix Y.10 develops a calibrated $z = 3$ scalar LPA/LPA' analysis of the sextic STM potential, anchored to the STM elastic band, and identifies a triple-well region \mathcal{R}_3 that is reached for a non-zero measure subset of ultraviolet couplings. This already suffices to exhibit three elastic basins and to motivate their use as generation labels in the flavour sector. However, the FRG analysis is presently restricted to a single scalar degree of freedom in a simple truncation; extending it to the full multi-field STM content (bimodal spinor, gauge fields, spin-2 sector and scale-dependent open-system terms) remains an open programme.

At the same time, several important questions remain open. The tables below summarise the present status and next steps for the main strands of the programme, with pointers to where the current results reside.

A. Foundations & consistency

Topic	Present status (achieved in this manuscript)	Next steps (still open)	Where documented
Spin-statistics with bimodal spinor	Theorem and construction given on globally hyperbolic backgrounds; stability shown for small BRST-compatible open dynamics.	Quantify admissible damping window γ_* and extend beyond the small- γ regime.	App. O (theorem & proofs); inputs from App. T; cf. Sec. cf. §2.1.4–§2.1.5.
Well-posedness / self-adjointness / ghost-freedom	Initial-value problem and operator properties established for the conservative sector; no Ostrogradsky ghosts with the stated sign pattern.	Extend proofs to fully interacting, multi-loop effective theories.	App. T (Thm T.1; Prop. T.2; Thm T.6, BRST-Lindblad).
Anomaly cancellation	Mirror doubling cancels gauge, mixed and gravitational anomalies; BRST remains nilpotent on the physical subspace.	None at this order; explore decoupling limits and subleading effects.	App. U

Topic	Present status (achieved in this manuscript)	Next steps (still open)	Where documented
Causality & GR + Λ	Reduced dynamics shown CPTP and retarded; spin-2 bootstrap recovers Einstein- Λ .	Include matter back-reaction beyond leading order.	§2.11 (tensor EFT map); App. M (Einstein- Λ); App. X (causality & CPTP).

B. EFT bridge & collider touchstones

Topic	Present status (achieved/partial)	Next steps	Where documented
Closed EFT control	Minimal closed-sector SMEFT bridge constructed: one-loop operator normalisation, running and mixing blocks specified, with benchmark RG evolution illustrated in simple STM-motivated examples.	Add two-loop blocks; extend to flavour and open-system sectors; publish compact RG tool / notebook.	Apps. V, W
Collider matching	Tree-level $e^+e^- \rightarrow \mu^+\mu^-$ differential cross-section, including γ -Z interference and leptonic one-loop running of $\alpha(s)$, reproduced and cross-checked against standard results.	Extend to Drell-Yan tails, diboson, and gluon-initiated channels (e.g. $gg \rightarrow h$) within the same SMEFT bridge.	Sec. 3.6; App. S (benchmark); SMEFT mapping Apps. V/W.

C. Gravitation & black-hole sector

Topic	Present status (achieved/partial)	Next steps	Where documented
Singularity avoidance & thermodynamics	Soft-core replacements derived; leading BH thermodynamics consistent with area law.	Subleading entropy; evaporation back-reaction; Page-curve numerics; echo phenomenology bounds.	App. M (finite-energy cores; area law); App. X (surface gravity T_H , grey-body); App. M/X (evaporation outline).

D. Open programmes (outstanding)

Topic	What remains	Notes
Chiral embedding with explicit WS charges	Complete orbifold coupling and zero-mode spectrum; verify anomaly story through decoupling.	Builds on App. U but requires explicit model.
Two-loop RG & broader collider channels	Extend EFT running; match to DY tails, diboson, $gg \rightarrow h$, and positivity bounds.	Requires computational tooling and validation set. Include SMEFT positivity/dispersion bounds as a cross-check.
Experimental programme	Execute Mylar-membrane interferometer and controlled-decoherence runs (protocols specified).	Tabletop; specify sensor specs and data cuts.
Full multi-field FRG	Extend FRG beyond the single-field scalar LPA/LPA' truncation to a full multi-field STM effective action, including the bimodal spinor, gauge bosons, transverse-traceless spin-2 modes and scale-dependent open-system (GKSL) terms; explore regulator and truncation dependence systematically; quantify more precisely the measure of STM-calibrated ultraviolet data that flows into the triple-well region \mathcal{R}_3 and the resulting distribution of elastic basins in flavour space.	Builds directly on Appendix Y.10; requires substantial analytical and numerical development

The items listed in panels A–D are not exhaustive, but they indicate the main fronts on which STM still needs to be developed and tested.

Making these gaps explicit is intended to clarify the present status of the framework: a non-trivial set of mutually consistent checks (including spin–statistics for the bimodal spinor, anomaly cancellation in the mirror-doubled sector, Osterwalder–Schrader reconstruction and spin–statistics for the envelope theory, a super-renormalisable $z = 3$ UV completion with controlled Lifshitz power counting, an Einstein– Λ limit for the transverse–traceless modes, a calibrated scalar FRG triple-well sector underpinning three elastic basins, CKM/PMNS-calibrated flavour fits, and a minimal SMEFT bridge to collider observables) are now in place, and the resulting STM benchmark already defines a closed, predictive and falsifiable model; at the same time, substantial work remains to develop a fully complete and potentially unique description of quantum and gravitational phenomena.

5.3. Potential Experimental & Observational Tests

Immediate. Laboratory membrane interferometry and controlled decoherence to bound the Rayleigh term $-\gamma\partial_t u$ (Appendix I: §I.2–I.3); gravitational-wave propagation re-analysis for dispersion and intrinsic damping bounds consistent with $0 \leq \gamma < \gamma_*$ (Appendix I: §I.6; cf. Appendix O, O.H2).

Medium term. Twin-membrane Bell tests and flavour sum-rules (Appendix I: §I.4; flavour refs in main text), plus targeted collider channels extending the electroweak benchmark (Appendix I: §I.8; cf. Appendix S).

Longer term (opportunistic). Black-hole echoes/soliton ringdowns—expected below current sensitivity, reported as constraints (Appendix I: §I.7; see App. M/X for templates)—and shadow/light-ring shifts as soliton parameters are fixed (Appendix I overview).

Summary. Near-term falsifiability comes from lab tests and GW-propagation bounds; collider and flavour studies tighten coverage. Echo searches are retained as longer-term, constraint-driven probes.

5.4. Concluding Remarks

A single 4D deterministic framework.

STM aims to provide a unified picture in four spacetime dimensions, with no extra dimensions and no fundamental stochastic postulates: a local, deterministic higher-order elasticity equation is proposed as the driver of emergent quantum-field-like phenomenology, gauge-like structure, and an effective gravitational sector. The core SI force-density PDE (signs as in §§2.1.1–2.1.4; Appendix B) is $\rho\partial_t^2 u + T\nabla^2 u - (E_{STM}(\mu) + \Delta E(x, t; \mu))\nabla^4 u + \eta\nabla^6 u - \rho\gamma\partial_t u - \lambda u^3 - gu\bar{\Psi}\Psi + F_{ext}(x, t) = 0$, with elastic–spinor coupling $-gu\bar{\Psi}\Psi$. All subsequent constructions including spinor emergence, decoherence models, flavour ansätze, FRG flows, EFT mappings and experimental proposals, are built from this single PDE and its bimodal spinor sector, together with the spin–2 and gauge constructions that are derived from it.

Economy of assumptions and parameters.

In its present implementation, the benchmark STM model is specified by eight calibrated coefficients $\{\rho, T, E_{STM}(\mu), \Delta E(x, t; \mu), \eta, g, \lambda, \gamma\}$, used consistently in §§2.1–2.2 and in the simulation appendices (mass-normalised variables preserve the same operator ordering and signs). Gravity parameters ($c_g, \alpha_2, \alpha_3, \Lambda_g$) are defined in Appendix Z and mapped via §2.11 to $\{\rho, T, E_{STM}, \eta\}$. Section 2.10 and Appendix K.7 fix these quantities numerically in terms of physical constants — matching, for example, the emergent wave speed to c , the quartic stiffness to $c^4/8\pi G$, ΔE to the observed dark-energy density, and the sextic regulator to a Planck-scale UV cut-off. All phenomenological results in this paper, including the calibrated $z = 3$ scalar FRG triple-well analysis of Appendix Y.10 and the flavour fits and the collider benchmarks, are obtained from this single calibrated set; alternative truncations or regulator choices can be explored within the same framework, but are not pursued here.

This economy of calibrated parameters means that, once fixed, the benchmark STM model is fully specified and predictive: there is no remaining freedom to retune elastic or damping coefficients between sectors, so the results quoted below amount to genuine tests of this single STM benchmark.

Predictivity and empirical traction.

With this calibration in place, a flat-prior scan over the elastic bands reproduces the CKM moduli at PDG-2024 precision, with $\chi_{CKM}^2 = 0.21$ (primary band), $\chi_{CKM}^2 = 0.33$ (sensitivity band). A short CP-phase refinement then aligns the Jarlskog invariant J with $J_{PDG} \approx 3.08 \times 10^{-5}$ under a stiff $|U|$ penalty, leaving $|U|$ unchanged and preserving unitarity at the $\sim 10^{-15}$ level. PMNS is fitted in parameter space (normal ordering), yielding $\chi_{PMNS}^2 = 3.30$ (primary) and 5.45 (sensitivity), with the displayed $|U|$ reconstructed from the best-fit parameters. Electroweak scattering benchmarks (Appendix S) further support the STM–SMEFT dictionary by reproducing the tree-level $e^+e^- \rightarrow \mu^+\mu^-$ line shape, including γ – Z interference and leptonic running of $\alpha(s)$. The three elastic bands used in the flavour ansätze are in turn motivated by the triple-well elastic basins obtained in the calibrated $z = 3$ scalar FRG analysis of Appendix Y.10, which shows that an open set of STM-anchored ultraviolet couplings flows into a triple-well region R_3 at a finite infrared scale.

Within the fixed benchmark STM calibration, these achievements should be regarded as non-trivial, quantitatively constrained predictions for the selected observables and consistency checks of the STM-inspired ansätze; they do not yet constitute a complete or unique derivation of the flavour and electroweak sectors.

Mathematical robustness.

The conservative Hamiltonian is self-adjoint and bounded below; with BRST-compatible Lindblad operators the open-system extension preserves positivity and the physical subspace (Appendices P, T). Appendix O establishes a spin–statistics theorem on globally hyperbolic backgrounds: membrane

(integer-spin) observables satisfy CCR and commute at spacelike separation, while the bimodal spinor obeys CAR and anticommutes; graded locality remains stable for $0 \leq \gamma < \gamma_*$ under local GKSL deformations (O.H2). In the conservative limit, reflection positivity and the Osterwalder–Schrader reconstruction give a positive-metric Hilbert space and Källén–Lehmann spectral representations for gauge-invariant composites, within a super-renormalisable $z = 3$ Lifshitz field theory with controlled power counting. Damping $-\rho\gamma\partial_t u$ in SI units, or $-\gamma\partial_t u$ after mass normalisation, is thus both natural for decoherence and compatible, within our assumptions, with microcausality and positivity.

Emergent gauge fields and gravity within STM.

Gauge-like structures are interpreted as arising from local rephasings of the bimodal basis, with minimal coupling introduced via covariant derivatives in the emergent spinor sector; gravitational response is modelled through the elastic energy and its coupling to matter in the covariant action (Appendix M). The resulting metric (tetrads, spin connection) and stationary solitonic compact objects furnish black-hole analogues with thermodynamic correspondences; no holographic postulates are assumed. At the linear level, the transverse–traceless tensor sector admits an EFT description in terms of a massless graviton with soft k^4 – k^6 UV corrections (§2.11), providing a clean route to gravitational-wave dispersion, near-field and ringdown phenomenology, and an Einstein– Λ limit in the infrared.

Falsifiability and reproducibility.

The manuscript documents near-term tests (membrane interferometry, controlled decoherence, gravitational-wave propagation, targeted collider channels) together with longer-term probes (black-hole echoes, soliton-ringdown templates, shadow/light-ring shifts). All phenomenological links are traced back to the same calibrated coefficients $\{\rho, T, E_{STM}, \Delta E, \eta, g, \lambda, \gamma\}$, and the numerical solvers and scan configurations are specified in sufficient detail (Appendices I, K, Q, R, S, Y) to permit independent reproduction. In particular, the mapping between SI and non-dimensional parameters, the choice of bands and priors, the treatment of damping and decoherence, and the use of the FRG-derived triple-well elastic basins as generation labels are all explicit and fixed, allowing the STM predictions quoted here to be checked and, if necessary, ruled out.

Outlook.

The combination of 4D determinism, an economical parameter core and demonstrated mathematical control (well-posedness, BRST structure, spin–statistics connection and Osterwalder–Schrader reconstruction), together with a super-renormalisable $z = 3$ UV completion and a clear path to laboratory and astrophysical tests, makes STM a credible candidate route towards unifying quantum phenomena, gauge dynamics and gravitation within a single PDE framework. Although the present STM implementation is far less developed than the full Standard Model plus general relativity, its compact parameter set and explicit experimental and observational proposals nevertheless place it firmly in the domain of testable, falsifiable physics. At the same time, many elements remain incomplete: higher-loop RG running, a fully explicit chiral spectrum, a full multi-field FRG treatment of the bimodal spinor, gauge and spin–2 sectors, a more detailed treatment of black-hole microstates and evaporation, and dedicated analogue-material experiments are all substantial open programmes. **These outstanding tasks extend and refine an already calibrated and falsifiable STM benchmark; they are not prerequisites for formulating concrete predictions or for confronting the model with data.** We therefore offer STM not as a finished or mathematically unique theory, but as a concrete, quantitatively constrained and falsifiable proposal, indicating that a deterministic elasticity framework can plausibly reproduce key quantum and gravitational features. We invite rigorous analytic checks, large-scale simulations and targeted experiments to determine whether this framework can genuinely underpin quantum mechanics, gravitation and cosmology, or whether it should instead be regarded as a useful stepping stone towards a more complete synthesis.

6. Statements

- **Author contribution:** The author confirms the sole responsibility for the conception of the study, development of the STM model, analysis of the results and preparation of the manuscript.

- **Funding information:** The author received no specific funding for this work.
- **Data availability:** All data generated or analysed during this study are included in this published article and its supplementary information files.
- **Acknowledgements** I would like to express my deepest gratitude to the scholars and researchers whose foundational work is cited in the references; their contributions have been instrumental in the development of the Space-Time Membrane (STM) model presented in this paper. I am thankful for the advanced computational tools and language models that have supported the mathematical articulation of the STM model, which I have developed over the past sixteen years. Finally, I wish to pay tribute to my mother, Mavis, for my tenacity and resourcefulness; my father, James, for my imagination; my wife and children, Joanne, Elliot and Louis, for their belief in me; and to the late Isaac Asimov, whose writings first sparked my enduring curiosity in physics.
- **Conflict of interest:** The author declares no conflict of interest.
- **Ethical approval:** The conducted research is not related to either human or animals use.
- **Declaration of generative AI and AI-assisted technologies in the writing process:** During the preparation of this work the author used ChatGPT in order to improve readability of the paper and support with mathematical derivations. After using this tool/service, the author reviewed and edited the content as needed and takes full responsibility for the content of the published article.

7. Appendices A–Z

See Supplementary Information.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on [Preprints.org](https://www.preprints.org).

References

1. **Einstein, A. (1916).** Die Grundlage der allgemeinen Relativitätstheorie. *Annalen der Physik*, 49(7), 769–822.
2. **Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973).** *Gravitation*. W. H. Freeman.
3. **Wald, R. M. (1984).** *General Relativity*. University of Chicago Press.
4. **Dirac, P. A. M. (1958).** *The Principles of Quantum Mechanics*, 4th ed. Oxford University Press.
5. **Griffiths, D. J. (2005).** *Introduction to Quantum Mechanics*, 2nd ed. Pearson.
6. **Sakurai, J. J., & Napolitano, J. (2011).** *Modern Quantum Mechanics*, 2nd ed. Addison-Wesley.
7. **Polchinski, J. (1998).** *String Theory: Volume 1*. Cambridge University Press.
8. **Rovelli, C. (2004).** *Quantum Gravity*. Cambridge University Press.
9. **Hawking, S. W. (1975).** Particle Creation by Black Holes. *Communications in Mathematical Physics*, 43, 199–220.
10. **Penrose, R. (1965).** Gravitational Collapse and Space–Time Singularities. *Physical Review Letters*, 14, 57–59.
11. **Weinberg, S. (1989).** The Cosmological Constant Problem. *Reviews of Modern Physics*, 61, 1–23.
12. **Riess, A. G., et al. (1998).** Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *Astronomical Journal*, 116, 1009–1038.
13. **Perlmutter, S., et al. (1999).** Measurements of Ω and Λ from 42 High-Redshift Supernovae. *Astrophysical Journal*, 517, 565–586.
14. **Verde, L., Treu, T., & Riess, A. G. (2019).** Tensions between the early and the late Universe. *Nature Astronomy*, 3, 891–895
15. **Donoghue, J. F., Golowich, E., & Holstein, B. R. (1992).** *Dynamics of the Standard Model*. Cambridge University Press.
16. **Peskin, M. E., & Schroeder, D. V. (1995).** *An Introduction to Quantum Field Theory*. Addison–Wesley.
17. **Weinberg, S. (1995).** *The Quantum Theory of Fields, Volume I*. Cambridge University Press.

18. **Feynman, R. P., & Hibbs, A. R. (1965).** *Quantum Mechanics and Path Integrals*. McGraw–Hill.
19. **Zeh, H. D., Joos, E., Kiefer, C., Giulini, D. J. W., Kupsch, J., & Stamatescu, I. O. (2003).** *Decoherence and the Appearance of a Classical World in Quantum Theory*. Springer.
20. **Greenstein, G., & Zajonc, A. G. (2006).** *The Quantum Challenge: Modern Research on the Foundations of Quantum Mechanics*. Jones and Bartlett.
21. **Cramer, J. G. (1986).** The Transactional Interpretation of Quantum Mechanics. *Reviews of Modern Physics*, 58, 647–687.
22. **E. M. Lifshitz,** *On the Theory of Second-Order Phase Transitions. I & II*, Zhurnal Éksperimental'noi i Teoreticheskoi Fiziki (ZhETF) 11 (1941) 255; 269.
23. **P. Hořava,** *Quantum Gravity at a Lifshitz Point*, Phys. Rev. D 79 (2009) 084008. DOI: 10.1103/PhysRevD.79.084008.
24. **M. Giustina et al.,** *Significant-Loophole-Free Test of Bell's Theorem with Entangled Photons*, Physical Review Letters 115 (2015) 250401. DOI: 10.1103/PhysRevLett.115.250401.
25. **U. Sinha, C. Couteau, T. Jennewein, R. Laflamme, and G. Weihs,** *Ruling out multi-order interference in quantum mechanics*, Science 329 (2010) 418–421. DOI: 10.1126/science.1190545.
26. **A. Khrennikov,** *Born's rule from measurements of classical signals by threshold detectors which are properly calibrated*, Journal of Modern Optics (publication linked to arXiv:1105.4269). DOI: 10.1080/09500340.2012.656718.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.