

Article

Not peer-reviewed version

---

# 1,4-Disubstituted-1,2,3-triazole Linked Cyclic Ketones Ascytotoxic Agents

---

Aranav Kumar , [Sujeet Kumar](#) , Arnika Das , [Basavraj Metikurki](#) , [Dominique Schols](#) , [Subhas S Somalingappa Karki](#) \*

Posted Date: 15 May 2025

doi: 10.20944/preprints202505.1137.v1

Keywords: 1,2,3-Triazole; cytotoxicity; galectin-1 (4Y24); molecular docking



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

## Article

# 1,4-Disubstituted-1,2,3-triazole Linked Cyclic Ketones Ascytotoxic Agents

Aranav Kumar <sup>1,2</sup>, Sujeet Kumar <sup>3,†</sup>, Arnika Das <sup>1,2,†</sup>, Basavraj Metikurki <sup>3</sup>, Dominique Schols <sup>4</sup>, Subhas S. Karki <sup>1,2,\*</sup>

<sup>1</sup> Department of Pharmaceutical Chemistry, KLE College of Pharmacy, (A constituent unit of KLE Academy of Higher Education and Research-Belagavi) Bengaluru-560010, Karnataka, India

<sup>2</sup> Dr Prabhakar B Kore Basic Science Research Centre, Off-Campus, Bengaluru-560010, Karnataka, India

<sup>3</sup> Department of Pharmaceutical Chemistry, Nitte College of Pharmaceutical Sciences (A constituent college of Nitte Deemed to be University), Bengaluru-560064, Karnataka, India

<sup>4</sup> Rega Institute for Medical Research, Department of Microbiology, Immunology and Transplantation, Laboratory of Virology and Chemotherapy, KU Leuven, B-3000 Leuven, Belgium

\* Correspondence: subhaskarki@gmail.com or subhashskarki@klepharmblr.org; Tel.: +91-8023325611

† Authors contributed equally.

**Abstract:** A series of 1,2,3-triazole-linked cyclic ketones (AR 1-14) were synthesized involving the condensation of triazole-aldehydes with cyclic ketones in tetrahydrofuran (THF) and potassium hydroxide (KOH). The synthesized derivatives were evaluated for their *in-vitro* cytotoxicity against retinal (hTERT RPE-1)-1), pancreatic (Capan-1), myeloid (K-562, Hap-1), colorectal (HCT 116), lung (NCI-H460), lymphoblastic (DND-41), and non-Hodgkin lymphoma (Z-138) cells. Among the tested derivatives, molecule 2,6-bis(4-((1-(4-fluorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-7) exhibited moderate cytotoxic property with an IC<sub>50</sub> of 11.8  $\mu$ M and 13.6  $\mu$ M against pancreatic adenocarcinoma and colorectal carcinoma cells respectively. Molecular docking against galectin-1 receptor (PDB ID: 4Y24) demonstrated a favourable binding interaction (-7.1 kcal/mol) indicating a strong receptor-ligand affinity.

**Keywords:** 1,2,3-Triazole; cytotoxicity; galectin-1 (4Y24); molecular docking

## 1. Introduction

Cancer is a group of complex diseases characterized by the uncontrolled growth. In India, the estimated number of new cancer cases in 2022 was 14,61,427, and one in nine individuals is likely to develop cancer during their lifetime. Alarmingly, the incidence of cancer cases in the country is projected to increase by 12.8% by 2025[1]. In recent decades, heterocyclic chemistry has played a pivotal role in the discovery and development of new anticancer agents. Among heterocycles, nitrogen-containing frameworks are particularly valued due to their low toxicity, enhanced reactivity, and strong receptor affinity [2]. Considering this, the present study focuses on the design and development of novel 1,2,3-triazole analogs with potential anticancer activity. It was synthesized via a 1,3-dipolar cycloaddition reaction between an azide and an alkyne or aldehyde, under metal-catalyzed or metal-free conditions [3]. It is widely employed in medicinal chemistry as a bioisostere due to its chemical stability and favorable pharmacokinetic properties [4].




Galectins are the promising molecular targets in cancer therapy representing a family of  $\beta$ -galactoside-binding soluble lectins composed of approximately 130 amino acids. These proteins are distributed throughout the cytosol and nucleus, where they mediate various biological functions via N- or O-glycosylation, interacting with carbohydrate recognition domains (CRDs) [5–8]. Out of the fifteen identified galectins, galectin-1 has garnered significant attention due to its overexpression in several cancers, where it plays a key role in metastasis, tumor progression, and immune-evasion. It binds to glycosylated receptors (such as glycoproteins like CD45, CD43, or integrins) on the surface


of immune cells, particularly T cells, dendritic cells, and natural killer (NK) cells [9–12]. These characteristics make galectin-1 an appealing molecular target for the design of novel anticancer therapeutics.

## 2. Results and Discussion

### 2.1. Chemistry

Fourteen derivatives of 1,2,3-triazole-linked cyclic ketones (AR1-14) were synthesized as outlined in the Scheme 1a-d. The synthetic route commenced with the reaction of 4-hydroxybenzaldehyde (**1**) and propargyl bromide (**2**) in dry dimethyl formamide (DMF) yielding 4-(prop-2-ynoxy)benzaldehyde (**3**) (Scheme 1a). In the subsequent step (Scheme 1b), compound **3** was subjected to a copper(I)-catalyzed azide-alkyne cycloaddition (CuAAC) with different benzyl azides (**4a-h**) in the presence of sodium ascorbate ( $\text{C}_6\text{H}_7\text{NaO}_6$ ) and copper sulphate pentahydrate ( $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ), affording the corresponding triazole-benzaldehydes (**5a-h**), Scheme 1b. The final derivatives AR1-14 were synthesized by condensation of **5a-h** with cyclic ketones **6** (Scheme 1c) and **7** (Scheme 1d) in THF and KOH. The synthesized compounds were confirmed by melting point (M.p), thin-layer chromatography (TLC), Fourier-Transform Infrared (FTIR), and  $^1\text{H}$  and  $^{13}\text{C}$  Nuclear Magnetic Resonance (NMR) spectroscopy. The FTIR spectra of compounds AR 1-14 had shown aromatic C-H stretching vibrations in the range of  $3096\text{--}3000\text{ cm}^{-1}$ , while aliphatic C-H stretches appeared between  $2998\text{--}2710\text{ cm}^{-1}$ . The carbonyl ( $\text{C}=\text{O}$ ) stretching bands were noted between  $1698\text{--}1652\text{ cm}^{-1}$ . The  $^1\text{H}$  NMR spectra ( $\delta$ , ppm) showed singlets for triazole ring protons between  $\delta$  8.37–8.25 ppm. The aromatic protons appeared in the range  $\delta$  8.24–6.44, while benzylic protons were observed between  $\delta$  7.58–7.38 ppm. Methylene protons from  $-\text{OCH}_2-$  and  $-\text{NCH}_2-$  groups resonated in the ranges of  $\delta$  5.80–5.50 and  $\delta$  5.24–5.18 ppm, respectively. Methoxy ( $-\text{OCH}_3$ ) protons were evident between  $\delta$  3.73–3.62 ppm in compounds AR2, AR3, AR4, AR8, and AR9. Methyl protons in AR6 and AR11 were identified as singlet at  $\delta$  2.27 ppm. Cycloalkyl protons resonated broadly in the range of  $\delta$  3.05–1.71 ppm. The  $^{13}\text{C}$  NMR spectra data for the compounds had displayed peaks in the range of  $\delta$  195 to 188 ppm for  $\text{C}=\text{O}$  of cyclic ketones, for aromatic carbons between  $\delta$  159–106 ppm, and for methoxy and methyl carbons in the range of  $\delta$  62 to 21 ppm.





**Scheme 1. a:** Synthesis of 4-(prop-2-yn-1-yloxy)benzaldehyde(3); **b:** Synthesis of 4-((1-benzyl-1H-1,2,3-triazol-4-yl)methoxy)benzaldehydes(5a-h); **c:** 2,6-Bis(4-((1-substituted benzyl-1H-1,2,3-triazol-4-yl)oxy)benzylidene)cyclohexanone (AR 1-8); **d:** 2,6-Bis(4-((1-substituted benzyl-1H-1,2,3-triazol-4-yl)oxy)benzylidene)cyclopentanone (AR 9-14).

2.2. Biology

2.2.1. Cytotoxic Study

The *in-vitro* cytotoxicity study was conducted on different human cancerous and non-cancerous (hTERT RPE-1) cell lines, following established protocols reported in the literature. Dimethyl sulfoxide (DMSO) was employed as the solvent control, while docetaxel and staurosporine were used as reference standards. The IC<sub>50</sub> values, as summarized in Table 1, highlight compound AR-7 as the moderately active molecule in the series, exhibiting an IC<sub>50</sub> of 11.8 and 13.6 μM against the Capan-1 and HCT-116 cell lines respectively while AR-9 had shown IC<sub>50</sub> of 17.4μM against HCT-116 cells. In contrast to docetaxel and staurosporine, none of the other molecules were found to be more cytotoxic to non-cancerous hTERT RPE-1 cells, indicating a higher degree of selectivity for cancerous cells. Molecules AR 1-3 were not tested during the study, hence not included in Table 1.

**Table 1.** Cytotoxicity data (*in-vitro*) of synthesized derivatives AR4-14 and reference molecules docetaxel and staurosporine.

Compou nd	R	IC <sub>50</sub>							
		hTERT RPE-1	Capan-1	Hap-1	HCT- 116	NCI- H460	DND-41	K-562	Z-138
		<i>retina (non cancero us)</i>	<i>pancreatic adenocarcin oma</i>	<i>chroni c myeloi d leuke mia</i>	<i>colorect al carcino ma</i>	<i>lung carcino ma</i>	<i>acute lymphobla stic leukemia</i>	<i>chroni c myeloi d leuke mia</i>	<i>non- Hodgki n lympho ma</i>
AR-4	3,4,5-tri- OCH <sub>3</sub>	>100	78.2	89.7	39.9	>100	>100	>100	>100
AR-5	4-NO <sub>2</sub>	>100	58.3	>100	47.3	24.7	>100	>100	>100
AR-6	4-CH <sub>3</sub>	30.9	31.5	38.2	27.7	21.2	24.4	>100	>100
AR-7	4-F	38.1	<b>11.8</b>	31.6	<b>13.6</b>	21.9	>100	>100	>100
AR-8	4-Cl	>100	>100	>100	>100	>100	>100	>100	>100
AR-9	OCH <sub>3</sub>	73.8	>100	84.3	17.4	76.7	>100	>100	>100
AR-10	3,4,5-tri- OCH <sub>3</sub>	48.5	31.5	45.6	80.3	40.5	68.5	>100	>100

AR-11	4-CH <sub>3</sub>	40.3	25.6	42.5	29.2	20.8	37.1	>100	>100
AR-12	4-NO <sub>2</sub>	53.8	20.1	43.1	20.5	47.4	>100	>100	>100
AR-13	4-Cl	42.9	47.2	51.2	48.3	53.6	>100	>100	>100
AR-14	4-F	51.8	50	>100	67	58.1	>100	>100	40.8
Standard	Docetaxel	13.5	6.3	1.6	0.8	0.1	1.9	3.4	1.9
d	(nM)								
Standard	Staurosporine	0.4	4.6	0.3	0.3	3.2	6.4	29.8	0.3
d	(nM)								


2.2.2. In-Silico Study

In support of *in-vitro* cytotoxicity data, the binding mechanism of the most active compound AR-7 was investigated using molecular docking approach. AutoDockVina 1.1.3 was employed to dock the molecule AR-7 into the active site of the target protein, human galectin-1 (PDB ID: 4Y24). The co-crystallized ligand TD-139(bound to galectin-1) was first extracted and prepared using AutoDock Tools. The prepared TD-139 ligand was re-docked into the binding pocket of galectin-1 using the same grid configuration applied for AR-7. The re-docking of TD-139 yielded a conformation closely matching the crystal structure, with a docking score of -7.1 kcal/mol, thereby confirming the reliability of the docking methodology (Figure 1). Subsequent docking of AR-7 revealed a favourable binding pose with a docking score of -7.6 kcal/mol, indicating a potentially stronger interaction compared to TD-139. The CRD) of galectin-1 is comprisedof five carbohydrate-binding sub-sites (CBS) designated A to E (Figure 2). The co-ligand TD-139 was found to interact with sub-sites B, C, D, and E, whereas PR-7, due to its extended structure engaged with sub-sites A, B, C, and E. Detailed interaction analysis showed that:


- The 4-fluorophenyl ring of AR-7 formed an electrostatic interaction with Asp123 in sub-site A.
- The triazole ring exhibited a hydrophobic  $\pi$ -alkyl interaction with Val31 in sub-site B, Trp68 in sub-site C and an electrostatic interaction with His44 in sub-site C.
- The oxygen linker between the triazole and benzene rings participated in hydrogenbonding with His44.
- Another triazole ring formed hydrogen bonding with Asp54 in sub-site A.
- Finally, a fluorophenyl ring at the terminal end of the molecule established an electrostatic  $\pi$ -cation interaction with Arg73 in sub-site A (Figure 3).

Collectively it can be summarized that, AR-7 interacted with all conserved residues in the CBS of galectin-1, similar to TD-139 involving three electrostatic, two hydrogen bonding, and two hydrophobic interactions, supporting its stronger binding affinity and potential for greater inhibition of galectin-1 compared to other synthesized molecules in the series.






**Figure 1.** Protein-co-crystal ligand (TD-139) against human galectin-1 (PDB ID: 4Y24). The co-crystallised ligand is shown in pink and its docked pose showed in cyan.



**Figure 2.** The residue belonging to the 5 subsites in CBS of CRD in human Galectin-1 is shown in surface and stick representation. Each subsites shown in different colors. Subsite A in purple, subsite B in yellow, subsite C in pink, subsite D in orange and subsite E in cyan.



**Figure 3.** Protein-AR-7 against human galectin-1 (PDB ID: 4Y24). The compound AR-7 is shown in yellow sticks. Electrostatic, hydrogen and hydrophobic interactions are shown in orange, grey and pink dotted line respectively.

### 3. Materials and Methods

#### 3.1. Chemistry

##### 3.1.1. General Information and Instrumentation

All reagents and solvents were of analytical grade and used as received unless otherwise stated. The purity of reagents and solvents was confirmed prior to use. The progress of reactions was monitored by TLC using pre-coated Aluchrosep silica gel 60/UV254 plates (Sd Fine-Chem Ltd.). M.p was determined using the open capillary tube method in heavy liquid paraffin and were reported uncorrected. FTIR spectra were recorded using the diffuse reflectance technique with IR-grade potassium bromide (KBr) on a JASCO 460+ spectrometer. The NMR spectra were obtained using Bruker Ultraspec AMX 400 and JEOL RESONANCE spectrometers operating at 400 ( $^1\text{H}$ NMR) and 500 ( $^{13}\text{C}$  NMR) MHz in deuterated dimethyl sulfoxide ( $\text{DMSO}-d_6$ ) and chloroform ( $\text{CDCl}_3$ ). The chemical shift ( $\delta$ ) values were reported in parts per million (ppm) relative to tetramethylsilane (TMS) as the internal standard. The intermediates 4-(prop-2-ynoxy)benzaldehyde (3), benzyl azide (4a), 4-nitrobenzyl azide (4b), 4-methylbenzyl azide (4c), 4-chlorobenzyl azide (4d) and the 4-((1-benzyl-1H-1,2,3-triazol-4-yl)methoxy)benzaldehydes (5a-h) were prepared as per the literature [13]. The compounds 4-fluoro (4e), 4-methoxy (4f), 3,5-dimethoxy (4g), and 3,4,5-trimethoxybenzyl azide (4h) were synthesized according to literature reference 14.

##### 3.1.2. Synthesis of (2E,6E)-2,6-bis(4-((1-benzyl-1H-1,2,3-triazol-4-yl)methoxy)benzylidene) Cyclic Ketones (AR 1-14).

A solution of triazole aldehyde (5a-h, 1 mmol) in THF (20 mL) was mixed with an equal molar volume of a methanolic KOH solution (20 mL). To this mixture, the corresponding cyclic ketone (6 or 7, 1 mmol) was added gradually with continuous stirring and stirred at 35 °C for 16 h. Progress of reaction was monitored by TLC. Upon completion, the reaction mixture was poured into cold water and neutralized with dilute hydrochloric acid. The resulting precipitate was filtered, washed with cold water, and recrystallized from an ethanol-chloroform mixture (80:20 v/v) to yield AR1-14.

*(2E,6E)-2,6-bis(4-((1-benzyl-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-1)*

Yellow crystals, 70% yield. M.p: 160-162 °C. IR  $\nu$  (KBr): 3061, 3007, 2945, 2729, 1655, 1596, 1558, 1455, 1297  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  8.29 (2H, s, triazole-H), 7.57 (2H, s, benzyl-H), 7.52 (4H, d,  $J=8.8$ , Ar.), 7.39-7.30 (10H, m, Ar.), 7.15 (4H, d,  $J=8.8$ , Ar.), 5.61 (4H, s, 2 -OCH $_2$ -), 5.19 (4H, s, 2 -N-CH $_2$ -), 2.87 (4H, t, 2 -CH $_2$ -),  $J=10.4$ ), 1.71 (2H, p, -CH $_2$ -),  $J=24$ ).

*(2E,6E)-2,6-bis(4-((1-(4-methoxybenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-2)*

Yellow amorphous powder, 55% yield, M.p: 128-130 °C. IR  $\nu$  (KBr): 3072, 3019, 2953, 2836, 1654, 1597, 1551, 1460, 1385, 1277  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  8.25 (2H, s, triazole-H), 7.58 (2H, s, benzyl-H), 7.52 (4H, d,  $J=7.2$ , Ar.), 7.31 (4H, d,  $J=6.8$ , Ar.), 7.12 (4H, d,  $J=6.8$ , Ar.), 6.93 (4H, d,  $J=6.8$ , Ar.) 5.53 (4H, s, 2 -OCH $_2$ -), 5.18 (4H, s, 2 -N-CH $_2$ -), 3.73 (6H, s, 2 -OCH $_3$ ), 2.87 (4H, q,  $J=10.4$ , 2 -CH $_2$ -), 1.73 (2H, p, -CH $_2$ -),  $J=19.2$ ) ppm.  $^{13}\text{C}$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  188.62 (s), 159.15 (s), 158.49 (s), 142.68 (s), 135.37 (s), 134.36 (s), 132.22 (s), 131.53 (s), 129.64 (s), 127.90 (s), 124.47 (s), 114.84 (s), 114.13 (s), 113.90 (s), 61.15 (s), 55.13 (s), 52.40 (s), 27.91 (s), 22.46 (s) ppm.

*(2E,6E)-2,6-bis(4-((1-(3,5-dimethoxybenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-3)*

Yellow crystals, 61% yield, M.p: 115-118 °C. IR  $\nu$  (KBr): 3076, 2998, 2840, 1661, 1596, 1560, 1466, 1352, 1295  $\text{cm}^{-1}$ .  $^1\text{H}$ -NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  8.29 (2H, s, triazole-H), 7.57 (2H, s, benzyl-H), 7.51 (4H, d,  $J=9.2$ , Ar.), 7.12 (4H, d,  $J=8.8$ , Ar.), 6.46-6.44 (6H, m, Ar.), 5.52 (4H, s, 2 -OCH $_2$ -), 5.20 (4H, s, 2 -N-CH $_2$ -), 3.70 (12H, s, 4 -OCH $_3$ ), 2.87 (4H, t, 2 -CH $_2$ -),  $J=10.8$ ), 1.71 (2H, p, -CH $_2$ -),  $J=24.8$ ) ppm.

*(2E,6E)-2,6-bis(4-((1-(3,4,5-trimethoxybenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-4)*

Yellow crystals, 65% yield, M.p: 180-184 °C. IR  $\nu$  (KBr): 3094, 2996, 2798, 1666, 1591, 1508, 1465, 1387, 1246  $\text{cm}^{-1}$ .  $^1\text{H}$ -NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  8.30 (2H, s, triazole-H), 7.57 (2H, s, benzyl-H), 7.51 (4H, d,  $J=8.8$ , Ar.), 7.11 (4H, d,  $J=8.8$ , Ar.), 6.69 (4H, s, Ar.), 5.50 (4H, s, 2 -OCH $_2$ -), 3.72 (12H, s, 4 -OCH $_3$ ), 3.62 (6H, s, 2 -OCH $_3$ ), 2.87 (4H, t, 2 -CH $_2$ -),  $J=10.8$ ), 1.71 (2H, p, -CH $_2$ -),  $J=24.8$ ) ppm.

*(2E,6E)-2,6-bis(4-((1-(4-nitrobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-5)*

Yellow crystals, 58 % yield, M.P: 180-182 °C. IR  $\nu$  (KBr): 3086, 2982, 2714, 1660, 1598, 1558, 1345, 1455, 1464, 1229  $\text{cm}^{-1}$ .  $^1\text{H}$ -NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  8.37 (2H, s, triazole-H), 8.24 (4H, d,  $J=8.4$ , Ar.), 7.57 (2H, s, benzyl-H), 7.54-7.47 (8H, m, Ar.), 7.12 (4H, d,  $J=8.8$ , Ar.), 5.80 (4H, s, 2 -OCH $_2$ -), 5.22 (4H, s, 2 -N-CH $_2$ -), 2.86 (4H, t, 2 -N-CH $_2$ -),  $J=10.4$ ), 1.71 (2H, p, -CH $_2$ -),  $J=24.8$ ) ppm.  $^{13}\text{C}$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  188.41 (s), 158.45 (s), 147.26 (s), 143.40 (s), 142.92 (s), 135.36 (s), 134.39 (s), 132.24 (s), 131.61 (s), 129.06 (s), 128.29 (s), 125.24 (s), 123.95 (s), 114.88 (s), 113.89 (s), 61.14 (s), 51.94 (s), 27.92 (s), 22.46 (s) ppm.

*(2E,6E)-2,6-bis(4-((1-(4-methylbenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-6)*

Creamy crystals, 65% yield, M.p: 148-150 °C. IR  $\nu$  (KBr): 3096, 3027, 2946, 2877, 1654, 1595, 1557, 1456, 1420, 1294  $\text{cm}^{-1}$ .  $^1\text{H}$ -NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  8.25 (2H, s, triazole-H), 7.57 (2H, s, triazole H), 7.51 (4H, d,  $J=8.8$ , Ar.), 7.21 (4H, d,  $J=8.0$ , Ar.), 7.17 (4H, d,  $J=8.0$ , Ar.), 7.11 (4H, d,  $J=8.8$ , Ar.), 5.54 (4H, s, 2 -OCH $_2$ -), 5.18 (4H, s, 2 -N-CH $_2$ -), 2.86 (4H, t, 2 -CH $_2$ -),  $J=10.4$ ), 2.27 (6H, s, 2 -CH $_3$ ), 1.71 (2H, p, -CH $_2$ -),  $J=19.2$ ) ppm.  $^{13}\text{C}$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  188.60 (s), 158.46 (s), 142.68 (s), 137.50 (s), 135.33 (s), 134.35 (s), 132.96 (s), 132.18 (s), 129.27 (s), 128.23 (s), 127.99 (s), 124.60 (s), 114.84 (s), 61.15 (s), 52.64 (s), 27.87 (s), 22.45 (s), 20.66 (s) ppm.

*(2E,6E)-2,6-bis(4-((1-(4-fluorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-7)*



Yellow crystals, 67% yield, M.p: 118-120°C. IR  $\nu$  (KBr): 3084, 3011, 2942, 2834, 1662, 1600, 1564, 1456, 1298  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  8.29 (2H, s, triazole-H), 7.57 (2H, s, benzyl-H), 7.52 (4H, d,  $J=8.8$ , Ar.), 7.41-7.36 (4H, m, Ar.), 7.21 (4H, t,  $J=16$ , Ar.), 7.11 (4H, d,  $J=8.8$ , Ar.), 5.60 (4H, s, 2 -  $\text{OCH}_2$ -), 5.19 (4H, s, 2 -  $\text{N-CH}_2$ -), 2.87 (4H, t, 2 -  $\text{CH}_2$ -,  $J=10.4$ ), 1.73 (2H, p, -  $\text{CH}_2$ -,  $J=24.8$ ) ppm.  $^{13}\text{C NMR}$  (500 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  188.99 (s), 163.66 (s), 161.57 (s), 159.05 (s), 143.36 (s), 135.94 (s), 134.98 (s), 132.79 (s), 130.93 (s), 130.87 (s), 128.84 (s), 125.29 (s), 116.28 (s), 116.11 (s), 115.45 (s), 61.73 (s), 52.63 (s), 28.48 (s), 23.04 (s) ppm.

(2E,6E)-2,6-bis(4-((1-(4-chlorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclohexanone (AR-8)

Creamy crystals, 60% yield, M.p: 188-190°C. IR  $\nu$  (KBr): 3082, 3004, 2962, 2837, 1698, 1598, 1513, 1464, 1255  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  8.26 (2H, s, triazole-H), 7.66 (4H, d,  $J=7.2$ , Ar.), 7.40 (2H, s, benzyl-H), 7.31 (4H, d,  $J=7.2$ , ar.), 7.16 (4H, d,  $J=7.2$ , Ar.), 6.93 (4H, d,  $J=7.2$ , Ar.) 5.53 (4H, s, 2 -  $\text{OCH}_2$ -), 5.20 (4H, s, 2 -  $\text{N-CH}_2$ -), 3.73 (6H, s, 2 -  $\text{OCH}_3$ ), 3.05 (4H, s, 2 -  $\text{CH}_2$ ) ppm.

(2E,5E)-2,5-bis(4-((1-(4-methoxybenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclopentanone (AR-9)

Creamy crystals, 58% yield, M.p: 210-212°C. IR  $\nu$  (KBr): 3072, 3000, 2942, 2837, 1689, 1595, 1508, 1459, 1351, 1243  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  8.31 (2H, s, triazole-H), 7.65 (4H, d,  $J=9.2$ , Ar.), 7.38 (2H, s, benzyl-H), 7.15 (4H, d,  $J=9.2$ , Ar.), 6.66 (4H, s, Ar.), 5.50 (4H, s, 2 -  $\text{OCH}_2$ -), 5.22 (4H, s, 2 -  $\text{N-CH}_2$ -), 3.72 (12H, s, 4 -  $\text{OCH}_3$ ), 3.62 (6H, s, 2 -  $\text{OCH}_3$ ), 3.04 (4H, s, 2 -  $\text{CH}_2$ -) ppm.  $^{13}\text{C NMR}$  (500 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  189.18 (s), 159.01 (s), 153.59 (s), 143.29 (s), 137.94 (s), 135.91 (s), 134.95 (s), 132.77 (s), 131.86 (s), 128.82 (s), 125.23 (s), 115.43 (s), 106.32 (s), 61.71 (s), 60.55 (s), 56.46 (s), 53.70 (s), 28.36 (s), 23.03 (s) ppm.

(2E,5E)-2,5-bis(4-((1-(3,4,5-trimethoxybenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclopentanone (AR-10)

Brown crystals, 68% yield. M.p: 138-140°C. IR  $\nu$  (KBr): 3095, 3025, 2958, 2711, 1652, 1593, 1552, 1455, 1351, 1295  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  8.26 (2H, s, triazole-H), 7.65 (4H, d,  $J=8.8$ , Ar.), 7.39 (2H, s, benzyl-H), 7.22 (4H, d,  $J=8$ , Ar.), 7.18 (4H, d,  $J=8$ , Ar.), 7.15 (4H, d,  $J=8.8$ , Ar.), 5.55 (4H, s, 2 -  $\text{OCH}_2$ -), 5.20 (4H, s, 2 -  $\text{N-CH}_2$ -), 3.05 (4H, s, 2 -  $\text{CH}_2$ -), 2.27 (s, 6H, 2 -  $\text{CH}_3$ ) ppm.  $^{13}\text{C NMR}$  (500 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  188.42 (s), 158.46 (s), 142.80 (s), 135.37 (s), 135.00 (s), 134.38 (s), 132.89 (s), 131.53 (s), 129.92 (s), 128.78 (s), 128.26 (s), 124.86 (s), 114.87 (s), 113.94 (s), 61.15 (s), 52.02 (s), 27.91 (s), 22.47 (s) ppm.

(2E,5E)-2,5-bis(4-((1-(4-methylbenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclopentanone (AR-11)

Yellow amorphous mass, % yield 72. M.p: 208-210°C. IR  $\nu$  (KBr): 3082, 3034, 2989, 2875, 1697, 1632, 1597, 1457, 1377, 1286  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  8.37 (2H, s, triazole-H), 8.24 (4H, d,  $J=8.8$ , Ar.), 7.66 (4H, d,  $J=8.8$ , Ar.), 7.54 (4H, d,  $J=8.8$ , Ar.), 7.39 (2H, s, benzyl-H), 7.16 (4H, d,  $J=8.8$ , Ar.), 5.80 (4H, s, 2 -  $\text{OCH}_2$ -), 5.24 (4H, s, 2 -  $\text{N-CH}_2$ -), 3.05 (4H, s, 2 -  $\text{CH}_2$ -) ppm.  $^{13}\text{C NMR}$  (500 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  195.50 (s), 159.62 (s), 143.21 (s), 138.11 (s), 136.22 (s), 133.58 (s), 133.07 (s), 132.61 (s), 129.88 (s), 128.98 (s), 128.61 (s), 125.27 (s), 115.80 (s), 61.78 (s), 53.23 (s), 26.49 (s), 21.28 (s) ppm.

(2E,5E)-2,5-bis(4-((1-(4-nitrobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclopentanone (AR-12)

Yellow crystals, 65% yield, M.p: 185-187°C. IR  $\nu$  (KBr): 3078, 2882, 2710, 1665, 1600, 1550, 1350, 1498, 1471, 1220.  $^1\text{H NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  8.30 (2H, s, triazole-H), 7.66 (4H, d,  $J=8.8$ , Ar.), 7.44 (4H, d,  $J=9.2$ , Ar.), 7.39 (2H, s, benzyl-H), 7.34 (4H, d,  $J=9.2$  Ar.), 7.15 (4H, d,  $J=9.2$ , Ar.), 5.62 (4H, s, 2 -  $\text{OCH}_2$ -), 5.21 (4H, s, 2 -  $\text{N-CH}_2$ -), 3.05 (4H, s, 2 -  $\text{CH}_2$ ) ppm.

(2E,5E)-2,5-bis(4-((1-(4-chlorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclopentanone (AR-13)

Yellow amorphous, 58% yield, M.p: 228-230°C. IR  $\nu$  (KBr): 3077, 3025, 2921, 2856, 1698, 1622, 1580, 1462, 1365, 1274  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ) ppm:  $\delta$  8.30(s,2H, triazole H),

7.66(d,4H,j=8.8),7.44(d,4H,j=9.2),7.39(s,2H,benzyl-H),7.34(d, 4H,j=9.2),7.15(d, 4H, j=9.2), 5.62(s,4H, 2-OCH<sub>2</sub>-), 5.21(s,4H, 2-CH<sub>2</sub>-), 3.05 (s, 4H,cyclopentanyl). <sup>13</sup>C NMR (400 MHz, DMSO-d<sub>6</sub>) ppm: δ 159.01, 142.72, 135.65, 134.99, 132.89, 132.48, 132.02, 129.92, 128.78, 128.42, 124.89, 115.22, 61.19, 52.02, 25.91.

(2E,5E)-2,5-bis(4-((1-(4-fluorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy)benzylidene)cyclopentanone (AR-14)

Yellow crystals, 59% yield, M.p: 208-210 °C. IR ν (KBr): 3040, 2962, 2847, 1692, 1620, 1585, 1472, 1363, 1245 cm<sup>-1</sup>. <sup>1</sup>H-NMR (400 MHz, DMSO-d<sub>6</sub>) :δ 8.30(s,2H, triazole H), 7.66 (d, 4H,j=8.8), 7.41-7.37(m,6H), 7.23-7.13 (td, 8H, j=16,8.8),5.60(s,4H, 2-OCH<sub>2</sub>-), 5.21(s,4H, 2-CH<sub>2</sub>-), 3.05(s, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-) ppm.<sup>13</sup>C-NMR (500 MHz, DMSO-d<sub>6</sub>) ppm: δ 16.91, 159.05, 142.69, 135.63, 132.46, 130.34, 130.27, 124.72, 115.69, 115.52, 115.19, 61.39, 52.03, 25.89.

### 3.2. Bio-Evaluation

#### 3.2.1. Cancer Cell Lines

The human cancer cell lines Capan-1, HCT-116, NCI-H460, Hap-1, K-562, Z-138 and non cancerous hTERT RPE-1 were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA). The DND-41 cell line was sourced from the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ Leibniz-Institut, Braunschweig, Germany). All cell lines were cultured according to the suppliers' recommendations. Culture media were purchased from Gibco (Gibco Life Technologies, Merelbeke, Belgium) and supplemented with 10% fetal bovine serum (HyClone, Cytiva, MA, USA). For all biological assays, the test compounds and reference drugs were dissolved in DMSO at a concentration of 100 μM.

#### 3.2.2. Cytotoxicity Assays

Adherent cell lines were seeded at densities ranging from 500 to 1500 cells per well in 384-well plates (Greiner Bio-One, Vilvoorde, Belgium). Following overnight incubation, cells were treated with seven different concentrations of the test compounds, ranging from 100 to 0.006 μM. Untreated cell lines (i.e., without compound treatment) were used as negative controls. Suspension cell lines were seeded at densities ranging from 2500 to 5000 cells per well in 384-well culture plates containing the test compounds at the same concentration points. All cell lines were incubated for 72 h with compounds and then analyzed using the CellTiter 96® AQueous One Solution Cell Proliferation Assay (MTS) reagent (Promega, Leiden, The Netherlands) according to the manufacturer's instructions. Absorbance of the samples was measured at 490 nm using a SpectraMaxPlus 384 (Molecular Devices, CA, USA), and OD values were used to calculate the 50% inhibitory concentration (IC<sub>50</sub>). Compounds were tested in two independent experiments [15].

#### 3.2.3. In-Silico Study

The binding interaction of the most active compound in the series (AR-7) with galectin-1 protein (PDB ID: 4Y24) was investigated using AutoDockVina 1.1.2. Prior to docking, the protein structure was prepared by removing water molecules, adding hydrogen atoms, and assigning appropriate atom types and partial charges to each atom. The chemical structure of PR-7 was constructed using ChemBioDraw Ultra 12.0, and its geometry was optimized using the MMFF94 force field. Energy minimization was carried out with a maximum of 1000 iterations and a root mean square (RMS) gradient threshold of 0.1 kcal/mol. The minimized structure was further prepared using the AutoDock Tools (ADT) package, where Gasteiger charges were assigned, hydrogens added, and rotatable bonds were defined. Both ligand and receptor structures were saved in .pdbqt format, as required for docking. Docking was performed within a defined binding site using a grid box of 40 × 80 × 40 points with a grid spacing of 0.375 Å. The centre of the grid box was set at coordinates X = 22.081, Y = 0.3, Z = -16.435. The docking simulations were run on a Linux-based multi-core CPU platform. Among the generated poses, the conformation with the lowest binding energy was selected

and aligned to the active site of the receptor crystal structure for visualization and interaction analysis [16].

#### 4. Conclusions

A series of fourteen 1,2,3-triazole-linked cyclic ketones (AR 1-14) were synthesized, purified, and structurally characterized. Molecular docking was performed on galectin-1 (PDB ID: 4Y24) with TD-139 as the reference ligand. Among the synthesized derivatives, PR-7 exhibited a docking score of -7.1 Kcal/mol comparable to that of TD-139 (-7.6 kcal/mol) indicating strong binding affinity toward the galectin-1 active site. Cytotoxicity screening was performed across eight human cell lines, including both cancerous and non-cancerous types. PR-7 demonstrated the highest inhibitory effect specifically against the Capan-1 pancreatic cancer cell line. However, none of the synthesized derivatives surpassed the cytotoxic efficacy of standard drugs docetaxel and staurosporine. These findings suggest that further structural optimization is required to enhance the anticancer potency of this scaffold.

**Supplementary Materials:** The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Figures S1-S16e: <sup>1</sup>H-NMR spectra, Figures S17-27d: <sup>13</sup>C-NMR spectra.

**Author Contributions:** Conceptualization& data curation, Dr. Subhas S Karki (SSK); Synthetic investigation, Aranav Kumar (AK)&Amika Das (AD); Data interpretation and original draftpreparation, Sujeet Kumar (SK); Software, BasavrajMetikurki (BM); *In-vitro* investigation, Dominique Schols(DS). All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** The data are available as supplementary file.

**Acknowledgments:** The authors would like to acknowledge KLE College of Pharmacy and Dr. Prabhakar B. Kore Basic Science Research Centre (Off-Campus) Bengaluru-560010, Karnataka, India for the support to carry-out the submitted work.

**Conflicts of Interest:** The authors declare no conflicts of interest.

#### Abbreviations

The following abbreviations are used in this manuscript:

ADT	AutoDock Tools
ATCC	American Type Culture Collection
CBS	Carbohydrate-Binding Sub-sites
CRD	Carbohydrate Recognition Domain
CuAAC	Copper(I)-catalyzed Azide-Alkyne Cycloaddition
DMF	Dimethyl Formamide
DMSO	Dimethyl Sulfoxide
DSMZ	Deutsche Sammlung von Mikroorganismen und Zellkulturen
FTIR	Fourier-Transform Infrared
M.p	Melting Point
NK	Natural Killer
NMR	Nuclear Magnetic Resonance
OD	Optical Density
PDB	Protein Data Bank
ppm	Parts Per Million
RMS	Root Mean Square
THF	Tetrahydrofuran
TLC	Thin-Layer Chromatography
TMS	Tetramethylsilane

## References

1. Sathishkumar, K.; Chaturvedi, M.; Das, P.; Stephen, S.; Mathur, P. Cancer incidence estimates for 2022 & projection for 2025: Result from National Cancer Registry Programme, India. *Indian J Med Res* 2022, 156, 598-607.
2. Kumar, A.; Singh, A.K.; Singh, H.; Vijayan, V.; Kumar, D.; Naik, J.; Thareja, S.; Yadav, J.P.; Pathak, P.; Grishina, M.; Verma, A.; Khalilullah, H.; Jaremko, M.; Emwas, A.H.; Kumar, P. Nitrogen containing heterocycles as anticancer Agents: a medicinal chemistry perspective. *Pharmaceuticals (Basel)* 2023, 14, 299.
3. Disha, P.V.; Ruturajsinh, M.V.; Hitendra, M.P. Versatile synthetic platform for 1,2,3-triazole chemistry. *ACS Omega* 2022, 7, 36945-36987.
4. Malik, M.S.; Ahmed, S.A.; Althagafi, I.I.; Ansari, M.A.; Kamal, A. Application of triazoles as bioisosteres and linkers in the development of microtubule targeting agents. *RSC Med Chem* 2020, 11, 327-348.
5. Johannes, L.; Jacob, R.; Leffler, H. Galectins at a glance. *J Cell Sci* 2018, 131, jcs208884.
6. Camby, I.; Le Mercier, M.; Lefranc, F.; Kiss, R. Galectin-1: a small protein with major functions. *Glycobiology* 2006, 16, 137R-157R.
7. Cedeno-Laurent, F.; Dimitroff, C.J. Galectin-1 research in T cell immunity: past, present and future. *Clin Immunol* 2012, 142, 107-16.
8. Rabinovich, G. Galectin-1 as a potential cancer target. *Br J Cancer* 2005, 92, 1188-1192.
9. Carlos, P.M.P.; Juan, B.C.; Santiago, D.L.; Marcelo, A.M. The structural biology of Galectin-ligand recognition: current advances in modeling tools, protein engineering, and inhibitor design. *Front Chem* 2019, 7, 1-14.
10. Sridhar, G.N. Non-carbohydrate galectin-1 inhibitors as promising anticancer agents: design strategies, structure activity relationship and mechanistic insights. *Eur J Med Chem Rep* 2024, 11, 100170.
11. Cousin, J.M.; Cloninger, M.J. The role of galectin-1 in cancer progression, and synthetic multivalent systems for the study of Galectin-1. *Int J Mol Sci* 2016, 17, 1566.
12. Huang, Y.; Wang, H.C.; Zhao, J.; Wu, M.H.; Shih, T.C. Immunosuppressive roles of galectin-1 in the tumor microenvironment. *Biomolecules* 2021, 11, 1398.
13. Das, A.; Greco, G.; Kumar, S.; Catanzaro, E.; Morigi, R.; Locatelli, A.; Schols, D.; Alici, H.; Tahtaci, H.; Ravindran, F.; Fimognari, C.; Karki, S.S. Synthesis, *in-vitro* cytotoxicity, molecular docking and ADME study of some indolin-2-one linked 1,2,3-triazole derivatives. *Comput Biol and Chem* 2022, 97, 107641.
14. Shaik, P.S.; Nayak, V.L.; Sultana, F.; SubbaRao, A.V.; Shaik, A.B.; Korrapati, S.B.; Ahmed, K. Design and synthesis of imidazo [2,1-b]thiazole linked triazole conjugates: microtubule-destabilizing agents. *Eur J Med Chem* 2017, 126, 36-51.
15. Walle, T.V.; Theppawong, A.; Grootaert, C.; Jonghe, S.D.; Persoons, L.; Daelemans, D.; Hecke, K.V.; Camp, J.V.; D'hooghe, M. Synthesis and cytotoxic evaluation of monocarbonylcurcuminoids and their pyrazoline derivatives. *Monatsh Chem* 2019, 150, 2045-2051.
16. Trott, O.; Olson, A.J. AutoDockVina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem* 2010, 31, 455-461.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.