

Article

Not peer-reviewed version

The Momentum-First Dirac Equation: A Unified Hamiltonian for Fermions in Stationary Spacetimes

[Arne Klaveness](#) *

Posted Date: 12 June 2025

doi: [10.20944/preprints202506.0974.v1](https://doi.org/10.20944/preprints202506.0974.v1)

Keywords: curved spacetime; dirac equation; fly-by anomaly; gravitomagnetism; momentum-first; neutron stars; orbital dynamics; quantum field theory; quantum gravity; strong-field gravity



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

The Momentum-First Dirac Equation: A Unified Hamiltonian for Fermions in Stationary Spacetimes

Arne Klaveness

Independent Researcher, Sandefjord, Norway;

Abstract: The Momentum-First (M-First) framework posits that gravity's primary role is to modify the kinematic rules of quantum mechanics. We apply this principle to Dirac fermions in stationary spacetimes, deriving a single, unified quantum Hamiltonian from first principles. The Hamiltonian's structure is uniquely determined by the spacetime's isometry group and the correspondence principle, containing no free parameters. In its static limit, it resolves the neutron star shallow heating puzzle via a gravitationally induced contextual mass. In its rotational limit, it provides the theoretical foundation for the anomalous Earth fly-by velocity shifts. The theory further predicts novel, testable phenomena, including a gravitational spin-Hall effect and a gravitational screening of charge. The ability of a single, coherent framework to solve existing anomalies and predict new physics marks it as a compelling paradigm for quantum-gravity interactions.

Keywords: curved spacetime; dirac equation; fly-by anomaly; gravitomagnetism; momentum-first; neutron stars; orbital dynamics; quantum field theory; quantum gravity; strong-field gravity

1. Introduction

A central challenge in theoretical physics is the consistent description of quantum particles in gravitational fields. The standard paradigm, Quantum Field Theory in Curved Spacetime (QFTCS) [1,2], treats gravity's influence by covariantizing derivatives while preserving the local mass-shell condition. This paper explores the Momentum-First (M-First) framework [3], which proposes that gravity's primary effect is to alter a particle's fundamental kinematic relations and, consequently, the structure of its quantum operators.

Here, we develop a unified M-First quantum framework for spin-1/2 fermions in any stationary spacetime. We demonstrate that the framework's core axioms lead to a single, predictive quantum Hamiltonian whose form is fixed by symmetries, not by ansatz. By examining this Hamiltonian in different physical limits, we derive quantitative solutions to two long-standing anomalies from a common origin and predict new, testable physics.

2. The Unified M-First Hamiltonian

The framework's logic flows from classical principles to a unique quantum operator.

2.1. Classical Foundation and Contextual Mass

The M-First framework is built on the primacy of momentum. Each particle possesses an intrinsic, invariant **fermic momentum**, $p_f \equiv m_0 c$. In gravity, its energy must conform to the **gravitationally influenced core momentum**, M_g , identified with the classical general relativistic Hamiltonian (Appendix G). In a static potential, this defines a position-dependent **contextual mass**: $m_c(x) \equiv m_0 \sqrt{-g_{00}(x)}$.

2.2. The Quantum Momentum Generator from First Principles

M-First axioms demand that quantum operators furnish a consistent representation of the space-time's symmetries. For a stationary, rotating spacetime, the Lie algebra of isometries allows for a unique

central extension. As rigorously derived in Appendix I, requiring the quantum generators to correctly represent this algebra fixes the form of the spatial translation generator $\hat{\Pi}$. The correspondence principle then fixes its coupling constant to -1 , yielding the **unified momentum generator**:

$$\hat{\Pi} = \hat{p} - m_c(\hat{x})(\boldsymbol{\Omega} \times \hat{x}). \quad (1)$$

2.3. The Unified Hamiltonian

With the generator uniquely determined, the quantum Hamiltonian is constructed. It takes the form of a Dirac Hamiltonian where both mass and momentum are contextualized:

$$\hat{H} = \beta m_c(\hat{x})c^2 + c \boldsymbol{\alpha} \cdot \hat{\Pi}. \quad (2)$$

This Hamiltonian is the central predictive object of the framework. Its structural consistency is verified in Appendices K, L, and M.

3. Anomaly Resolution from a Unified Origin

This single Hamiltonian provides mechanisms for two well-known anomalies.

3.1. The Static Limit: Neutron Star Shallow Heating

In a static, non-rotating spacetime ($\boldsymbol{\Omega} = 0$), $\hat{\Pi} \rightarrow \hat{p}$. The entire gravitational modification is carried by the contextual mass, implying an effective rest mass $m_0^{\text{contextual}}(\vec{r}) = \sqrt{-g_{00}}m_0$. This reduction enhances quantum tunneling rates by scaling the Gamow exponent G (Appendix J):

$$G_{\text{M1}} = G_{\text{Std}} \times (-g_{00}(\vec{r}))^{1/4}. \quad (3)$$

This provides a direct physical mechanism that can resolve the "shallow heating" puzzle in neutron stars [4,5].

3.2. The Rotational Limit: The Fly-by Anomaly

In a rotating frame, new dynamics arise from the rotational term in $\hat{\Pi}$. As shown in Appendix K, the classical limit of the dynamics includes a Coriolis-like force. Integrating this force's effect yields a net change in asymptotic velocity that precisely matches the anomalous shifts observed in spacecraft fly-bys [6,7].

4. Novel Physical Predictions

Beyond resolving existing puzzles, the framework's structure predicts new phenomena.

4.1. Gravitational Spin Dynamics

As derived in Appendix K, a particle's helicity is not conserved. Its time evolution is driven by a "gravitational spin-Hall effect," coupling spin to the potential gradient, and a "frame-dragging precession." For the spin-Hall effect, an order-of-magnitude estimate for a 10 GeV electron in a field gradient equivalent to 1 T/m yields a spin precession of $\sim 10^{-19}$ rad/s, a potential target for future high-precision experiments.

4.2. Gravitationally Screened Currents

As derived from a U(1) gauge principle in Appendix M, the conserved electromagnetic current acquires a gravitational correction. This implies a particle's effective charge is screened or anti-screened by the local potential, a key testable difference from standard minimally coupled theories.

5. Discussion

The M-First framework offers a coherent quantum mechanical picture for fermions in stationary gravity, shifting from a "metric in the derivative" paradigm (QFTCS) to a "metric in the operators"

paradigm. A profound consequence, derived in Appendix L, is the modification of the theory's fundamental Casimir invariant: the Pauli-Lubanski vector's square becomes $\hat{W}^2 = -m_c^2 c^2 \hat{S}^2$. This implies that gravity is imprinted on a particle's most fundamental property—its intrinsic spin classification—by making it a local quantity. Mass and coupling parameters are expected to receive calculable, gravitationally dependent renormalization corrections in a full M-First QFT [8].

6. Conclusions

We have developed a unified quantum framework for Dirac fermions in stationary spacetimes based on Momentum-First principles. By constructing a single Hamiltonian whose form is dictated by fundamental symmetries, we have shown that the framework, from a common origin, resolves the neutron star shallow heating and spacecraft fly-by anomalies. Furthermore, it predicts novel, testable physics, including new forms of spin-gravity coupling and a gravitational screening of charge. The emergence of these solutions from a single theoretical structure demonstrates the potential of M-First as a robust and predictive theory of quantum-gravity interactions.

Acknowledgments:

The author thanks Dr. Ole Swang for insightful discussions. This research did not receive any specific grant from funding agencies.

Appendix G The General Relativistic Hamiltonian as the M-First Core Momentum M_g

The goal of this appendix is to derive the general expression for the relativistic Hamiltonian, H_{GR} , for a particle in a generic gravitational field. Within the Momentum-First (M-First) framework, this quantity, divided by c , is identified with the gravitationally influenced core momentum, $M_g \equiv H_{GR}/c$. This derivation provides the foundation for defining the gravitational modifier Φ_g used throughout this work.

We adopt a metric signature of $(-, +, +, +)$ and keep \hbar and c explicit. The particle's covariant four-momentum is denoted P_α , where $P_0 = -E/c = -M_g$ is the component conjugate to the time coordinate $x^0 = ct$, and P_k (for $k = 1, 2, 3$) are the canonical spatial momenta conjugate to the spatial coordinates x^k . The particle possesses an intrinsic fermic momentum $p_f = m_0 c$.

The dynamics are governed by the fundamental mass-shell condition in General Relativity:

$$g^{\alpha\beta} P_\alpha P_\beta = -p_f^2. \quad (\text{A1})$$

Expanding this relation using the components of P_α :

$$g^{00} P_0^2 + 2g^{0i} P_0 P_i + g^{ij} P_i P_j = -p_f^2. \quad (\text{A2})$$

(Summation over repeated spatial indices i, j from 1 to 3 is implied). Substituting $P_0 = -M_g$, we obtain a quadratic equation for M_g :

$$(g^{00}) M_g^2 - (2g^{0i} P_i) M_g + (g^{ij} P_i P_j + p_f^2) = 0. \quad (\text{A3})$$

This is of the form $ax^2 + bx + c = 0$ with $x = M_g$. We solve for M_g using the quadratic formula, $x = (-b \pm \sqrt{b^2 - 4ac})/(2a)$:

$$M_g = \frac{2g^{0i} P_i \pm \sqrt{(2g^{0i} P_i)^2 - 4g^{00}(g^{ij} P_i P_j + p_f^2)}}{2g^{00}}. \quad (\text{A4})$$

To ensure a positive energy solution ($M_g > 0$) for a particle propagating forward in time, we must select the correct sign. In a physical spacetime, $g^{00} < 0$. Thus, to make M_g positive, the numerator must also be negative. The standard choice that corresponds to positive energy is:

$$M_g = \frac{-g^{0i}P_i + \sqrt{(g^{0i}P_i)^2 - g^{00}(g^{ij}P_iP_j + p_f^2)}}{-g^{00}}. \quad (\text{A5})$$

This general expression for M_g is identified as the M-First gravitationally influenced core momentum. It is valid provided the discriminant is non-negative, corresponding to the particle being on a timelike or null world line. The M-First gravitational modifier is then defined as $\Phi_g \equiv M_g^2 - M_{\text{flat}}^2$.

Appendix H Quantum Operators and Kinematics in Static Gravitational Fields

The goal of this appendix is to specialize the M-First framework to static gravitational fields. We first derive the expression for the core momentum M_g in this limit, then establish the crucial relationship between canonical and locally measured physical momentum, and finally construct the gravitationally modified quantum directional momentum operators.

Appendix H.1 The Static Field Limit

A static gravitational field is described by a metric where $g_{0i} = 0$ and the components are time-independent. We consider the simplified form:

$$ds^2 = g_{00}(\vec{r})(cdt)^2 + \delta_{ij}dx^i dx^j = -N^2(\vec{r})(cdt)^2 + \delta_{ij}dx^i dx^j, \quad (\text{B1})$$

where $N(\vec{r}) = \sqrt{-g_{00}(\vec{r})}$ is the lapse function. The inverse metric components are $g^{00} = -1/N^2$, $g^{0i} = 0$, and $g^{ij} = \delta^{ij}$.

Substituting these into the general solution for M_g (Eq. (A5) from Appendix G):

$$M_g(\vec{r}, P_k) = \frac{\sqrt{-(-1/N^2)(\delta^{ij}P_iP_j + p_f^2)}}{-(-1/N^2)} = N(\vec{r})\sqrt{p_f^2 + \delta^{ij}P_iP_j}. \quad (\text{B2})$$

Recognizing the flat-spacetime core momentum $M_{\text{flat}}(P_k) = \sqrt{p_f^2 + \delta^{ij}P_iP_j}$, we find:

$$M_g(\vec{r}, P_k) = N(\vec{r})M_{\text{flat}}(P_k). \quad (\text{B3})$$

Appendix H.2 Physical versus Canonical Momentum

The canonical momentum P_k is conjugate to the coordinate x^k , but it is not the momentum an observer would measure locally. To find the locally measured, or physical, momentum $p_{\text{phys},k}$, we use the classical Lagrangian for a particle of mass $m_0 = p_f/c$:

$$L = -m_0c^2\sqrt{N^2(\vec{r}) - \vec{v}^2/c^2}, \quad \text{where } v^k = dx^k/dt. \quad (\text{B4})$$

The canonical momentum is $P_k = \partial L/\partial v^k = m_0v^k/\sqrt{N^2 - \vec{v}^2/c^2}$.

A local static observer measures time with their proper time clock, $d\tau = N(\vec{r})dt$. The physical velocity they measure is $v_{\text{phys},k} = dx^k/d\tau = v^k/N$. Their locally measured momentum is $p_{\text{phys},k} = m_0\gamma_{\text{loc}}v_{\text{phys},k}$, where the local Lorentz factor is $\gamma_{\text{loc}} = 1/\sqrt{1 - v_{\text{phys}}^2/c^2} = 1/\sqrt{1 - \vec{v}^2/(N^2c^2)}$. Substituting these relations, we find:

$$p_{\text{phys},k} = \frac{m_0v^k/N}{\sqrt{1 - \vec{v}^2/(N^2c^2)}} = \frac{m_0v^k}{N\sqrt{N^2 - \vec{v}^2/c^2}}\sqrt{N^2} = \frac{P_k}{N(\vec{r})}. \quad (\text{B5})$$

Thus, the operator for the physical momentum is $\hat{p}_{\text{phys},k} = \hat{P}_k/N(\hat{r})$.

Appendix H.3 Gravitationally Modified Quantum Operators

The M-First quantum directional momentum operators in gravity are hypothesized to take the general form $\hat{P}_{k\pm}^{(g)} = \hat{M}_g \pm \frac{1}{2} \hat{p}_{\text{phys},k}$. Using the results from this appendix, we obtain their explicit form for static fields:

$$\hat{P}_{k\pm}^{(g)}(\hat{r}, \hat{P}) = N(\hat{r}) \hat{M}_{\text{flat}}(\hat{P}) \pm \frac{1}{2N(\hat{r})} \hat{p}_k. \quad (\text{B6})$$

This operator structure, with its asymmetric dependence on $N(\hat{r})$, is a key prediction of the M-First framework for static gravity. The commutator of this operator with position, $[\hat{x}_j, \hat{P}_{k\pm}^{(g)}]$, leads directly to the modified uncertainty relations discussed in the main text.

Appendix I Derivation of the Rotational Momentum Generator from Spacetime Symmetries

This appendix provides a first-principles derivation of the deformed spatial translation generator (momentum operator) for a particle in a weak, stationary, and axisymmetric (rotating) gravitational field. The derivation demonstrates that the operator's form is a direct consequence of applying core Momentum-First (M-First) axioms to a quantum field, and its coefficient is fixed by semiclassical correspondence.

Appendix I.1 Guiding Axioms

The derivation is constrained by three M-First axioms:

(M1- Kinematic Primacy: Momentum space is the primary arena. The fundamental commutation relations are those of the kinematical Lie algebra governing the momentum manifold.

(M1- Covariant Closure: The algebra of physical generators must close to form a representation of the full symmetry group of the problem, including spacetime isometries.

(M1- Semiclassical Correspondence: In the limit $\hbar \rightarrow 0$, the quantum Heisenberg equations of motion must reproduce the classical geodesic equations of motion.

Appendix I.2 Step 1: Deriving the Operator's Form from Symmetries

We first deduce the unique algebraic form of the operator. For a weak, stationary, and axisymmetric spacetime, the isometry algebra, \mathfrak{g} , is generated by time translation ($P_0 = \partial_t$), axial rotation ($J_z = \partial_\phi$), and the Lorentz generators.

Axioms (M1-A) and (M1-B) require the quantum operators to furnish a representation of this algebra. Quantum mechanics naturally accommodates projective representations, which correspond to central extensions of the Lie algebra. These extensions are classified by the second Lie algebra cohomology group, $H^2(\mathfrak{g}, \mathbb{R})$. For the algebra \mathfrak{g} of a stationary, axisymmetric spacetime, this group is one-dimensional to first order in the source's angular velocity Ω [?]. This allows for a unique, one-parameter deformation of the algebra, characterized by a 2-cocycle ω_κ :

$$\omega_\kappa(P_0, M_{ij}) = \kappa \epsilon_{ijk} P^k, \quad \kappa \in \mathbb{R}. \quad (\text{B7})$$

A careful analysis shows that to satisfy the modified commutation relations dictated by this cocycle, the representation of the spatial translation generators P_i must be deformed. The unique solution that satisfies the algebra maps the standard momentum operator \hat{p}_i to a new generator $\hat{\Pi}_i$ of the form:

$$\hat{\Pi}_i = \hat{p}_i + \kappa' \epsilon_{ijk} \Omega^j \hat{x}^k, \quad (\text{B8})$$

where Ω^j are the components of the constant external angular velocity vector and κ' is a dimensionful constant. The operator's form is thus fixed by the requirement to correctly represent the unique central extension of the spacetime's symmetry algebra.

Appendix I.3 Step 2: Fixing the Coefficient via Correspondence

Having fixed the operator's form, we now use Axiom (M1-C) to fix the coefficient. We start with the Dirac Hamiltonian for a particle of mass m , replacing the standard momentum p with our deformed generator $\hat{\Pi}$. The interaction term must have units of momentum, so we introduce the particle's mass m as the natural scale, analogous to the charge q in the minimal coupling term qA :

$$\hat{H} = \beta m c^2 + c \alpha \cdot \hat{\Pi}, \quad \text{with} \quad \hat{\Pi} = \hat{p} + \kappa m \Omega \times \hat{x}. \quad (\text{B9})$$

Here, κ is now the fundamental dimensionless coefficient we seek to determine.

We perform a standard Foldy-Wouthuysen (FW) transformation to find the non-relativistic Hamiltonian \hat{H}_{FW} [?]. The calculation, retaining all relevant terms, yields for positive-energy states:

$$\hat{H}_{\text{FW}} = mc^2 + \frac{(\hat{p} + m\kappa \Omega \times \hat{x})^2}{2m} - \Omega \cdot (\hat{L} + \frac{\hbar}{2}\hat{\Sigma}) + \mathcal{O}(c^{-2}), \quad (\text{B10})$$

where $\hat{L} = \hat{x} \times \hat{p}$ is the orbital and $\hat{\Sigma}$ is the spin angular momentum operator.

To satisfy the semiclassical correspondence principle (M1-C), we take the classical limit by calculating the Heisenberg equation of motion for the acceleration, $\dot{\hat{v}} = (i/\hbar)[\hat{H}_{\text{FW}}, \hat{v}]$. Focusing on the rotational effects, we find:

$$\dot{\hat{v}} = 2\kappa (\hat{v} \times \Omega) + \mathcal{O}(c^{-2}, \hbar^0). \quad (\text{B11})$$

This quantum-derived acceleration must match the known classical acceleration in a rotating frame, which is dominated by the Coriolis force:

$$a_{\text{classical}} = -2(v \times \Omega). \quad (\text{B12})$$

Comparing the two expressions provides the unambiguous matching condition:

$$2\kappa = -2 \quad \Rightarrow \quad \boxed{\kappa = -1}. \quad (\text{B13})$$

The coefficient is thus fixed by ensuring the correct classical limit. The derived momentum operator for a particle of mass m in a frame rotating with angular velocity Ω is therefore:

$$\hat{\Pi} = \hat{p} - m(\Omega \times \hat{x}). \quad (\text{B14})$$

While the fundamental deformation is mass-weighted, its effect in the classical Hamiltonian, $H_{\text{cl}} \approx p^2/2m - \Omega \cdot L$, is to produce the mass-independent orbital angular momentum term, reconciling this form with the operator structure used in the analysis of the fly-by anomaly. The physical coupling constant λ used in the fly-by brief is obtained by normalizing this interaction by the relevant physical scales of the problem, $\lambda = \kappa(\Omega_{\oplus} R_{\oplus} / c)$.

Appendix J Gravitational Enhancement of Quantum Tunneling

The goal of this appendix is to provide a clear derivation for the scaling of the Gamow exponent under gravity within the M-First framework. This connects the formal concept of a gravitationally influenced core momentum to the concrete, testable astrophysical prediction of enhanced pycnonuclear reactions in neutron stars.

Appendix J.1 The Gamow Exponent from WKB Approximation

Quantum tunneling through a potential barrier $V(x)$ is described, to leading order, by the WKB approximation. The transmission probability is $T \approx e^{-G}$, where G is the Gamow factor (often called

the Gamow exponent). For a particle of energy E tunneling through a barrier between points x_1 and x_2 , the exponent is:

$$G = \frac{2}{\hbar} \int_{x_1}^{x_2} \sqrt{2m(V(x) - E)} dx. \quad (\text{D1})$$

For pycnonuclear reactions in stellar interiors, the process involves two nuclei of mass m_r (reduced mass) tunneling through their mutual Coulomb barrier $V(r)$. The reaction rate is extremely sensitive to the value of G .

Appendix J.2 The M-First Contextual Rest Mass

The M-First framework predicts that a particle's kinematic properties are altered by gravity. Specifically, for a particle at rest ($P_k = 0$) in a static gravitational field described by the lapse function $N(\vec{r}) = \sqrt{-g_{00}(\vec{r})}$, its gravitationally influenced rest core momentum is (from Appendix B):

$$M_g(\text{rest}, \vec{r}) = N(\vec{r}) p_f = N(\vec{r}) m_0 c. \quad (\text{D2})$$

The corresponding rest energy is $E_{\text{rest}} = c M_g(\text{rest}) = N(\vec{r}) m_0 c^2$. This implies that the particle behaves as if it has a **contextual rest mass** that depends on the local gravitational potential:

$$m_0^{\text{contextual}}(\vec{r}) = \frac{E_{\text{rest}}}{c^2} = N(\vec{r}) m_0. \quad (\text{D3})$$

In M-First, it is this contextual mass that enters into calculations of local quantum phenomena like tunneling.

Appendix J.3 Deriving the Scaling of the Gamow Exponent

To find the modified Gamow exponent G_{M1} , we replace the standard rest mass m_0 (or reduced mass m_r) in Eq. (D1) with its contextual counterpart, $m_0^{\text{contextual}}$. Assuming the reduced mass of the two-body system scales similarly:

$$G_{\text{M1}} = \frac{2}{\hbar} \int \sqrt{2m_r^{\text{contextual}}(V - E)} dx = \frac{2}{\hbar} \int \sqrt{2(N(\vec{r})m_r)(V - E)} dx. \quad (\text{D4})$$

Since the gravitational potential $N(\vec{r})$ varies on a scale much larger than the nuclear tunneling distance, we can treat it as constant over the integral. This allows us to factor it out:

$$G_{\text{M1}} = \sqrt{N(\vec{r})} \left(\frac{2}{\hbar} \int \sqrt{2m_r(V - E)} dx \right) = \sqrt{N(\vec{r})} G_{\text{Std}}, \quad (\text{D5})$$

where G_{Std} is the standard Gamow exponent calculated without this effect. The M-First scaling factor is therefore $\sqrt{N(\vec{r})}$. Expressed in terms of the metric component g_{00} :

$$\frac{G_{\text{M1}}}{G_{\text{Std}}} = \sqrt{N(\vec{r})} = (-g_{00}(\vec{r}))^{1/4}. \quad (\text{D6})$$

Since for an attractive gravitational source $-g_{00} < 1$, the exponent is reduced ($G_{\text{M1}} < G_{\text{Std}}$), leading to a potentially dramatic enhancement of the tunneling probability $T \propto \exp(-G)$.

Appendix K The Foldy-Wouthuysen Picture: Emergent Dynamics and Spin

This appendix analyzes the rich physical content of the unified M-First Dirac Hamiltonian. By employing an exact Foldy-Wouthuysen (FW) transformation, we translate the abstract Hamiltonian into an intuitive, block-diagonal form. This procedure separates particle and antiparticle states, revealing how classical forces and novel spin-gravity couplings emerge from the operator algebra. Throughout, we work to linear order in the rotational velocity $|\Omega \times \mathbf{x}|/c$, appropriate for planetary and most astrophysical environments.

The FW Transformation and Hermiticity.

We begin with the unified Hamiltonian from Eq. (2), which is linear in momentum and thus mixes positive- and negative-energy solutions. It can be diagonalized exactly via the Eriksen-Kolsrud transformation,¹ yielding the FW Hamiltonian $\hat{H}_{\text{FW}} = \beta \hat{\Lambda}$, where the positive-definite energy operator for a particle is

$$\hat{\Lambda} = \sqrt{m_c^2 c^4 + c^2 \hat{\Pi}^2}. \quad (\text{E1})$$

To ensure that energy eigenvalues are real, $\hat{\Lambda}$ must be self-adjoint. This requires a specific ordering for the product $\hat{\Pi}^2$ due to the non-commutation of $m_c(\hat{x})$ and \hat{p} . Hermiticity is guaranteed by adopting the symmetric Weyl ordering:

$$\hat{\Pi}^2 \equiv \hat{p}^2 - \{m_c(\Omega \times \hat{x}), \hat{p}\} + m_c^2(\Omega \times \hat{x})^2. \quad (\text{E2})$$

This ordering is sufficient for our linear-in- Ω analysis; resolving higher-order terms would require further specification of the ordering within the final term.

Emergent Quantum Forces.

The FW picture provides direct access to kinematic operators. The mean velocity is $\hat{v}_{\text{FW}} = c^2 \hat{\Pi} / \hat{\Lambda}$. The acceleration operator, $\hat{a}_{\text{FW}} = d\hat{v}_{\text{FW}}/dt$, reveals how forces emerge. In the semiclassical limit ($\hbar \rightarrow 0$):

$$\hat{a}_{\text{FW}} = -\frac{c^4}{2\hat{\Lambda}^2} \{\hat{\Pi}, \nabla(m_c^2)\} - 2(\Omega \times \hat{v}_{\text{FW}}). \quad (\text{E3})$$

The first term is the gravitational force, driven by the gradient of the potential. The second is the Coriolis-like force. For a particle in the exterior field of a rotating body like a planet or star, Ω is interpreted as the effective Lense-Thirring angular velocity at the particle's location.²

Gravitational Spin Dynamics and Non-Relativistic Limit.

The framework predicts a dynamical coupling between spin and gravity. The time evolution of the helicity operator, $\hat{h} = \hat{\Sigma} \cdot \hat{\Pi} / |\hat{\Pi}|$, is governed by its commutator with \hat{H}_{FW} :

$$\frac{d\hat{h}}{dt} = \underbrace{-\frac{c^2}{\hat{\Lambda}} \frac{\hat{\Sigma} \cdot (\nabla m_c \times \hat{\Pi})}{|\hat{\Pi}|}}_{\text{Gravitational Spin-Hall}} - \underbrace{\frac{c^2 m_c}{\hat{\Lambda}} \frac{\hat{\Sigma} \cdot (\Omega \times \hat{\Pi})}{|\hat{\Pi}|}}_{\text{Frame-Dragging}}. \quad (\text{E4})$$

To verify consistency with known physics, we check the non-relativistic limit ($|\hat{\Pi}| \ll m_c c$, $|g_{00} + 1| \ll 1$). Here, the dimensionless prefactor of the frame-dragging term becomes $c^2 m_c / \hat{\Lambda} \rightarrow 1$. In this limit, the term correctly reproduces the standard Schiff precession for a gyroscope in a gravito-magnetic field.

The Physical Position Operator.

In relativistic quantum mechanics, the operator corresponding to a particle's measurable position is not the canonical \hat{x} (which exhibits unphysical "Zitterbewegung"), but the Newton-Wigner (NW) operator, \hat{X} .³ The NW operator represents the center of energy of a wavefunction and is defined by the transformation $\hat{X} = U \mathcal{P}_+ \hat{x} \mathcal{P}_+ U^\dagger$. A key property is that its components commute, $[\hat{X}_i, \hat{X}_j] = 0$, guaranteeing that a particle's position is simultaneously localizable in all three spatial dimensions.

¹ E. Eriksen and M. Kolsrud, Nuovo Cimento Suppl. **18**, 1 (1960).

² See, e.g., C. W. Misner, K. S. Thorne, and J. A. Wheeler, "Gravitation" (W. H. Freeman, 1973), §33.4 for the classical analogue.

³ T. D. Newton and E. P. Wigner, Rev. Mod. Phys. **21**, 400 (1949).

Appendix L Relativistic Symmetries and the Gravitationally Deformed Poincaré Algebra

A fundamental test of any relativistic quantum theory is the consistency of its spacetime symmetry generators. This appendix demonstrates that the M-First framework yields a consistent, albeit non-trivially deformed, representation of the Poincaré algebra. This deformation reveals gravity's deep role in shaping a particle's fundamental invariants.

Angular Momentum and Rotational Symmetry.

We first confirm that the theory respects the spacetime's rotational symmetry. The total angular momentum operator, $\hat{J} = \hat{L} + \hat{S}$, is constructed from the orbital part $\hat{L} = \hat{x} \times \hat{\Pi}$ and the spin part $\hat{S} = (\hbar/2)\hat{\Sigma}$. A direct calculation verifies that \hat{J} satisfies the $\mathfrak{su}(2)$ Lie algebra, $[\hat{J}_i, \hat{J}_j] = i\hbar\epsilon_{ijk}\hat{J}_k$. It is also a conserved quantity, as it commutes with the Foldy-Wouthuysen (FW) Hamiltonian derived in Appendix K:

$$[\hat{J}, \hat{H}_{\text{FW}}] = 0. \quad (\text{F1})$$

This confirms that total angular momentum is conserved in the stationary, axisymmetric spacetime, as required.

Boost Generators and the Centrally Extended Algebra.

Lorentz boost generators are more subtle. Their form is fixed by the requirement that they correctly generate boosts for the physical (Newton-Wigner) position operator \hat{X} . In the FW picture, this leads to the generator:

$$\hat{K} = \frac{1}{2c^2}\{\hat{X}, \hat{H}_{\text{FW}}\} - \frac{\hbar c^2(\hat{\Sigma} \times \hat{\Pi})}{2\hat{\Lambda}(\hat{\Lambda} + m_c c^2)} - t\hat{\Pi}. \quad (\text{F2})$$

A careful evaluation of the commutator $[\hat{K}_i, \hat{K}_j]$ reveals a non-trivial deformation of the standard Poincaré algebra:

$$[\hat{K}_i, \hat{K}_j] = -i\hbar \frac{\epsilon_{ijk}}{c^2} \hat{J}_k + \frac{\hbar}{c^2} \epsilon_{ijk} \Omega_k (\hat{H}_{\text{FW}} - m_c c^2). \quad (\text{F3})$$

The first term is the standard Wigner rotation. The second is a central extension proportional to $\hbar\Omega$. This term reveals a purely quantum effect: two successive boosts fail to commute back to a simple rotation, instead acquiring an additional term dependent on the background frame-dragging. This signifies that the geometry of boosts is itself curved by the background rotation.

The Pauli-Lubanski Invariant and the Localization of Spin.

The deformation of the algebra mandates a re-evaluation of its Casimir invariants—the quantities that classify irreducible representations and thus define a particle's intrinsic properties. The primary invariant is the square of the Pauli-Lubanski pseudovector, $\hat{W}^\mu = -\frac{1}{2}\epsilon^{\mu\nu\rho\sigma}\hat{J}_{\nu\rho}\hat{P}_\sigma$, where $\hat{P}^\mu = (\hat{H}_{\text{FW}}/c, \hat{\Pi})$. A full calculation of $\hat{W}^2 = \hat{W}^\mu\hat{W}_\mu$ yields a profound result:

$$\hat{W}^2 = -m_c^2(\hat{x})c^2\hat{S}^2 = -m_0^2c^2(-g_{00}(\hat{x}))s(s+1)\hbar^2. \quad (\text{F4})$$

This result demonstrates a remarkable consequence of the deformed algebra: the Poincaré invariant that defines a particle's spin magnitude is no longer a universal constant. It becomes a local field, scaled by the gravitational potential. Gravity is thus woven into the very fabric of a particle's identity. This *localization of a Casimir invariant* is a unique and fundamental prediction of the M-First framework.

Appendix M Interactions, Currents, and Geometric Observables

This final appendix connects the M-First formalism to key physical observables. We derive expressions for the stress-energy tensor, the electromagnetic current, and the geometric phase acquired

during adiabatic evolution, revealing novel predictions for how M-First particles source and respond to external fields.

Appendix M.1 Stress-Energy Tensor and Energy Conditions

The canonical stress-energy tensor for the M-First Dirac field is obtained by varying the action with respect to the spacetime metric. Applying the Belinfante-Rosenfeld procedure to obtain a symmetric tensor, we can analyze its components in the FW picture. The operator for the energy density is given by:

$$\hat{T}^{00}(\mathbf{x}) = \hat{H}_{\text{FW}}\delta^{(3)}(\mathbf{x} - \hat{\mathbf{X}}), \quad (\text{F5})$$

where $\hat{\mathbf{X}}$ is the Newton-Wigner position operator. For any positive-energy state ($H_{\text{FW}} = \Lambda > 0$), the expectation value of the energy density is manifestly positive, $\langle T^{00} \rangle \geq 0$. The M-First framework thus satisfies the weak energy condition.

Appendix M.2 Gravitationally Modified Electromagnetic Current

A key prediction of the framework is that gravity alters how fermions couple to gauge fields. This is derived by applying a U(1) gauge principle to the M-First Dirac action. The conserved Noether current associated with this symmetry is found to be:

$$\hat{J}^\mu = qc\bar{\psi}\gamma^\mu\left(1 + \frac{\Phi_g(\hat{\mathbf{x}})}{m_0^2c^2}\right)\psi + \mathcal{O}(\Phi_g^2), \quad (\text{F6})$$

where $\Phi_g = M_g^2 - M_{\text{flat}}^2$ is the M-First gravitational modifier. This result implies that a particle's effective electric charge is screened or anti-screened by the local gravitational potential. This non-universal coupling is a distinctive and testable prediction of the framework, differing from standard minimally coupled theories.

Appendix M.3 Geometric Phase from Adiabatic Evolution

When the background fields (g_{00}, Ω) vary slowly in time, a quantum state acquires a geometric (Berry) phase. This phase is generated by the Berry curvature, a 2-form in the parameter space of the background fields. A formal calculation shows that the components of this curvature are directly determined by the same off-diagonal Hamiltonian matrix elements that drive the spin precession dynamics derived in Eq. (E4) of Appendix K.

This provides a deep link between spin dynamics and geometry: the forces causing spin precession are the agents that curve the particle's state space, leading to the accumulation of a geometric phase. For the gravitational spin-Hall effect, we can make an order-of-magnitude estimate: a 10 GeV electron in a region with a gravitational gradient equivalent to a 1 T/m magnetic field gradient would experience a spin precession of roughly 10^{-19} rad/s. While extremely small, this points towards a new class of high-precision experiments that could probe such effects. A full treatment of these couplings at the loop level is the subject of a forthcoming paper on M-First Quantum Field Theory [8].

References

1. Birrell, N.D.; Davies, P.C.W. *Quantum Fields in Curved Space*; Cambridge University Press, 1982.
2. Wald, R.M. *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*; University of Chicago Press, 1994.
3. Klavness, A. Momentum Is All You Need. OSF Preprint, 2025. https://doi.org/10.31219/osf.io/jdr3q_v2.
4. Klavness, A. Enhanced Pycnonuclear Reactions in Neutron Star Crusts from Momentum-First Gravity. (unpublished) Brief Communication, manuscript in preparation.
5. Brown, E.F.; Cumming, A. Mapping the neutron star interior: Constraints from the crustal cooling time scale. *Astrophys. J.* **2009**, *698*, 1020, [arXiv:0901.1092]. <https://doi.org/10.1088/0004-637X/698/2/1020>.
6. Klavness, A. A First Principles Derivation of the Earth Fly-by Velocity Shift. (unpublished) Brief Communication, manuscript in preparation.

7. Anderson, J.D.; Campbell, J.K.; Ekelund, J.E.; Ellis, J.; Jordan, J.F. Anomalous Orbital-Energy Changes Observed during Spacecraft Flybys of Earth. *Phys. Rev. Lett.* **2008**, *100*, 091102. <https://doi.org/10.1103/PhysRevLett.100.091102>.
8. Klaveness, A. Momentum-First Quantum Field Theory: Emergent Interactions from Internal Momentum Structure. (unpublished) Manuscript in preparation.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.