

Article

Not peer-reviewed version

Unifying Penrose Process and Blandford–Znajek Mechanism using Negative Phase Velocity

[Sandi Setiawan](#)*

Posted Date: 30 October 2025

doi: 10.20944/preprints202510.2341.v1

Keywords: Negative phase velocity (NPV); Penrose process; Blandford–Znajek mechanism; Superradiance; Kerr black hole / ergosphere; Negative Killing energy; Energy extraction from black holes; Poynting flux / electromagnetic jets; Frame-dragging / horizon physics; GRMHD diagnostics / numerical simulations; Wave amplification / mode scattering



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Unifying Penrose Process and Blandford–Znajek Mechanism Using Negative Phase Velocity

Sandi Setiawan

Life Member, Clare Hall, University of Cambridge, UK; sandi.setiawan@cantab.net

Abstract

The extraction of rotational energy from a Kerr black hole admits several celebrated descriptions, including the Penrose process, wave superradiance, and the Blandford–Znajek mechanism. Traditionally treated as distinct, these processes are shown here to be unified through a single criterion: the existence of negative Killing energy flux across the horizon, locally manifested as negative phase velocity (NPV) in the ergosphere. I demonstrate that (i) the Penrose process corresponds to particle trajectories with $E_\chi < 0$, (ii) superradiance corresponds to wave modes with $\omega < m\Omega_H$, and (iii) the Blandford–Znajek mechanism corresponds to electromagnetic field lines with $\Omega_F < \Omega_H$. In all cases, the horizon absorbs negative energy while positive energy escapes to infinity. By expressing these mechanisms through the NPV condition $\mathbf{S} \cdot \mathbf{k} < 0$, we present a unified theoretical framework for black hole energy extraction.

Keywords: negative phase velocity (NPV); Penrose process; Blandford–Znajek mechanism; superradiance; Kerr black hole / ergosphere; negative Killing energy; energy extraction from black holes; Poynting flux / electromagnetic jets; frame-dragging / horizon physics; GRMHD diagnostics / numerical simulations; wave amplification / mode scattering

1. Introduction

Rotating black holes provide one of the most remarkable natural laboratories for relativistic energy transfer. The ergosphere—the spacetime region outside the horizon where the timelike Killing vector becomes spacelike—permits negative-energy states relative to infinity. This peculiar feature allows rotational energy to be tapped, an idea pioneered by Penrose (1969).

Over the decades, three paradigms have emerged:

- **Penrose process:** a test particle entering the ergosphere can split into fragments, one of which falls into the black hole with negative Killing energy, allowing the other to escape with enhanced energy.
- **Superradiance:** classical or quantum fields scattered by a rotating black hole can emerge amplified if their frequency satisfies $\omega < m\Omega_H$.
- **Blandford–Znajek mechanism:** magnetic field lines threading the horizon, coupled to a force-free magnetosphere, can extract rotational energy and drive astrophysical jets.

Although derived in different contexts—particles, waves, and electromagnetic fields—these processes share the same root: *the absorption of negative energy by the horizon*. This work demonstrates that a local diagnostic, *negative phase velocity* (NPV), provides a unifying description. In all channels, energy extraction occurs precisely when the phase velocity points opposite to the energy flux, i.e., $\mathbf{S} \cdot \mathbf{k} < 0$.

Table 1. Connection between Penrose process, superradiance, and Blandford–Znajek (BZ) mechanism, and their relation to negative phase velocity (NPV).

Phenomenon	Description	Relation to NPV
Penrose process	Particle in the ergosphere splits; one fragment with negative energy falls into the hole, the other escapes with greater-than-incoming energy.	Negative-energy state corresponds to energy flow opposing momentum (phase direction) — a mechanical analog of NPV.
Superradiance	Incident waves scattering off a rotating black hole are amplified when the superradiant condition holds.	Amplification arises because, within the ergosphere, local phase and energy flow oppose each other, creating an NPV region that extracts rotational energy.
Blandford–Znajek (BZ)	Magnetic field lines threading the rotating hole drive currents; energy is extracted as outward Poynting flux.	Magnetosphere behaves as a macroscopic NPV-like system: outward Poynting flux with co-rotating field structure; horizon acts like a rotating conductor with effectively negative impedance.

2. Theoretical Framework

2.1. Geometry and Conserved Currents

Consider a Kerr black hole with mass M and angular momentum $J = aM$. The horizon generator is

$$\chi^\mu = t^\mu + \Omega_H \phi^\mu, \quad \Omega_H = \frac{a}{2Mr_+}, \quad (1)$$

with r_+ the horizon radius.

For a stress-energy tensor $T_{\mu\nu}$, the conserved flux of energy across the horizon is

$$\dot{E}_H = \int_{\mathcal{H}} T_{\mu\nu} \chi^\mu d\Sigma^\nu, \quad (2)$$

while the angular momentum flux is

$$\dot{J}_H = - \int_{\mathcal{H}} T_{\mu\nu} \phi^\mu d\Sigma^\nu. \quad (3)$$

Horizon regularity imposes the relation

$$\dot{E}_H = \Omega_H \dot{J}_H. \quad (4)$$

Thus, if $\dot{J}_H < 0$, the horizon absorbs negative energy: rotational energy is extracted.

2.2. Negative Phase Velocity Condition

In a locally non-rotating (ZAMO) frame, the Poynting vector \mathbf{S} gives the direction of energy flow, while the wavevector \mathbf{k} sets the direction of phase propagation. The *NPV condition* is

$$\mathbf{S} \cdot \mathbf{k} < 0, \quad (5)$$

which coincides with the condition that the horizon absorbs negative Killing energy.

In ordinary media, the phase velocity (direction of wavefront propagation) and the energy flow (given by the Poynting vector) point in the same direction. In negative-phase-velocity (NPV) or left-handed media, the phase velocity points opposite to the energy flow. Such media exhibit negative refraction, reverse Doppler, and reverse Cherenkov effects. This can also arise not just in meta-materials, but in curved spacetime near rotating black holes, where the spacetime metric itself acts as an effective medium with exotic constitutive parameters.

In Kerr geometry (i.e. rotating black holes), the frame-dragging of spacetime inside the ergosphere produces regions where the effective electromagnetic constitutive tensor allows NPV propagation. This means that Waves can have phase velocity directed inward while energy flow (Poynting vector)

points outward, and the local observer sees “backward” wave propagation relative to the global energy flux. This is the electromagnetic analog of negative energy orbits in the Penrose process.

3. Results: Three Channels of Energy Extraction

3.1. Penrose Process (Particle Channel)

A particle with 4-momentum p^μ has Killing energy

$$E_\chi = -p_\mu \chi^\mu. \quad (6)$$

In the ergosphere, E_χ can become negative. If a particle splits such that one fragment has $E_\chi < 0$, the escaping fragment must carry away energy in excess of the original. This is the particle realization of the NPV condition: momentum direction anti-aligned with energy flux.

3.2. Superradiance (Wave Channel)

For a scalar field mode

$$\Phi \sim e^{-i\omega t + im\phi}, \quad (7)$$

the horizon flux evaluates to

$$\dot{E}_H \propto (\omega - m\Omega_H) |\Phi|^2. \quad (8)$$

Amplification occurs when

$$\omega < m\Omega_H, \quad (9)$$

so that $\dot{E}_H < 0$. Locally, this corresponds to NPV: the phase velocity associated with \mathbf{k} is opposite to the outward energy flux \mathbf{S} .

3.3. Blandford–Znajek Mechanism (Electromagnetic Channel)

In a magnetosphere, magnetic field lines rotate with angular velocity Ω_F . Each azimuthal Fourier mode behaves as if $\omega = m\Omega_F$. The horizon energy flux is then

$$\dot{E}_H \propto (\Omega_F - \Omega_H) \dot{J}_H. \quad (10)$$

When $\Omega_F < \Omega_H$, the flux is negative and outward Poynting power is produced. The classic BZ power law is

$$P_{\text{BZ}} \sim \kappa \Omega_H^2 \Phi_B^2 \left(\frac{\Omega_F}{\Omega_H} \right) \left(1 - \frac{\Omega_F}{\Omega_H} \right), \quad (11)$$

with Φ_B the magnetic flux and κ a dimensionless constant.

Table 2. Unified physical picture across particle, wave, and field/plasma regimes, highlighting how NPV manifests in each.

Process	Mechanism	NPV manifestation
Penrose (particle)	Negative-energy orbits in the ergosphere enable net energy extraction.	Momentum/phase direction vs. energy flow are misaligned (mechanical analog of NPV).
Superradiance (wave)	Wave amplification by a rotating horizon under the superradiant condition.	Phase velocity opposes energy (Poynting) flow within the ergosphere (wave-level NPV).
Blandford–Znajek (field/plasma)	Magnetically driven jet (outward Poynting flux) powered by black-hole spin.	Effective negative electromagnetic impedance; macroscopic NPV-like energy extraction.
NPV (EM/geometric)	Phase–energy opposition enabled by frame-dragging and constitutive effects.	Underlies all of the above energy-extraction phenomena.

4. Discussion

The above results demonstrate that all known energy extraction processes from a Kerr black hole are governed by a single criterion: *the absorption of negative Killing energy*, equivalently expressed as *negative phase velocity propagation* in the ergosphere.

- **Penrose process:** NPV arises in the momentum-energy misalignment of particles.
- **Superradiance:** NPV is realized when the reflected wave has group velocity outward but phase velocity inward.
- **BZ mechanism:** NPV manifests as field lines with pattern speed below the horizon angular velocity, leading to reversed energy flux across the horizon.

This unification has several consequences:

1. It clarifies that BZ is not fundamentally different from superradiance but its *steady, force-free, large-scale limit*.
2. It provides a diagnostic for simulations: identifying regions with $\mathbf{S} \cdot \mathbf{k} < 0$ can signal ongoing energy extraction in numerical GRMHD models.
3. It suggests that analogous processes may occur in other rotating systems with ergoregions, such as analogue gravity experiments or rotating compact stars.




Figure 1. Penrose Process vs. Blandford-Znajek Mechanism

5. Conclusion

We have demonstrated that the Penrose process, superradiant scattering, and the Blandford-Znajek mechanism are three manifestations of a unified energy-extraction phenomenon from rotating black holes. The ergosphere permits negative Killing energy states, whose local signature is *negative phase velocity*.

- In the particle channel, this condition allows the Penrose process.
- In the wave channel, it yields superradiant amplification.
- In the electromagnetic channel, it drives the Blandford-Znajek jet.

This unified perspective highlights the ergosphere as the essential stage and NPV as the diagnostic connecting microscopic, mesoscopic, and macroscopic energy extraction processes.

Data Availability Statement: No data were generated or analyzed in support of this research.

References

1. Blandford, R. D., & Znajek, R. L., 1977, MNRAS 179, 433.
2. Misner, C. W., 1972, Phys. Rev. Lett. 28, 994.
3. Penrose, R., 1969, Rivista del Nuo. Cim. 1, 252.
4. Setiawan, S., Mackay, T. G., Lakhtakia, A, 2005, Phys.Lett. A341 15-21; Erratum-ibid. 2007, A361, 529-533.
5. Teukolsky, S. A., & Press, W. H., 1974, ApJ 193, 443.
6. Thorne, K. S., Price, R. H., & Macdonald, D. A., 1986, Black Holes: The Membrane Paradigm, Yale University Press.
7. Zel'dovich, Y. B., 1971, JETP Lett. 14, 1.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.