

Genetic Variants in Potassium Channel Genes and Their Clinical Implications in Kazakhstani Patients with Cardiac Arrhythmias

[Ayaulym Chamoieva](#) , [Saule Rakhimova](#) * , Zhannur Abilova , [Ainur Akhmetova](#) , Gulbanu Akilzhanova , [Madina Zhalbinova](#) , [Asset Danjyarov](#) , [Kenes Akilzhanov](#) , Askhat Molkenov , [Ulykbek Kairov](#) , Anargul Kuanysheva , Nurlan Shaimardanov , [Ayan Abdrakhmanov](#) , Makhabbat Bekbossynova , [Ainur Akilzhanova](#) *

Posted Date: 26 November 2025

doi: 10.20944/preprints202511.2094.v1

Keywords: cardiac arrhythmia; potassium channel genes; *KCNH2*; genetic testing; next-generation sequencing; sudden cardiac death; clinical genetics; Kazakhstan



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Genetic Variants in Potassium Channel Genes and Their Clinical Implications in Kazakhstani Patients with Cardiac Arrhythmias

Ayaulym Chamoieva ¹, Saule Rakhimova ^{1,2,*}, Zhannur Abilova ¹, Ainur Akhmetova ¹, Gulbanu Akilzhanova ³, Madina Zhalbinova ¹, Asset Daniyarov ^{1,4}, Kenes Akilzhanov ³, Askhat Molkenov ¹, Ulykbek Kairov ^{1,2}, Anargul Kuanysheva ⁵, Nurlan Shaimardanov ⁶, Ayan Abdrakhmanov ⁷, Makhabbat Bekbossynova ⁸ and Ainur Akilzhanova ^{1,2,4,7,9,*}

¹ Center for Life Sciences, National Laboratory Astana, Nazarbayev University, Astana 010000, Kazakhstan

² Eurasian Society of Personalized Medicine, Astana, 010000, Kazakhstan

³ Department of Medicine, Semey Medical University, Pavlodar Branch, Pavlodar 140000, Kazakhstan

⁴ Department of General Biology and Genomics, L. N. Gumilyov Eurasian National University, Astana, 010000, Kazakhstan

⁵ Medical Centre Hospital of President's Affairs Administration of the Republic of Kazakhstan, Astana, 010000, Kazakhstan

⁶ Department of Medicine, Semey Medical University, Semey 071400, Kazakhstan

⁷ Heart Rhythm Institute, Medical University Astana, Astana, 010000, Kazakhstan

⁸ Heart Center, Corporate fund "University Medical Center", Nazarbayev University, Astana 010000, Kazakhstan

⁹ The Academy of Public Administration under the President of the Republic of Kazakhstan, Astana, Republic of Kazakhstan

* Correspondence: akilzhanova@nu.edu.kz; akilzhanainur@gmail.com; Tel.: +7-7172-706501; saule.rakhimova@nu.edu.kz.

Abstract

Background/Objectives: Cardiac arrhythmias are among the leading causes of sudden cardiac death (SCD). Pathogenic variants in potassium channel genes play a key role in inherited arrhythmia syndromes, yet their contribution in Central Asian populations remains poorly characterized. **Methods:** We performed targeted next-generation sequencing (NGS) using a 96-gene custom Haloplex panel in 79 Kazakhstani patients with clinically diagnosed arrhythmias, including atrioventricular block, sick sinus syndrome, and atrial fibrillation. Detected variants in potassium channel genes were classified according to ACMG guidelines and correlated with clinical phenotypes. **Results:** A total of 52 variants were identified across 11 potassium channel genes. Two likely pathogenic variants (*KCNH2* p.Cys66Gly and p.Arg176Trp) and six variants of uncertain significance (VUS) in *KCNQ1*, *KCNE2*, *KCNE3*, and *KCNJ8* were detected. Two novel previously unreported variants were found in *KCNE5* and *KCND3*. Patients harboring pathogenic variants commonly presented with early-onset arrhythmias or a positive family history of cardiovascular disease. Carriers of *KCNH2* variants exhibited mild QT prolongation and recurrent syncope. **Conclusions:** This is the first genetic study of potassium channel gene mutations in Kazakhstani patients with cardiac arrhythmias. Detection of pathogenic and novel variants highlights the clinical utility of integrating genetic testing into diagnostic and management pathways for arrhythmia syndromes. Population-specific genomic data are essential for improving risk stratification, guiding medication safety, and enabling cascade family screening in Central Asia.

Keywords: cardiac arrhythmia; potassium channel genes; *KCNH2*; genetic testing; next-generation sequencing; sudden cardiac death; clinical genetics; Kazakhstan

1. Introduction

Cardiovascular diseases (CVDs) remain the leading cause of mortality globally, accounting for approximately 18 million deaths in 2019 [1], with low- and middle-income countries such as Kazakhstan disproportionately affected. Cardiac arrhythmias are a major cause of morbidity and mortality worldwide, contributing significantly to sudden cardiac death (SCD) and heart failure. Cardiac arrhythmias represent a clinically heterogeneous group of disorders diagnosed based on an electrocardiographic pattern demonstrating depolarization and repolarization of the atria and ventricles. Despite advances in diagnostic and therapeutic strategies, a substantial proportion of arrhythmias remain idiopathic, with unclear underlying mechanisms [2-3]. Increasing evidence demonstrates that genetic variants, particularly in ion channel genes, are key determinants of both inherited and acquired arrhythmia susceptibility. Hereditary arrhythmia syndromes include primary electrical heart disorders known as channelopathies. Channelopathies, the dysfunction of ion channels, are caused by mutations in genes coding pore-forming α -subunit of ion channels [4]. Patients with channelopathies are usually unaware of their disorders and have a substantial risk of SCD. Therefore, early genetic diagnostics should not be neglected [5]. Cardiac channelopathies include a group of conditions as syncope, ventricular arrhythmias and even SCD [6].

Among these, potassium channel genes play a pivotal role in regulating cardiac action potential repolarization. Notably, nearly 80 genes encode pore-forming subunits of potassium channels, representing the largest family of pore-forming channel proteins in the human genome. Pathogenic variants in *KCNQ1*, *KCNH2*, *KCNE1*, and related genes are established causes of long QT syndrome (LQTS), short QT syndrome (SQTS), Brugada syndrome (BrS), and atrial fibrillation (AF) [3-4, 7]. Moreover, mutations in potassium channel genes can alter cardiac repolarization by disrupting normal ion flow across the cell membrane. Typically, loss-of-function mutations lead to delayed repolarization and are associated with LQTS, while gain-of-function mutations may shorten repolarization, contributing to AF and SQTS. Heterozygous missense variants are the most frequent and often disturb channel function through mechanisms such as protein misfolding, defective trafficking, or abnormal gating. Identification of above-mentioned variants not only aids in confirming diagnosis but also informs risk stratification, pharmacogenomic guidance, and family screening [8-10].

To date, most genetic studies of arrhythmia syndromes have focused on European, East Asian, or North American populations. Data from Central Asia, including Kazakhstan, are scarce. Given the unique ethnic and genetic background of the Kazakh population, characterizing the local variant spectrum is critical for accurate interpretation and clinical translation of genomic findings. The development of next-generation sequencing (NGS) has become a valuable tool for uncovering disease-causing variants, particularly when targeting specific genomic regions [11].

This study aimed to identify pathogenic and novel variants in potassium channel genes among Kazakhstani patients with clinically diagnosed cardiac arrhythmias using targeted NGS. We further sought to correlate genetic findings with clinical phenotypes and discuss implications for personalized arrhythmia management.

2. Materials and Methods

2.1. Study Population

The study was performed with the principles of Declaration of Helsinki. The research protocol was approved by the Ethics Committees of the National Laboratory Astana, Nazarbayev University (№ 16, from 11 March 2015) and the National Research Cardiac Surgery Center, NRCC, (since 2023-Heart Center, Corporate Fund "University Medical Center", Nazarbayev University), Astana, Kazakhstan (№ 16, from 24 February 2015). Written consent was obtained from all participants of our study. To protect the rights of patients and do not disclose their personal information patient's personal information was encoded and all the data was depersonalized during database collection.

A total of 79 unrelated patients with arrhythmic disorders were included in our study group, including atrioventricular (AV) block, sick sinus syndrome (SSS), and atrial fibrillation (AF). Among AF cases, both idiopathic atrial fibrillation (iAF) and coronary heart disease-associated atrial fibrillation (CHD-AF) were represented. The clinical diagnosis of arrhythmic disorders was based on the medical history of patients, clinical representation, all instrumental findings and made by

clinicians. All study participants were recruited during 2015-2016 during their treatment at the NRCC. Demographics (age, gender, and ethnicity), family anamnesis and clinical features, results of the blood tests, electrocardiogram, 24-hour Holter electrocardiogram monitoring and echocardiogram, were obtained from medical records (Supplementary Table S1). Moreover, 234 healthy Kazakh individuals from the population database of whole-genome sequencing were used as a control.

2.2. DNA Extraction

Peripheral blood samples were collected, and genomic DNA was extracted from whole blood samples using the Wizard® Genomic DNA Purification kit (Promega, Madison, WI, USA) according to the manufacturer's instructions. The quality (concentration and purity) of extracted DNA was measured by NanoDrop™ Spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). The qualitative and quantitative analyses of samples were undertaken using spectrophotometer Qubit 2.0 (Thermo Fisher Scientific, Waltham, MA, USA) and gel electrophoresis (2% agarose gel). The total volume of isolated DNA was 45 µl, the concentration of DNA in 1 microliter was at least 5 ng/µl. Enrichment Control DNA was used as a control.

2.3. Targeted Sequencing

Targeted sequencing was performed using the Agilent SureSelect/ Haloplex custom panel encompassing 96 genes associated with cardiac arrhythmias, including 11 potassium channel genes (Table 1).

Table 1. Selected potassium channel genes associated with arrhythmia.

Gene	Protein	Chromosome	Transcript	OMIM	Phenotype
KCNA5	Kv1.5	12p13.32	NM_002234	176267	AF
KCND3	Kv4.3	1p13.2	NM_172198	605411	BrS
KCNE1	Kv7.1	21q22.12	NM_000219	176261	LQTS, Jervell and Lange Nielsen syndrome
KCNE2	Kv7.2	21q22.11	NM_172201	603796	LQTS, AF
KCNE3	Kv7.3	11q13.4	NM_005472	604433	BrS
KCNE5	Kv7.5	Xq23	NM_012282	600681	BrS
KCNH2	Kv11.1/hERG	7q36.1	NM_172057/ NM_000238	152427	LQTS, SQTS
KCNJ2	Kir2.1	17q24.3	NM_000891	600681	Andersen-Tawil syndrome, SQTS, AF
KCNJ5	Kir 3.4	11q24.3	NM_000890	600734	LQTS
KCNJ8	Kir 6.1	12p12.1	NM_00498	600935	BrS, SQTS, IVF associated, ERS
KCNQ1	Kv7.1	11p15.5-p15.4	NM_181798	607542	LQTS, SQTS, AF, Jervell and Lange Nielsen syndrome

*AF, atrial fibrillation; BrS, Brugada syndrome; LQTS, long QT syndrome; SQTS, short QT syndrome; IVF, idiopathic ventricular fibrillation; ERS, early repolarization syndrome.

DNA libraries for all individuals in our group were prepared using the HaloPlex Custom Panel Tier 1 kit (Agilent Technologies) according to the protocol "HaloPlex Target Enrichment System for Illumina Sequencing" (v.D5, May 2013). The protocol is available online (<https://www.agilent.com>). The Haloplex protocol is optimized for digestion of 225 ng of genomic DNA. Prepared DNA-libraries were checked on 2100 BioAnalyzer (Agilent Technologies, Santa Clara, CA, US) for quality using the High Sensitivity DNA Assay kit (Thermo Fisher Scientific). Then, pooled samples were loaded for sequencing on Illumina HiSeq2000 platform (San Diego, USA). The design was adapted to the paired-

end 150 bp sequencing technology. The size of target region made up 463.767 kbp. Concerning the realization of the NGS technology and the accuracy of sequences, the mean target coverage was 99.46 %. We considered that it was correctly covered and then suitable for further data analysis.

2.4. Bioinformatics and Variant Annotation

Raw data were processed using standard bioinformatics pipelines (BWA for alignment, GATK for variant calling). Genetic variants were analyzed using the online software SureCall Design (Agilent Technologies, Santa Clara, CA, USA). The obtained variants corresponded to the Human Gene Mutation Database (HGMD) and were annotated with ANNOVAR. The clinical significance of the variants was interpreted in accordance with the guidelines developed by the American College of Medical Genetics and Genomics (ACMG) and the Association of Molecular Pathology (AMP) in 2015 and classified as pathogenic (P), likely pathogenic (LP), a variant of uncertain significance (VUS), likely benign (LB), or benign (B) [12]. Additionally, the obtained variants were checked in online databases of Clinvar (<http://www.ncbi.nlm.nih.gov/clinvar>) and Franklin Genoox (<https://franklin.genoox.com/>). Population frequencies were verified against international research databases, the 1000 Genomes (1000G), the Genome Aggregation Database (gnomAD), Exome Aggregation Consortium (ExAC), Online Mendelian Inheritance in Man (OMIM), Mutation Taster, NCBI, etc. Additionally, allele frequencies of variants were compared with a cohort of 234 healthy Kazakh individuals.

All sequence data presented in this study are deposited in the NCBI Sequence Read Archive (SRA) repository and are publicly available under accession BioProject number PRJNA908657 (<https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA908657>).

2.5. Statistical Analysis

Statistical analysis was performed using IBM SPSS Statistics 23.0 program (SPSS, Chicago, IL, USA). Continuous variables were presented as mean \pm standard deviation, and categorical variables as numbers (n) or percentages (%).

3. Results

3.1. Clinical Characteristics

The cohort included 79 patients. The most common diagnoses were atrial fibrillation (59.5%), AV block (20%), and sick sinus syndrome (20%). The clinical characteristics of patients are provided in Table 2.

Table 2. Patient characteristics.

Characteristics	Total, n = 79	AV block, n=16	SSS, n=16	iAF, n=31	CHD AF, n=16
Age, years	47.5 \pm 17.5	45,6 \pm 23.8	47,7 \pm 17.6	42,7 \pm 13.9	58,4 \pm 11.9
Gender, M/F	52/27	6/10	9/7	22/9	15/1
BMI, kg/m ²	26.8 \pm 5.4	26,1 \pm 6	25,5 \pm 5.4	27,3 \pm 5.3	28,2 \pm 5.4
Family history CVD, N	23 (29%)	3	4	8	8
Diabetics, N	11 (14%)	2	1	5	3
Syncope, N	12 (15%)	7	4	0	1
Pacemaker implantation, N	31 (39%)	14	16	0	1
Prolonged QT, N	14 (18%)	5	6	0	3

The mean age at onset in the study group was 47.5 \pm 17.5 years; 5 patients (6%) were under 18 years of age. The cohort included 52 (65.8%) male patients. The research group consists of Kazakhs (64.2%), Russians (17.3%) and other nationalities (18.5%). Besides, 14 (18%) of the patients had a

prolonged QT interval (QT \geq 440 ms), and 12 (15%) reported experiencing syncope in their medical history. Of 79 patients with arrhythmias, 23 individuals (29%) had a predisposition to cardiovascular diseases in their family history. Due to serious cardiac risks accompanying life-threatening arrhythmias, 31 (39%) patients underwent pacemaker implantation, namely almost all of the AV block and SSS patients. The average body mass index (BMI) of the patients was 26.8 ± 5.4 kg/m², and 11 (14%) patients had type 2 diabetes. In addition, 9 patients with CHD AF had suffered a myocardial infarction. None of the patients had a clinical death (cardiac arrest).

3.2. Genetic Findings

A total of 52 variants were identified across 11 potassium channel genes; full details are provided in Supplementary Table S2. According to ACMG classification [12], these included 2 likely pathogenic variants in *KCNH2* (p.Cys66Gly, p.Arg176Trp), 6 VUS in *KCNQ1*, *KCNE2*, *KCNE3*, and *KCNJ8* (Table 3).

Table 3. List of patients with rhythm disorders carrying rare disease-causing variants.

Case ID	Sex	Age	Group	Clinical phenotype	LVE F (%)	QT interval, ms	Familial history	Gene	Nucleotide	AA change	gnomeAD	Exonic effect	HGM D	ACMG score
#516	M	55	CHD AF	Myocardial infarction	44	378	yes	KCNH2	NM_00238.3:c.196T>A	p.Cys66Gly	-	missense	DM	4
#334	F	20	iAF	WPW syndrome, Paroxysmal orthodromic AV-reentry tachycardia	61	402	no	KCNH2	NM_00238.3:c.196T>A	p.Cys66Gly	-	missense	DM	4
#377	M	24	iAF	A blood thrombus in the auricle of the left atrium	64,71	320	no	KCNH2	NM_00238.3:c.196T>A	p.Cys66Gly	-	missense	DM	4
#202	M	76	iAF	CHF I, Paroxysmal AF	64,71	340	no	KCNH2	NM_00238.3:c.526C>T	p.Arg176Trp	0.000618	missense	DM	4
#573	M	41	iAF	EHRA II, CHF, NYHA I, Paroxysmal AF	52	392	no	KCNH2		p.Arg176Trp	0.000618	missense	DM	4
#464	F	30	iAF	Paroxysmal AF, EHRA I. Left atrial flutter, Atrial extrasystole	58	380	no	KCNE2	NM_172201.1:c.29C>A	p.Thr10Lys	0.000968	missense	DM?	3
#80	M	57	SSS	Arterial hypertension	60	No path	yes	KCNE3		-	0.000646	missense	DM	3

#282	M	22	iAF	Paroxysmal AF	-	400	no	KCNQ1	NM_002182:c.1128+4C>T	-	0.0001	intron	N/A	3
#93	F	47	AV block	Arterial hypertension	66	No path	yes	KCNQ1	NM_002182:c.1033-4C>T	-		intron	N/A	3
#333	M	30	iAF	CHF NYHA II, persistent AF	46	360	yes	KCNJ8	NM_004982.3:c.263C>G	p.Ala88Gly	-	missense	N/A	3
#579	M	64	CHD AF	CHF NYHA II, EHRA I. Persistent AF	36	386	yes	KCNJ8	NM_004982.3:c.1145A>G	p.Lys382Arg	-	missense	N/A	3

*LVEF, left ventricular ejection fraction; AA, amino acid; CHF, congestive heart failure; DM, disease-causing mutation;

The variants can be considered very rare with a minor allele frequency (MAF) threshold of $\leq 0.02\%$ in all used population databases (gnomeAD, ExAC, dbSNP).

Categorization of all potassium channel genes according to ACMG classification indicated that the number of mutations in the iAF study group is higher than in other groups, as well as the number of patients. AV block, SSS, and CHD AF patients share almost the same number of genetic variants in potassium channel genes.

Notably, a prolonged QT interval (>440 ms) was not observed in any patient carrying disease-causing variants.

Overall, 17 variants in potassium channel genes were presented in the HGMD database (accessed August 2025, Table 4).

Table 4. Potassium channel gene variants in HGMD.

Gene	HGMD mutation	Variant class	dbSNP identifier	gnomeAD	ClinVar
KCNH2	Tyr652Ter	DM	rs1137617	-	Not provided
KCNQ1	Tyr662Ter	DM	rs1161907	0.000649	Conflicting interpretations of pathogenicity
KCNH2	Cys66Gly	DM	rs199473416	-	Not provided
KCNE3	Thr4Ala	DM	rs200856070	0.000646	Conflicting interpretations of pathogenicity
KCNQ1	Lys393Asn	DM	rs12720457	0.000387	Conflicting interpretations of pathogenicity
KCNH2	Arg176Trp	DM	Rs36210422	0.000618	Conflicting interpretations of pathogenicity
KCNE2	Thr10Met	DM?	rs199473648	0.000968	Conflicting interpretations of pathogenicity
KCNE5	Tyr81His	DM?	Rs199924386	0.000324	Benign
KCNQ1	Pro448Arg	DM?	Rs12720449	0.005302	Benign/Likely benign
KCNA5	Pro307Ser	DFP	Rs17215409	0.002235	Conflicting interpretations of pathogenicity
KCNH2	Arg1047Leu	DFP	Rs36210421	0.026377	Conflicting interpretations of pathogenicity
KCNH2	Lys897Thr	DFP	Rs1805123	0.185003	Benign
KCNE1	Ser38Gly	DFP	rs1805127	0.659157	Benign
KCNJ2	Leu382Leu	DF	Rs173135	0.118301	Benign
KCNE3	Phe66Phe	DF	rs2270676	0.138846	Benign
KCNJ5	Glu282Gln	FP	rs7102584	0.015136	Benign

*DM, disease-causing mutation; DFP, disease-causing mutation with functional proof; DF, disease-associated polymorphism with functional evidence; FP, in vitro/laboratory or in vivo functional polymorphism.

Among them, 7 variants had previously been reported as disease-causing (DM) in the HGMD, and 2 additional variants (KCNE2 p.Thr10Met and KCNE5 p.Tyr81His) were labeled as probable disease mutations (DM?), indicating uncertain but possible pathogenicity. Two variants (KCNA5 p.Pro307Ser and KCNH2 p.Arg1047Leu) were categorized as disease-causing with functional proof (DFP). Furthermore, 5 variants were listed as functionally relevant polymorphisms (DF/FP), yet were not considered clinically pathogenic, including KCNH2 p.Lys897Thr and KCNE1 p.Ser38Gly, which showed relatively high allele frequencies (>0.01) in gnomAD and were marked as benign in ClinVar.

In this study, we identified 2 variants in KCND3, KCNE5 that were not present in dbSNP, gnomAD, ClinVar, and HGMD databases, suggesting they may represent novel or extremely rare variants (Table 5).

Table 5. Novel (previously unreported) variants in potassium-channel genes.

Case ID	Gene	Transcript	Nucleotide	Protein change	Geno type	Chr	Exonic effect	SIFT score	PolyPhe n-2 score	CADD phred
472	KCND3	NM_17219	c.1928C>A	p.Pro643His	0/1	1	missense tolerated	0.8741,	0.961,	22.5,

							damagin deleterio	g	us
							0.138,		
							possibly		
							0.7802,	damagin	8.025,
513	KCNE5	2	c.378G>T	s	0/1	X	missense tolerated	g	tolerated

*SIFT, Sorting Intolerant From Tolerant; CADD, Combined Annotation Dependent Depletion.

The frequency filtering was performed by comparing two unreported variants against a control cohort of 234 healthy Kazakh individuals to assess population-specific novelty. As a result, these variants were not observed in the control group. Also, the variants were evaluated in silico prediction tools. KCND3:p.Pro643His showed high CADD phred score (≥ 20), consistent with potential deleteriousness. KCNE5 p.Gln126His yielded mixed or tolerated predictions across SIFT, PolyPhen-2 and CADD phred.

4. Discussion

We evaluated the contribution of genetic variants in potassium channel genes in a cohort of 79 patients with cardiac arrhythmias/ In this study. Particularly, we used a targeted NGS panel consisting of 96 genes relevant to arrhythmias and cardiomyopathies to overcome the limitations of unselective sequencing. Despite the broader scope of whole-genome/whole-exome sequencing, a targeted gene enrichment approach provides full coverage, high accuracy and high sensitivity for specific genes [13].

In our study, the most commonly noted LQTS-causing variants were found in the KCNH2 and KCNQ1 genes, which encode the α subunits of potassium channels Kv11.1/hERG and Kv7.1/KvLQT1, respectively. Nearly 75% of LQTS disorder is caused by the main LQTS genes - KCNQ1, KCNH2, and SCN5A [14-15]. The first symptoms of LQTS patients are syncope and then cardiac arrest. Those symptoms are the result of ventricular arrhythmias [16]. Around 30–35% patients with LQTS have mutations in KCNQ1, while 25–40% have mutations in KCNH2, highlighting the clinical significance of potassium channels and their inclusion in diagnostic panels [17]. Namely, we observed 13 mutated alleles of KCNH2 gene, and 8 variants representing KCNQ1 gene. These genes are responsible for LQTS type 1 and LQTS type 2, respectively.

Our study demonstrates that genetic variants in potassium channel genes are detectable in a significant proportion of Kazakhstani patients with arrhythmias, even in the absence of overt QT prolongation or structural heart disease. This underscores the value of genetic testing as an adjunct to clinical evaluation, particularly in idiopathic arrhythmias or early-onset AF. Approximately 18% of patients in our cohort exhibited prolonged QT, and 15% reported syncope – both key risk markers for SCD. Normal QT is considered ≤ 440 ms for men and ≤ 460 ms for women. QT $> 450/470$ ms indicates prolongation; ≥ 480 ms is used diagnostically in symptomatic individuals, and ≥ 500 ms confers high risk [9, 15, 18]. However, none of the patients carrying disease-causing variants showed QT interval prolongation in our cohort. A recent studies show that a significant number of carriers of pathogenic LQTS gene variants have a baseline QT within the normal range. In genotyped LQTS patients, approximately half of them have no lifetime symptoms, and from 10 to 50% observe no visible QT prolongation in an electrocardiographic pattern [15-16, 19]. Among study participants, a patient (#89) diagnosed with AV block (III stage) has LQTS (QT=506 ms), including symptoms of syncope. Also, the patient's diagnosis is mediated by ventricular arrhythmia. However, disease-causing variants in potassium channel genes were not observed.

Glazer et al. (2021) reported strong associations between pathogenic/likely pathogenic variants in KCNQ1, KCNH2 and KCNE1 and arrhythmias (adjusted ORs 20–25) in an unselected population cohort. Importantly, Glazer and colleagues used a PM2 threshold of MAF < 0.00005 (0.005%) in gnomAD when assessing rarity for arrhythmia syndromes, including LQTS [20]. Notably, among the rare variants identified in our cohort, the VUS KCNQ1 c.1128+4C>T (gnomAD MAF 0.00001) met the PM2 rarity cutoff (PM2 = ACMG/AMP criterion for absence or extreme rarity in population databases [12]).

In our study, among 52 variants in 11 potassium channel genes, particularly, variants with pathogenicity status were found in *KCNH2* (p.Cys66Gly, p.Arg176Trp), *KCNQ1* p.Tyr662Ter, and *KCNE3* p.Thr4Ala. According to ACMG classification, patients #516, #334 and #377 carry a rare mutation in a likely pathogenic variant *KCNH2* p.Cys66Gly, NM_000238.4):c.196T>G. The variant is located within the PAS-domain, a critical regulatory segment of the HERG (*KCNH2*) channel that governs inactivation of *I_{Kr}* current. As described by Gustina et al. (2012), mutations in this domain, particularly within residues 26–135, impair *I_{Kr}* inactivation and promote delayed repolarization [21]. Patient #516 with CHD AF had suffered a myocardial infarction, reduced LVEF (44%) and a familial predisposition to CVD, which suggests a potential modifier effect, contributing to arrhythmogenic risk despite the absence of evident LQTS. A recent population-based cohort study confirmed a strong association between family history and increased occurrence of AF [22]. The presence of AF in relatives approximately doubles the risk of AF and is associated with an earlier disease onset [23]. Consistent with this, the number of disease-associated variants is highest among individuals with AF onset before age 30 [24]; accordingly, our patients (#377, #282, #334) with AF before age 30 fit this pattern of hereditary early-onset AF. Our findings demonstrate that clinically relevant variants in potassium channel genes are detectable in a considerable proportion of Kazakhstani patients with arrhythmias, supporting the growing recognition that ion channel dysfunction contributes not only to classical long QT or Brugada syndromes but also to more common arrhythmia phenotypes such as atrial fibrillation and conduction disorders. Patient # 334 with iAF has a manifesting paroxysmal orthodromic AV-reentry tachycardia and Wolff-Parkinson-White (WPW) syndrome. The latter is associated with increased risk of supraventricular tachyarrhythmia and SCD [25]. According to ACMG classification, both patients (#202 and #573) carry a likely pathogenic variant *KCNH2* p.Arg176Trp (rs36210422), class 4 (LP). Notably, both individuals presented with paroxysmal forms of AF, had no documented family history, and demonstrated normal LVEF (64.7%, 52%). Nevertheless, the allele frequency in gnomAD is quite low, 0.000618 ($\approx 0.062\%$), which indicates that the variant is extremely rare in the general population and further supports its disease-causing classification. Rare variants in ion channel genes, including potassium channels, have been linked to monogenic forms of AF through mechanisms that alter atrial refractoriness, action potential duration, or intercellular conduction [26]. Such electrophysiological disturbances may explain the occurrence of AF itself. Despite the absence of structural heart disease or QT prolongation, the presence of this variant in two unrelated cases supports its potential pathogenic role in arrhythmogenesis. A recent large-scale cohort study confirmed that *KCNH2* p.Arg176Trp was significantly enriched among individuals referred for LQTS genetic testing, particularly those with no identified genetic cause, suggesting that this variant may act as a risk-modifying allele under specific triggers such as medication or stress [27].

Based on the HGMD database, several rare variants were found in key potassium channel genes (such as *KCNQ1* Tyr662Ter and *KCNH2* p.Tyr652Ter, p.Cys66Gly, p.Arg176Trp), suggesting evidence for strong associations with arrhythmia phenotypes. (Table 4). Although ClinVar shows mixed interpretations, their classification as disease-causing in HGMD and their very low frequency in the population (gnomAD < 0.001) support their potential pathogenic impact. Two additional variants (Pro307Ser in *KCNA5* and Arg1047Leu in *KCNH2*) have been labeled as functionally abnormal, DFP, strengthening their clinical importance. Other variants (Lys897Thr and Ser38Gly) with DF/FP status, are likely benign but could still influence risk depending on the patient's genetic background. Interestingly, several of the identified variants in our cohort, including *KCNQ1* p.Tyr662Ter, *KCNH2* p.Tyr652Ter, and *KCNE3* p.Thr4Ala, have been reported in previous studies as disease-causing mutations primarily in European or East Asian populations [28-29]. However, their detection in a Central Asian cohort highlights the potential for both shared and population-specific genetic architectures of arrhythmic disorders. This underscores the importance of expanding genetic studies to include underrepresented populations, as founder effects and local allele frequencies may influence variant interpretation and clinical management. Two novel variants detected in *KCNE5* and *KCND3* may represent population-specific polymorphisms or low-frequency pathogenic alleles. Establishing regional genomic reference data is essential for accurate variant

interpretation under ACMG criteria and to prevent misclassification of benign ethnic-specific variants.

Additionally, VUS detected in *KCNQ1*, *KCNE2*, *KCNE3*, *KCNJ8*. Although the pathogenicity of these variants requires further functional validation, prior studies indicate that up to 40–45% of VUS in ion-channel genes may exhibit functional defects [30]. Notably, 4 out of 6 patients carrying VUS have a family history predisposition to CVD. VUS and DFP variants may be relevant depending on the clinical context and additional functional analysis. Thus, in symptomatic or familial cases, VUS should not be disregarded but rather integrated into risk assessment models and periodically re-evaluated as population data evolve.

It is important to mention that as multigene panels become more widely used, rare variants are increasingly found in individuals without clear disease. Mutation databases are full of benign/likely benign variants that were previously known to be disease-causing; consequently, interpretation of genetic variants is quite complicated [13]. Therefore, in our study we systematically reannotated identified variants and updated their classifications by cross-checking population databases.

As outlined in the 2022 European Society of Cardiology (ESC) guidelines and the Heart Rhythm Society (HRS) consensus statement, genetic testing is recommended for patients with unexplained syncope, idiopathic ventricular tachycardia, or family history of SCD. Integrating genetic results into cardiology practice offers several benefits. Namely, carriers of *KCNH2* or *KCNQ1* variants should avoid QT-prolonging drugs (antiarrhythmics, certain antibiotics, psychotropics). Regular ECG or Holter monitoring is necessary to detect repolarization changes. Patient's at-risk relatives are advised to undergo family cascade testing for early detection of carrier variants. Finally, individualized management strategies based on genotype–phenotype correlations can be integrated [9, 13]. Our findings provide empirical support for extending these recommendations to the Central Asian population.

Several limitations of this study should be acknowledged. First, the relatively small cohort size (79 patients) limits the generalizability and statistical power to detect genotype–phenotype correlations or estimate variant frequencies with precision. Second, although we used a comprehensive 96-gene targeted NGS panel providing high coverage and sensitivity, this approach does not detect structural variants, deep intronic changes, large copy-number alterations, or variants in genes outside the panel that may also contribute to arrhythmogenic phenotypes. Third, segregation analysis was not systematically performed in affected families, which restricts the ability to confirm variant pathogenicity through inheritance patterns. Fourth, functional validation of the identified variants, particularly those of uncertain significance (VUS) and novel findings, was not conducted; therefore, their predicted effects remain speculative. Fifth, QT interval measurements and clinical data were collected at a single time point, and longitudinal follow-up could provide additional insight into variable expressivity or age-related penetrance. Finally, population-specific allele frequencies in Central Asian groups remain poorly characterized, which complicates the interpretation of variant rarity and potential founder effects. We are performing large scale populational study in Kazakhstan to establish Kazakhstani Reference Database of genetic variants. Future studies with larger multiethnic cohorts, extended family analyses, and electrophysiological characterization of novel variants are warranted to validate and expand upon these findings.

5. Conclusions

This study provides the first comprehensive assessment of potassium channel gene variants in Kazakhstani patients with cardiac arrhythmias. Identification of pathogenic and novel variants demonstrates the genetic contribution to arrhythmia pathogenesis and underscores the clinical utility of NGS-based testing. Integration of genetic diagnostics into cardiology practice can improve patient stratification, optimize therapy, and enable family-based prevention strategies in Central Asia.

We identified several clinically significant variants and two previously unreported potassium channel variants in a cohort of 79 patients with cardiac arrhythmias using a 96-gene targeted NGS panel (Haloplex). None of these novel variants were detected in 234 healthy Kazakh controls. Variant classification followed ACMG/AMP criteria, and unreported variants were further assessed using multiple in silico prediction tools; however, only a subset met rarity thresholds in population

databases. These results highlight the utility of high-coverage targeted gene panels for detecting rare, potentially population-specific ion channel variants. To establish their pathogenicity and clinical relevance, segregation analyses and in vitro functional validation using patient-derived iPSC cardiomyocyte models are warranted. Functional validation of novel variants and family-based segregation analysis will be crucial next steps toward implementing precision cardiology in Central Asia.

6. Patents

Patent of the Republic of Kazakhstan for Utility Model No. 4574. Method for detecting genetic predisposition to cardiac arrhythmias based on a developed cardiogenetic panel of 96 candidate genes / Akilzhanova A.R., Abilova Zh.M., Akhmetova A.Zh., Kozhamkulov U.A., Rakhimova S.E., Kairov U.E. Published on 26.12.2019, application No. 2019/0880.2 dated 11.10.2017.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/doi/10.3390/pre25112094>, Table S1. Clinical characteristics of patients; Table S2: List of genetic variants in potassium channel genes.

Author Contributions: Conceptualization, A.C. and A.Ak.; methodology, S.R., A.C., Z.A., M.Z., A.Ak.; bioinformatics data analysis and software, U.K., A.D., A.M.; clinical data and patient treatment, G.A., K.A., A.K., N.S., A.A., M.B.; writing—original draft preparation, A.C. A.Ak.; writing—review and editing, S.R., A.Ak. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Committee of Science of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP19677442, AP23490249, BR24993023, BR24992841, BR27199879) and Nazarbayev University CRP (211123CRP1608, Ainur Akilzhanova).

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Local Ethics Committee of National Laboratory Astana, Nazarbayev University (pro tocol № 16, from 11 March 2015) and by the Ethics Committee of National Research Cardiac Surgery Center (NRCC), Astana, Kazakhstan (protocol № 16, from 24 February 2015).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: All sequence data presented in this study are deposited in the NCBI Sequence Read Archive (SRA) repository and are publicly available under accession BioProject number PRJNA908657 (<https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA908657>). The data presented in this study are available on request from the corresponding author. The data are not publicly available due to ethical and privacy restrictions indicated in informed consent.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

CVD	Cardiovascular diseases
NGS	Next-generation sequencing
SCD	Sudden cardiac death
LQTS	Long QT syndrome
SQTS	Short QT syndrome
BrS	Brugada syndrome
AF	Atrial fibrillation
CHD	Coronary heart disease
HGMD	Human Gene Mutation Database
ACMG	American College of Medical Genetics and Genomics

References

1. World Health Organization. [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
2. Tan, S. M. L., Lim, S. L., Ong, M. E., Leong, K. M. Genetics of Sudden Cardiac Arrest: Overview of Genetic Risk Factors and Aetiologies. *Arrhythmia & Electrophysiology Review*, **2025**, *14*, e18.
3. Zheng, J., Zhang, J., Danioko, S., Yao, H., Guo, H., Rakovski, C. A 12-lead electrocardiogram database for arrhythmia research covering more than 10,000 patients. *Scientific data*, **2020**, *7*(1), 48.
4. Dib Nehme, R., Sinno, L., Shouman, W., Ziade, J.A., Ammar, L.A., Amin, G., Booz, G.W. and Zouein, F.A. Cardiac Channelopathies: Clinical Diagnosis and Promising Therapeutics. *Journal of the American Heart Association*, **2025**, *14*(9), e040072.
5. Schwartz, P. J., Ackerman, M. J., Antzelevitch, C., Bezzina, C. R., Borggrefe, M., Cuneo, B. F., Wilde, A. A. Inherited cardiac arrhythmias. *Nature reviews Disease primers*, **2020**, *6*(1), 58.
6. Pesaresi, M., Bernini Di Michele, A., Melchionda, F., Onofri, V., Alessandrini, F., Turchi, C. Sudden Cardiac Death and Channelopathies: What Lies behind the Clinical Significance of Rare Splice-Site Alterations in the Genes Involved?. *Genes*, **2024**, *15*(10), 1272.
7. Guelly, C., Abilova, Z., Nuralinov, O., Panzitt, K., Akhmetova, A., Rakhimova, S., Kozhamkulov, U., Kairov, U., Molkenov, A., Ashenova, A. Trajanoski, S. Patients with coronary heart disease, dilated cardiomyopathy and idiopathic ventricular tachycardia share overlapping patterns of pathogenic variation in cardiac risk genes. *PeerJ*, **2021**, *9*, e10711.
8. Priori S.G., Blomström-Lundqvist C., Mazzanti A., Blom N., Borggrefe M., Camm J., Elliott P.M., Fitzsimons D., Hatala R., Hindricks G., Kirchhof P., Kjeldsen K. ESC Scientific Document Group. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: The Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC). *European heart journal*, **2015**, *36*(41), 2793–2867.
9. Zeppenfeld, K., Tfelt-Hansen, J., De Riva, M., Winkel, B.G., Behr, E.R., Blom, N.A., Charron, P., Corrado, D., Dagres, N., De Chillou, C. and Eckardt, L. 2022 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: Developed by the task force for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death of the European Society of Cardiology (ESC) Endorsed by the Association for European Paediatric and Congenital Cardiology (AEPC). *European heart journal*, **2022**, *43*(40), 3997-4126
10. Zhu, W., Bian, X., Lv, J. From genes to clinical management: A comprehensive review of long QT syndrome pathogenesis and treatment. *Heart Rhythm O2*, **2024**, *5*(8), 573-586.
11. Broendberg, A. K., Christiansen, M. K., Nielsen, J. C., Pedersen, L. N., Jensen, H. K. Targeted next generation sequencing in a young population with suspected inherited malignant cardiac arrhythmias. *European journal of human genetics : EJHG*, **2018**, *26*(3), 303–313.
12. Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., Grody, W. W., Hegde, M., Lyon, E., Spector, E., Voelkerding, K., Rehm, H. L., & ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in medicine : official journal of the American College of Medical Genetics*, **2015**, *17*(5), 405–424.
13. Wilde, A. A. M., Semsarian, C., Márquez, M. F., Shamloo, A. S., Ackerman, M. J., Ashley, E. A., Sternick, E. B., Barajas-Martinez, H., Behr, E. R., Bezzina, C. R., Breckpot, J., Charron, P., Chockalingam, P., Crotti, L., Gollob, M. H., Lubitz, S., Makita, N., Ohno, S., Ortiz-Genga, M., Sacilotto, L. Developed in partnership with and endorsed by the European Heart Rhythm Association (EHRA), a branch of the European Society of Cardiology (ESC), the Heart Rhythm Society (HRS), the Asia Pacific Heart Rhythm Society (APHRS), and the Latin American Heart Rhythm Society (LAHRS). European Heart Rhythm Association (EHRA)/Heart Rhythm Society (HRS)/Asia Pacific Heart Rhythm Society (APHRS)/Latin American Heart Rhythm Society (LAHRS) Expert Consensus Statement on the state of genetic testing for cardiac diseases. *Europace : European pacing, arrhythmias, and cardiac electrophysiology : journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology*, **2022**, *24*(8), 1307–1367.

14. Crotti, L., Odening, K. E., & Sanguinetti, M. C. Heritable arrhythmias associated with abnormal function of cardiac potassium channels. *Cardiovascular research*, **2020**, *116*(9), 1542-1556.
15. Krahn, A. D., Laksman, Z., Sy, R. W., Postema, P. G., Ackerman, M. J., Wilde, A. A., Han, H. C. Congenital long QT syndrome. *Clinical Electrophysiology*, **2022**, *8*(5), 687-706.
16. Wilde, A. A., Amin, A. S., Postema, P. G. Diagnosis, management and therapeutic strategies for congenital long QT syndrome. *Heart*, **2022**, *108*(5), 332-338.
17. Offerhaus, J. A., Bezzina, C. R., Wilde, A. A. Epidemiology of inherited arrhythmias. *Nature Reviews Cardiology*, **2020**, *17*(4), 205-215.
18. Vink, A.S., Neumann, B., Lieve, K.V., Sinner, M.F., Hofman, N., El Kadi, S., Schoenmaker, M.H., Slaghekke, H.M., De Jong, J.S., Clur, S.A.B., Blom, N.A. Determination and interpretation of the QT interval: comprehensive analysis of a large cohort of long QT syndrome patients and controls. *Circulation*, **2018**, *138*(21), 2345-2358.
19. European Society of Cardiology. How to measure the QT interval. *Cardiovascular Genomics Insight*, Volume 9. Available online: <https://www.escardio.org/Councils/Council-on-Cardiovascular-Genomics/Cardiovascular-Genomics-Insight/Volume-9/how-to-measure-the-qt-interval>. Accessed July 4, 2025.
20. Glazer, A.M., Davogustto, G., Shaffer, C.M., Vanoye, C.G., Desai, R.R., Farber-Eger, E.H., Dikilitas, O., Shang, N., Pacheco, J.A., Yang, T., Muhammad, A. Arrhythmia variant associations and reclassifications in the eMERGE-III sequencing study. *Circulation*, **2022**, *145*(12), 877-891.
21. Gustina, A.S. and Trudeau, M.C. HERG potassium channel regulation by the N-terminal eag domain. *Cellular signalling*, **2012**, *24*(8), 1592-1598.
22. Shah, A.S., Ongtengco, A., Qiao, V., Chen, Y., Diaz, A., Hill, M., Bhan, A., Tofovic, D.S. Darbar, D. Association Between Family History and Early-Onset Atrial Flutter Across Racial and Ethnic Groups. *Journal of the American Heart Association*, **2024**, *13*(10), e032320.
23. Chang, S.H., Kuo, C.F., Chou, I.J., See, L.C., Yu, K.H., Luo, S.F., Huang, L.H., Zhang, W., Doherty, M., Wen, M.S. Kuo, C.T. Association of a family history of atrial fibrillation with incidence and outcomes of atrial fibrillation: a population-based family cohort study. *JAMA cardiology*, **2017**, *2*(8), 863-870.
24. Yoneda, Z.T., Anderson, K.C., Quintana, J.A., O'Neill, M.J., Sims, R.A., Glazer, A.M., Shaffer, C.M., Crawford, D.M., Stricker, T., Ye, F. and Wells, Q.. Early-onset atrial fibrillation and the prevalence of rare variants in cardiomyopathy and arrhythmia genes. *JAMA cardiology*, **2021**, *6*(12), 1371-1379.
25. Bowles, N.E., Jou, C.J., Arrington, C.B., Kennedy, B.J., Earl, A., Matsunami, N., Meyers, L.L., Etheridge, S.P., Saarel, E.V., Bleyl, S.B., Yost, H.J. Exome analysis of a family with Wolff-Parkinson-White syndrome identifies a novel disease locus. *American Journal of Medical Genetics Part A*, **2015**, *167*(12), 2975-2984.
26. Hayashi, K., Konno, T., Tada, H., Tani, S., Liu, L., Fujino, N., Nohara, A., Hodatsu, A., Tsuda, T., Tanaka, Y. and Kawashiri, M.A.. Functional characterization of rare variants implicated in susceptibility to lone atrial fibrillation. *Circulation: Arrhythmia and Electrophysiology*, **2015**, *8*(5), 1095-1104.
27. Vatta, M., Truty, R., Garcia, J., Callis, T.E., Hatchell, K., Rojahn, S., Morales, A., Aradhya, S. and Nussbaum, R. Common variants in KCNE1, KCNH2, and SCN5A may impact cardiac arrhythmia risk. *Circulation: Genomic and Precision Medicine*, **2021**, *14*(1), p.e003206.
28. Nakajima, T., Wu, J., Kaneko, Y., Ashihara, T., Ohno, S., Irie, T., Ding, W.G., Matsuura, H., Kurabayashi, M., Horie, M. KCNE3 T4A as the genetic basis of Brugada-pattern electrocardiogram. *Circulation Journal*, **2012**, *76*(12), 2763-2772.
29. Kotta, M.C., Torchio, M., Bayliss, P., Cohen, M.C., Quarrell, O., Wheeldon, N., Marton, T., Gentilini, D., Crotti, L., Coombs, R.C. and Schwartz, P.J. Cardiac genetic investigation of sudden infant and early childhood death: a study from victims to families. *Journal of the American Heart Association*, **2023**, *12*(17), p.e029100.33
30. Ye, D., Gao, X., Bains, S., Neves, R.A., Zhou, W., Tester, D., Giudicessi, J.R. Ackerman, M.J. PO-01-240 UTILITY OF IN VITRO FUNCTIONAL CHARACTERIZATION IN THE PROMOTION OR DEMOTION OF CLINICALLY ENCOUNTERED VARIANTS OF UNCERTAIN SIGNIFICANCE IN THE THREE MOST COMMON LONG QT SYNDROME-CAUSATIVE GENES: KCNQ1, KCNH2, AND SCN5A. *Heart Rhythm*, **2023**, *20*(5), S166.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.