

Essay

Not peer-reviewed version

Escalating Dengue in Bangladesh: An Analytical Assessment of Environmental and Socioeconomic Drivers

[Abdul Kader Mohiuddin](#)*

Posted Date: 29 December 2025

doi: 10.20944/preprints202512.2526.v1

Keywords: dengue epidemiology; *Aedes* mosquito breeding; pandemic outbreaks; climate-driven transmission; urbanization impact; insecticide resistance; public health burden; micro plastic environmental risk



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Essay

Escalating Dengue in Bangladesh: An Analytical Assessment of Environmental and Socioeconomic Drivers

Abdul Kader Mohiuddin

Alumnus, Faculty of Pharmacy, Dhaka University, Bangladesh; dr.m.nasirullah.trust@gmail.com

Abstract

Dengue has emerged as a persistent and escalating public health crisis in Bangladesh, reflecting both local vulnerabilities and a broader global threat. Between 2018 and 2025, the outbreaks have caused substantial morbidity and mortality across all age groups, driven by rapid urbanization, dense populations, poor sanitation, and climatic changes. Environmental degradation, including loss of green cover, pollution, and plastic accumulation, has further intensified mosquito proliferation, while demographic shifts reveal increasing vulnerability among children, juveniles, and young adults. Immediate, coordinated action—including strengthened public health infrastructure, vector control, and community awareness—is imperative to curb the ongoing outbreaks and mitigate the potential for regional and global spread.

Keywords: dengue epidemiology; *Aedes* mosquito breeding; pandemic outbreaks; climate-driven transmission; urbanization impact; insecticide resistance; public health burden; micro plastic environmental risk

Introduction

Each year, mosquitoes wage a silent yet devastating war—infesting nearly 700 million people and claiming more than a million lives across the globe [1]. Mosquito-borne viruses like dengue, chikungunya, and Zika have devastated 166 countries over the last five decades, costing nearly \$100 billion and surging fourteen-fold between 2013 and 2022 [2]. While malaria continues to devastate Africa—accounting for over 90% of cases reported in the WHO African Region [3]—Asia is grappling with dengue, which is responsible for nearly 70% of global infections [4], with Southeast Asia bearing the heaviest burden [5]. Although the COVID-19 pandemic momentarily disrupted this trajectory, the post-pandemic resurgence of dengue infections reveals its persistent grip on the region [6]. In Bangladesh, dengue remained relatively rare before 2018 but surged thereafter, following global trends, briefly paused during the COVID-19 pandemic, and emerged as the deadliest infectious disease in the post-COVID era, peaking in 2023 (Figure 1). This alarming rise, driven by a combination of meteorological changes and overlooked socioeconomic factors, forms the central focus of this paper.




Figure 1. Trends in Dengue Cases and Deaths in Bangladesh, 2000–8 December 2025 (Source: The Institute of Epidemiology, Disease Control and Research, IEDCR) / Directorate General of Health Services, DGHS). The figure depicts a pronounced increase in dengue cases and deaths in Bangladesh during the post-COVID period, showing a closely aligned temporal pattern between case numbers and fatalities.

Methodology

This review synthesizes evidence from leading global health datasets, peer-reviewed literature, major international reports, and national sources including the DGHS, IEDCR, and the Bangladesh Meteorological Department to assess recent trends in dengue transmission in Bangladesh. When contemporary scholarly data were unavailable or incomplete, rigorously verified reports from reputable news media were incorporated to provide timely contextual updates. Environmental and socioeconomic determinants of dengue transmission were systematically integrated with key climatic variables—temperature, humidity, and rainfall—to construct a multidimensional understanding of transmission dynamics. Given the rapidly evolving nature of dengue epidemiology in Bangladesh, the analysis also employed cautiously framed predictive inferences informed by both local and global research. This approach enabled the identification of several under-recognized drivers of dengue virus transmission that may influence future outbreak patterns.

Literature Review

The Current Global Landscape of Dengue

Global dengue incidence is rising at an alarming rate. While reported patterns vary, the surge is undeniable. According to WHO, 6.5 million cases and 8,791 deaths were reported globally in 2023 [7],

rising to 7.6 million cases and 3,000 deaths by April 2024 [8]. Notably, the US CDC and WHO's global dengue surveillance system, launched in May 2024, reported 12–13 million cases and over 7,000 deaths in the Americas alone in 2024 [9,10]. Ranked among WHO's top ten global health threats, dengue affected approximately 90 countries in 2024, with Brazil bearing the highest burden, followed by Argentina and Mexico [11]. Current WHO estimates suggest that dengue causes up to 400 million infections annually [12], with incidence having increased thirtyfold over the past fifty years [13], now placing 3.9–5.6 billion people—more than half of the world's population—at risk [14,15]. Moreover, the US CDC reports that half of the global population lives in dengue-risk areas, putting both residents and travelers at risk [16]. Dengue, primarily transmitted by *Aedes aegypti* and *Aedes albopictus*, disproportionately affects the southern hemisphere, making effective tetravalent vaccines critical for global health [17]. Current vaccine development faces challenges in ensuring protection against all serotypes, addressing varied immune responses, and adapting to emerging strains.

Meteorological and Socioeconomic factors of Dengue Transmission

Environmental and socioeconomic factors jointly drive global dengue transmission, with climate conditions like temperature, rainfall, and humidity shaping mosquito habitats, while urbanization, population density, poverty, and inadequate sanitation increase human exposure and vulnerability, as discussed in Table 1.

Table 1. Determinants of Dengue Transmission: Climatic, Environmental, and Geographic Factors.

Factor	Key Points	Evidence / Findings
Climate	Shapes dengue ecology by influencing vector dynamics, Relationships between climate variables and dengue virus development, and transmission are complex [18]. mosquito–human interactions.	
Temperature	Rising temperatures increase dengue risk.	Regions with notable warming, such as sub-Saharan Africa and Oceania, show higher dengue incidence [19].
Rainfall	Provides breeding sites for mosquitoes.	Excess rainfall can wash away breeding sites, affecting outbreak patterns [20,21].
Humidity	Higher humidity supports mosquito survival and virus transmission.	Humidity $\geq 60\%$ and temperatures $>27^\circ\text{C}$ elevate dengue risk; mosquitoes rarely survive below 60% humidity [21,22].
Wind Speed	Influences mosquito activity and breeding.	Higher wind speeds reduce transmission by limiting mosquito flight, host-seeking, and breeding-site availability [23].
Environmental Conditions	Impact mosquito breeding and dengue transmission.	Key factors include water storage, waste disposal, housing, drainage, vegetation, urbanization, seasonal variation, and water supply [24].
Geography	Vector thrives in tropical and subtropical regions.	Spread is dictated by climate, urbanization, and population movement [12].
Latitude	Determines the global range of dengue.	<i>Aedes aegypti</i> thrives between $\sim 35^\circ\text{N}$ – 35°S ; warming expands risk to higher latitudes, including Africa, South America, southern China, and the U.S. [19,25].
Altitude	Limits mosquito habitats.	Mosquitoes generally stay below 6,500 ft; common up to 1,700 m, rare between 1,700–2,130 m; warming may increase risk at higher elevations [16,25,26].

Key Risk Factors for Dengue in Bangladesh: Insights from Previous Research

Bangladesh, a tropical country in South Asia situated between 20° – 27°N and 88° – 93°E , occupies the world's largest and most densely populated delta—the low-lying Ganges–Brahmaputra Delta,

formed by rivers originating in the Himalayas. With more than 1,000 rivers spanning roughly 24,140 km and a coastline along the Bay of Bengal, the nation experiences high humidity and heavy rainfall, making it particularly vulnerable to floods and tropical cyclones [27–29]. Over the past four decades, temperatures in Bangladesh have risen by approximately 0.5 °C, which has lengthened the dengue season and accelerated transmission, with case numbers doubling roughly every decade since 1990. The World Bank notes that dengue infections increase significantly at temperatures between 25 °C and 35 °C, peaking around 32 °C, while global mosquito transmission capacity has grown by up to 9.5% since 1950 [30].

A study published in Oxford Academic on dengue transmission in Bangladesh (2000–2022) found that rising temperatures and shifts in rainfall patterns between 2011 and 2022 were closely linked to increased cases and deaths [31]. Recent climatic shifts have further intensified the risk. From 2022 to 2025, October rainfall in Bangladesh has been highly variable, with a trend toward heavier late-monsoon precipitation. Such changes have prompted entomologists to warn of prolonged dengue outbreaks [32–34]. A comprehensive review of three major medical databases—PubMed, Embase, and Web of Science—up to December 5, 2024, indicates that climate change is reshaping temperature, rainfall, and humidity patterns, thereby expanding the geographic range of dengue and altering exposure risks across different populations [35]. Figure 2 shows that, following the global pattern, dengue cases rise in tandem with increases in temperature, rainfall, and humidity.

Evaluating the various dengue determinants in Bangladesh from 2000 to 2023, researchers from Begum Rokeya University used XGBoost and LightGBM with explainable AI to identify population density, precipitation, temperature, and land-use as key predictors, aligning with recent studies and supporting early-warning systems [36]. A comparative analysis with Singapore found that rainfall fueled dengue transmission in Bangladesh while humidity and sunshine suppressed it—whereas in Singapore, warmer temperatures drove infections and rainfall and humidity helped curb spread [37]. Hossain et al. (2023) identified rapid urbanization, climatic suitability, and the persistent presence of *Aedes* mosquitoes as key drivers of increased human–vector contact and the expanding geographic reach of dengue. Periodic serotype shifts, weak surveillance, limited healthcare capacity, and low public awareness further intensify these risks [38]. Building on this, Khan et al. (2024) highlighted possible post-COVID immune effects, climate variability, dominant viral serotypes, and systemic failures in patient management as contributors to Bangladesh's recent high fatality rates, underscoring the need for stronger clinical care, more trained personnel, improved vector control, and investment in One Health–based prevention [39].

Examining seasonal dengue patterns from January 2008 to November 2024, Alam et al. (2025) showed that incidence is tightly linked to meteorological conditions, with peaks strongly correlated with higher temperatures, humidity, rainfall, and wind speed. Their study emphasized the need for future models to integrate real-time meteorological inputs along with urbanization and socioeconomic factors [40]. Islam et al. (2023) similarly argued that combining climate projections with human mobility and socio-environmental variables is essential for forecasting outbreaks and guiding effective prevention strategies [41]. Supporting this, Islam and Hu (2024) identified rapid human movement as a major transmission driver in Bangladesh, with festival gatherings, increased mobility, and post-lockdown shifts all associated with higher case burdens [42]. Ogieuhi et al. (2025) noted that poor sanitation, insecticide resistance, limited vaccine access, low public awareness, and mounting healthcare pressures, combined with climate change and rapid urbanization, collectively heighten dengue risks, especially for vulnerable populations [43].




Figure 2. Trend in Dengue Cases in Bangladesh, 2018–November 2025 (Source: DGHS/Bangladesh Meteorological Department). The figure illustrates a clear upward trend in dengue transmission in Bangladesh since 2019, with a brief pause in 2020 consistent with global patterns. From June onwards, cases begin to rise sharply, closely following periods of elevated temperature, humidity, and rainfall. Peaks in dengue incidence consistently occur between September and November in recent years, a pattern that meteorologists attribute to the unusually high rainfall observed in October from 2022 to 2025 (visualized using Canva Illustrator).

Common Public Perception Vs Reality

In Bangladesh, dengue perception shows a mix of high awareness of its severity (it's deadly) but low personal risk (susceptibility), leading to inconsistent prevention, with educated urban dwellers often better informed than rural populations. Table 2 offers an overview of dengue-related knowledge, perception, and attitudes across different Bangladeshi populations.

Table 2. Knowledge, Perception, and Attitudes Towards Dengue in Various Bangladeshi Populations.

Study Place/ Population	Knowledge	Perception & Attitude
1,358 youths of capital Dhaka	Higher climate change knowledge; links with dengue awareness	Positive attitude toward dengue–climate connection; socio-demographic/lifestyle factors influence awareness [44]
Students via social media survey	Strong climate-change awareness; weak dengue-prevention knowledge	Solid attitudes; past dengue experience predicts preventive behaviors [45]
1,010 respondents across 9 regions	Widespread awareness; educated/urban/better-off higher knowledge	Misconceptions persist (e.g., <i>Aedes</i> breed hadin dirty water); weak preventive practices [46]
Dhaka university students	Good knowledge/practices; gaps in transmission, breeding sites, pregnancy-related risks	Strong attitudes; mixed-unit residents showed weakest preparedness [47]
745 slum dwellers of Dhaka	Recognized dengue severity and transmission	Low perceived personal risk; 60% inadequate preventive measures [48]
1,905 Northern-region residents	Limited awareness; poor understanding of climate-disease link	Perception and attitude not well- developed [49]
401 rural residents, Savar	Moderate knowledge; influenced by education, age, gender, occupation, health beliefs	High perceived severity; preventive practices unsatisfactory [50]
364 rural adults from Puthia & Paba upazila	48.4% had sufficient knowledge; higher education → better awareness	Gaps in understanding transmission/prevention; attitude not emphasized [51]
Scoping review of 27 studies	Moderate knowledge overall; rural/slum populations lower	Varying perception; rural/slum communities had weak preventive practices [52]
484 adults of Cox's Bazar	Average knowledge (84.3%)	Positive attitude (63%); knowledge/attitude linked to preventive practices [53]

The recent dengue outbreaks, driven by shifting climate patterns, rapid urbanization, dense populations, insecticide resistance, and low public awareness, have severely strained Bangladesh's healthcare system and economy. While climate change strongly shapes dengue (*Flavivirus*) transmission, insecticide misuse and rising resistance also play critical roles. WHO has warned that fogging is ineffective against *Aedes* mosquitoes, underscoring city corporations' misplaced reliance on mass spraying instead of source reduction, targeted larviciding, and proper vector control. Compounding the problem, widespread metabolic resistance and common kdr mutations have greatly reduced the effectiveness of pyrethroid insecticides, producing very low mosquito mortality even at elevated doses [54–56].

Rainfall influences mosquito growth in complex ways. While light rain creates standing water ideal for breeding, heavy rainfall can destroy breeding sites or wash away larvae, limiting mosquito development. Additionally, wind speed was found to be weakly positively correlated with dengue incidence in Bangladesh. Although many attributed the 2025 outbreak to heavy rainfall, the persistence of dengue had already been evident, with over 320,000 infections and 1,700 deaths recorded in 2023—figures considerably higher than those observed in 2025 (Figure 3). Interestingly, a study conducted in Dhaka revealed that dengue cases actually declined with increasing levels of both rainfall and sunshine, contradicting common public perception [57]. Experts warn that prolonged monsoons and poor waste management have created stagnant water and ecological imbalance, enabling mosquitoes to breed more extensively and intensifying the outbreaks [58].




Figure 3. Monthly Incidence of Dengue Cases and Dengue-Related Deaths in Bangladesh up to November 2025 (Source: DGHS). The figure shows a sharp rise in dengue cases and deaths beginning in June, with reported infections nearly quadrupling by October. The situation peaked in November 2025, when more than 24,500 cases and 100 deaths were recorded—over a quarter of the year’s total burden concentrated in a single month.

Discussion

Escalating Dengue Burden in Bangladesh

Bangladesh is at the epicenter of the crisis, grappling with unprecedented challenges. By 21 September 2025, deaths had surged 150% and cases had doubled from the previous year [59], and just two months later, by 23 November, infections had topped 90,000 with fatalities reaching 364 [60]—70% higher than six weeks earlier [61]. Hospital admissions, according to dynamic data from the Directorate General of Health Services (DGHS) [62], nearly quadrupled from 5,951 in June to 22,520 in October 2025, pushing an already fragile healthcare system to the brink (Figure 3).

November 2025 brought the crisis to a new peak: on 18 November alone, over 900 viral fever patients flooded hospitals, joining nearly 3,000 dengue cases already under treatment [63]. Since 2023, more than half a million Bangladeshis have been infected and over 2,670 have died—marking the deadliest dengue toll in the nation’s history. By the end of November, total cases had surpassed 94,300, hospitalizations had exceeded 92,000, and deaths had risen to 382. November alone recorded more than 24,500 cases and 104 fatalities, meaning that over one-quarter of the year’s infections and deaths occurred in a single, devastating month [62].

Historical data magnify the crisis. Between 2000 and 2022, Bangladesh recorded 853 dengue-related deaths [39,64], yet 2023 alone more than doubled that total, with 1,705 fatalities and over 321,000 infections—the largest annual outbreak on record (Figure 1). The demographic landscape is shifting. In 2023, women represented roughly 40% of dengue cases but accounted for 57% of deaths [65]. By December 8, 2025, men experienced nearly twice as many cases and over half of all deaths

(Figure 4). Notably, in 2023, older adults faced disproportionately severe dengue and higher mortality due to immune vulnerability and comorbidities, with each additional decade raising fatality by 30%, whereas by 2025, young adults aged 21–30 accounted for over a quarter of both cases and deaths [62,65]. However, older adolescents and young adults also represented more than half of all cases during the 2016, 2018, and 2019 outbreaks [65].




Figure 4. Demographic shifts in male and female cases and deaths, 2023–December 8, 2025 (Data Source: DGHS). The figure illustrates a pronounced demographic transition: male cases surged sharply in 2024 before leveling off in 2025, whereas female cases initially declined and later partially rebounded. Concurrently, male deaths exhibited a steady rise, surpassing female deaths by 2025—a striking reversal from previous years (visualized using Canva Illustrator).

Results and Findings

The Overlooked Drivers of Bangladesh's Escalating Dengue Crisis

The recent dengue outbreaks, fueled by changing climate patterns, rapid urbanization, high population density, insecticide resistance, and low public awareness, have placed a severe strain on Bangladesh's healthcare system and economy. While climate change and urban growth are widely acknowledged as major drivers of the rising dengue burden, several less-discussed factors—often tied to uncontrolled urbanization—have intensified the crisis; these interconnected issues, highlighted in recent international research and media, remain largely overlooked by the public due to limited awareness.

Vegetation Loss, and Rising Temperature

Warmer temperatures accelerate mosquito aging, shortening their lifespan and altering infection patterns [66]. Yet, over successive generations, heat-exposed mosquitoes can develop greater

tolerance to viruses without losing vitality, a recent study shows [67]. Global warming has thus become a “perfect storm” for mosquito-borne diseases, affecting every stage of transmission [68].

Urbanization-driven loss of natural vegetation further elevates dengue risk, as areas with reduced green cover provide ideal conditions for mosquito breeding and disease spread, as demonstrated in studies from Mexico [69] and Brazil [70]. In Amazonian Brazil, for example, deforestation of just one square kilometer was linked to 27 additional malaria cases [71].

Between 1989 and 2020, Dhaka lost more than half of its green cover due to rapid urban growth, triggering a significant rise in temperatures [72]. Over three decades, the number of extreme heat days ($\geq 35^\circ\text{C}$) nearly doubled, making Dhaka one of the fastest-warming cities in the world, according to the International Institute for Environment and Development [73]. Furthermore, the World Bank reports that the city’s heat index has increased more than 65% faster than the national average [74]. These hotter, denser conditions let *Aedes* mosquitoes adapt to heat, building stronger virus tolerance and becoming even more efficient carriers [63]. A climate projection from a decade ago indicates that, without adaptation, a 3.3°C increase by 2100 could result in more than 16,000 additional dengue cases [75].

Population Density, Poor Sanitation, and Waste Disposal

Rapid urbanization and extreme population density in Bangladesh are creating ideal conditions for intensified dengue transmission. In overcrowded cities with inadequate sanitation, stagnant water accumulates easily, offering abundant breeding sites for *Aedes* mosquitoes. Dhaka—home to more than 75,000 people per square mile [76]—is now the world’s second most densely populated city [77], and its tightly packed, human-built landscape accelerates *Aedes aegypti* growth, reproduction, and survival far more than suburban or rural settings [78]. Monsoon-season spikes in heat, humidity, and rainfall further amplify this risk, with 2019 data showing that nearly 90% of dengue cases erupted between June and October, overwhelmingly concentrated in the city’s hottest, most densely built neighborhoods [79]. Dengue hotspots consistently emerge where population density is highest, particularly in Thanas such as Badda, Jatrabari, Kadamtali, Mirpur, Mohammadpur, Sobujbagh, Shyampur, Tejgaon, Dhanmondi, and Uttara, where close human-mosquito contact further amplifies transmission [80].

In Bangladesh, roughly 40% of the population lives in urban areas, with over half residing in densely packed slums [81]. Communities without adequate sanitation—especially in these overcrowded settlements—are highly vulnerable to mosquito-borne diseases such as dengue and chikungunya [82]. Dhaka’s congested neighborhoods, compounded by poor sanitation, provide abundant stagnant water, creating ideal breeding grounds for mosquitoes. More than one-third of the population still lacks access to safely managed sanitation, and UNICEF estimates that about 230 tons of fecal waste enter Dhaka’s 4,500-kilometer drainage network every day. The system is already 70% clogged with trash and debris because of poor infrastructure and longstanding neglect, according to the Institute of Water Modelling [83,84]. As a result, even moderate rainfall creates stagnant, mosquito-infested pools—a problem further intensified by flooding and extreme weather across both urban and rural areas [85]. Additionally, in many dense urban neighborhoods, inconsistent water supply forces residents to store water in containers [43], a practice well documented in neighboring India, further increasing the risk of mosquito-borne diseases [86].

Poor waste management is a critical driver of dengue risk among both children and adults—and in urban Bangladesh this threat looms large. Shockingly, 55% of solid waste in urban areas remains uncollected, creating ideal breeding grounds for the mosquitoes that spread the disease [87]. Evidence from urban Thiruvananthapuram, South India, indicates that inadequate waste management infrastructure can be associated with a 40% higher incidence of dengue and chikungunya cases [88]. Likewise, studies in informal urban settlements in Indonesia and Fiji reported that by age 4–5, over half of children had already been infected, highlighting how insufficient waste disposal accelerates early exposure to dengue [89].

Pollution as a Trigger for Viral Resistance and Mosquito Dynamics

The WHO estimates that nearly a quarter of human diseases and deaths stem from long-term exposure to pollution [90]. While research on environmental impacts on dengue in Bangladesh remains limited, international studies underscore their significance. Recent findings from cities in Taiwan [91], Singapore [92], Guangzhou [93], Upper Northern Thailand [94], Melaka, Malaysia [95], and Greater São Paulo [96] demonstrate that air pollutants—such as particulate matter PM_{2.5}, SO₂, O₃, CO, and NO_x—interact with climate factors to influence mosquito populations, viral activity, and human immunity to the virus. These impacts, however, vary depending on pollutant type, concentration, and region, often producing complex, non-linear effects on mosquito dynamics. Interestingly, a study covering 76 provinces in Thailand from 2003 to 2021 found that higher surface concentrations of SO₂ and PM_{2.5} were generally associated with lower incidences of dengue, malaria, chikungunya, and Japanese encephalitis, likely due to adverse effects on mosquito survival and behavior [97]. These findings highlight the need for further research.

A *Lancet* study reported that improperly discarded plastics accumulate stagnant water, creating ideal breeding sites for *Aedes* mosquitoes that transmit dengue, Zika, chikungunya, and yellow fever, thereby directly increasing vector populations. Indirectly, plastic debris also clogs drainage systems, producing large stagnant pools that promote mosquito proliferation and elevate the risk of diseases such as malaria [98]. Bangladesh is now experiencing an alarming rise in micro plastic pollution. Just three rivers—Meghna, Karnaphuli, and Rupsha discharge nearly one million metric tons of mismanaged plastic each year [99]. In total, 36 rivers in Bangladesh are among the 1,656 waterways worldwide responsible for 80% of global riverine plastic emissions [100]. Per-capita plastic consumption has tripled—from 9 kg in 2005 to 2020—while COVID-19 contributed an additional 78,000 tons in a single year, according to a 2021 report by the Environment and Social Development Organization (ESDO) [101].

In Dhaka, per-capita use reaches 24 kg, and nearly one-eighth of all plastic waste ends up in canals and rivers. An estimated 23,000 to 36,000 tons of plastic waste accumulate annually across 1,212 dumping hotspots surrounding the Buriganga, Turag, Balu, and Shitalakhsya rivers, a trend highlighted by a former World Bank country director during a program in Dhaka [102]. Beyond environmental degradation, this rising plastic burden may intensify mosquito-borne disease risks: researchers from the Beijing Institute of Microbiology and Epidemiology show that mosquitoes exposed to micro plastics can transfer them to mammals, develop altered gut microbiomes, experience delayed development, and exhibit reduced insecticide susceptibility—factors that could heighten disease transmission [103]. Also, micro plastics can adsorb pyrethroid insecticides such as deltamethrin, reducing the concentration available to act on mosquitoes. However, because the findings rely on a single study and other research shows conflicting results, more evidence is needed to clarify how micro plastic exposure influences mosquito dynamics and dengue transmission.

Construction Sites and High-Rises: Major Breeding Grounds Driving Dengue in Dhaka

Dhaka's rapid and largely unplanned urban expansion has transformed the city into a highly conducive environment for *Aedes* mosquito proliferation. Numerous under-construction buildings, left exposed to the elements, now serve as prime breeding grounds for the vectors of dengue. Surveys indicate that, in the decade preceding 2016, an average of 95,000 new structures were erected annually within the jurisdiction of the Rajdhani Unnayan Kartripakkha (RAJUK). Over the subsequent fifteen years, at least 64,000 additional buildings were constructed across the capital [104,105]. In July 2020, inspections conducted by the Dhaka North City Corporation (DNCC) revealed that nearly 70% (8,764 out of 12,619) of homes and construction sites surveyed across 55 wards harbored potential *Aedes* breeding sources [106]. These inspections were carried out in collaboration with the National Malaria Elimination and *Aedes* Transmitted Disease Control Programme under the Directorate General of Health Services (DGHS).

The following year, the situation deteriorated further. A 2021 DGHS study covering 70 areas of Dhaka reported alarming *Aedes* densities, with the Breteau Index (BI)—the number of water-holding

containers infested with larvae per 100 houses—rising to 23.3 in Lalmatia and Iqbal Road (Ward 32, DNCC) and 20.0 in Sayedabad and Uttar Jatrabari (Ward 48, DSCC). High-rise buildings accounted for over 45% of breeding sites, followed by under-construction structures at nearly 35% [107]. In 2024, the former Mayor of DSCC warned that construction would be halted wherever *Aedes* larvae were detected and that dengue control drives would be launched in advance of the rainy season, alongside the government's seven-year National Dengue Prevention and Control Strategy [108]. The most recent pre-monsoon survey, conducted jointly by the DGHS Communicable Disease Control Programme and the Institute of Epidemiology, Disease Control and Research (IEDCR), presents a similarly concerning picture: multistory buildings accounted for almost 60% of *Aedes* larvae, with a further 20% found in under-construction sites [109].

From Neglect to Epidemic: How Policy Failures Worsened Dengue in Bangladesh

Bangladesh's authorities have repeatedly failed to curb *Aedes* populations, persisting with outdated chemical approaches while neglecting structural determinants and community-level interventions. Government action has remained fragmented and reactive; in 2023, officials proved unable to control *Aedes* mosquitoes, opting instead to fault households and impose ethically questionable fines. Such mismanagement and flawed strategies have allowed dengue transmission to escalate unchecked, rendering official prevention efforts largely performative (Figure 5). Transparency International Bangladesh has identified several drivers of high mortality, including inadequate hospital staffing, delayed diagnoses, false-negative NS1 results, weak vector-control measures, and limited healthcare capacity beyond Dhaka [110]. Experts further warn that the absence of strategic planning, non-adherence to WHO guidelines, and the failure to involve qualified public-health professionals have deepened the crisis. By 2024, South Asia was experiencing its most severe dengue epidemic on record, with Bangladesh and India reporting thousands of deaths as hospitals were overwhelmed. Concerns have mounted over inadequate anti-mosquito measures and the near absence of public awareness campaigns, shortcomings partly attributed to the lack of elected union parishad leadership under the interim government. Yet Dhaka's two city corporations have spent more than BDT 1,000 crore (over USD 81 million) on mosquito-control programs in the past decade, even as the capital continues to account for the majority of infections and fatalities [111]. In 2023 alone, Dhaka recorded more than half of all cases and nearly 70 per cent of fatalities, underscoring that vector-borne outbreaks transcend partisan boundaries [112]. In FY 2024–25, Dhaka South City Corporation spent less than 40 per cent of its overall budget despite increasing its mosquito-control allocation by 19% [113]. Weak implementation, poor coordination, obsolete operational strategies, and persistent shortages of chemicals and manpower have severely undermined larviciding, mosquito-control, and drain-cleaning activities.




Figure 5. Key Drivers of Bangladesh's Rising Dengue Surge. Bangladesh's dengue surge reflects a dangerous convergence of climate stress, rapid urbanization, dense settlements, and chronic sanitation failures, which together have created ideal conditions for *Aedes* mosquitoes to flourish. Shifting infection patterns—rising male fatalities, high hospital admissions, and a disproportionate burden on young people—underscore a worsening public-health emergency driven by environmental degradation, waste mismanagement, and construction-related breeding sites. The infographic illustrates how these interconnected pressures—heat, overcrowding, poor waste disposal, irregular water supply, declining green cover, and ineffective vector control—are fueling an escalating nationwide epidemic.

Conclusions

In Bangladesh, rising temperatures, unplanned urban expansion, and worsening pollution have created conditions that strongly favor mosquito proliferation, turning rapid development into a relentless battle against one of the country's deadliest tiny predators. The persistent and evolving threat of dengue underscores the need for timely hospitalization—because the illness can deteriorate quickly—as well as systematic research to understand how environmental pollution, climate variability, and extensive pesticide use are shaping viral resistance and mosquito behavior. Media coverage has largely failed to capture the severity of the crisis, and domestic research remains limited, often attributing outbreaks only to erratic rainfall, monsoon shifts, and stagnant water.

Yet evidence from regions with similar dengue patterns points to several overlooked drivers, including air pollution, pesticide and micro plastic resistance, and the complex interactions between rapid urbanization and mosquito ecology. With low levels of health literacy, even strong research rarely translates into public awareness or policy reform, and progress in evidence-based studies remains slow. Coordinated efforts that combine early clinical care with rigorous scientific investigation are therefore essential to mitigating the country's growing dengue burden.

This national tragedy is part of a much larger global shift. A study in *Nature* warns that by 2080, nearly three in five people could be at risk of dengue [114]. Last year alone, more than fourteen million people were infected worldwide—twice the previous year and twelve times higher than a decade ago [115,116]. As climate instability, unplanned urbanization, and expanding mosquito habitats intensify, dengue is no longer a regional challenge—it is an emerging pandemic that demands urgent international action. The time to act is now, before a greater catastrophe unfolds and more lives are lost.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

Funding: None.

Institutional Review Board Statement: N/A.

Data Availability Statement: The data used in this study were obtained from the Dengue Dynamic Dashboard for Bangladesh, maintained by the Directorate General of Health Services (DGHS), Health Emergency Operation Center & Control Room. These data are publicly available at: https://dashboard.dghs.gov.bd/pages/heoc_dengue_v1.php.

Conflicts of Interest: None.

Abbreviations

BI	Breteau Index
DGHS	Directorate General of Health Services
DNCC	Dhaka North City Corporation
DSCC	Dhaka South City Corporation
IEDCR	The Institute of Epidemiology, Disease Control and Research
NS1	Nonstructural Protein 1
RAJUK	Rajdhani Unnayan Kartripakkha
WHO	World Health Organization

References

1. Jackson A. World mosquito Day 2025 - A Global Health Crisis. World Mosquito Program. 2025 Aug 11.
2. Roiz D, Pontifes PA, Jourdain F, et al. The rising global economic costs of invasive *Aedes* mosquitoes and *Aedes*-borne diseases. *Sci Total Environ*. 2024;933:173054. doi:10.1016/j.scitotenv.2024.173054
3. WHO Africa Region. Malaria. World Health Organization. 2023.
4. United Nations Office for Disaster Risk Reduction (UNDRR), International Science Council (ISC). UNDRR–ISC Hazard Information Profiles – 2025 Update: BI0207 Dengue. United Nations Office for Disaster Risk Reduction; International Science Council; 2025.
5. Ahmad LC, Gill BS, Sulaiman LH, et al. Molecular epidemiology of dengue in Southeast Asia (SEA): Protocol of systematic review and meta-analysis. *BMJ Open*. 2025;15(4):e088890. doi:10.1136/bmjopen-2024-088890
6. Weng SL, Hung FY, Li ST, et al. Dengue epidemiology in 7 Southeast Asian countries: 24-year, retrospective, multicountry ecological study. *Interact J Med Res*. 2025;14:e70491. doi:10.2196/70491
7. Bishen S. The world is in the grip of a record dengue fever outbreak. What's causing it and how can it be stopped? World Economic Forum. November 5, 2024.
8. World Health Organization. Dengue - Global situation. *Disease Outbreak News*. May 30, 2024.
9. CDC. Current dengue outbreak. Centers for Disease Control and Prevention. Updated July 29, 2025.
10. eClinicalMedicine. Dengue as a growing global health concern. 2024;77:102975. doi:10.1016/j.eclinm.2024.102975.
11. Zhang, W.-X., Zhao, T.-Y., Wang, C.-C., He, Y., Lu, H.-Z., Zhang, H.-T., Wang, L.-M., Mao, Z., Li, C.-X., & Deng, S.-Q. (2025). Assessing the global dengue burden: Incidence, mortality, and disability trends over

- three decades. *PLoS Neglected Tropical Diseases*, 19(3), e0012932. <https://doi.org/10.1371/journal.pntd.0012932>
12. WHO. Dengue fact sheet. World Health Organization. 2025 Aug 21.
 13. Wei S, Zhang T, Sun S, et al. The shift in mosquito-borne disease incidence across the Asia-Pacific region (1992–2021): Insights from an age-period-cohort analysis using the Global Burden of Disease Study 2021. *BMC Public Health*. 2025;25:3373. doi:10.1186/s12889-025-24765-y
 14. Kim JH, Lim AY, Kim SH. Evaluating the effectiveness of dengue surveillance in the tropical and sub-tropical Asian nations through dengue case data from travelers returning to the five western Pacific countries and territories. *Travel Med Infect Dis*. 2025;64:102802. doi:10.1016/j.tmaid.2025.102802
 15. Lim A, Shearer FM, Sewalk K, Pigott DM, Clarke J, Ghouse A, Judge C, Kang H, Messina JP, Kraemer MUG, Gaythorpe KAM, de Souza WM, Nsoesie EO, Celone M, Faria N, Ryan SJ, Rabe IB, Rojas DP, Hay SI, Brownstein JS, Golding N, Brady OJ. The overlapping global distribution of dengue, chikungunya, Zika and yellow fever. *Nat Commun*. 2025;16(1):3418. doi:10.1038/s41467-025-58609-5
 16. CDC. Areas with Risk of Dengue. Centers for Disease Control and Prevention; Updated December 8, 2025.
 17. Ulgheri FM, Bernardes BG, Lancellotti M. Decoding Dengue: A Global Perspective, History, Role, and Challenges. *Pathogens*. 2025;14(9):954. doi:10.3390/pathogens14090954.
 18. Morin CW, Comrie AC, Ernst K. Climate and dengue transmission: evidence and implications. *Environ Health Perspect*. 2013;121(11-12):1264-1272. doi:10.1289/ehp.1306556
 19. Fengliu F, Ma Y, Qin P, Zhao Y, Liu Z, Wang W, Cheng B. Temperature-Driven Dengue Transmission in a Changing Climate: Patterns, Trends, and Future Projections. *GeoHealth*. 2024;8(10):e2024GH001059. doi:10.1029/2024GH001059.
 20. Benedum CM, Seidahmed OME, Eltahir EAB, Markuzon N. Statistical modeling of the effect of rainfall flushing on dengue transmission in Singapore. *PLoS Negl Trop Dis*. 2018;12(12):e0006935. doi:10.1371/journal.pntd.0006935.
 21. Sophia Y, Roxy MK, Murtugudde R, et al. Dengue dynamics, predictions, and future increase under changing monsoon climate in India. *Sci Rep*. 2025;15:1637. doi:10.1038/s41598-025-85437-w.
 22. Monintja TCN, Arsin AA, Amiruddin R, Syafar M. Analysis of temperature and humidity on dengue hemorrhagic fever in Manado Municipality. *Gaceta Sanitaria*. 2021;35(Suppl):S330-S333. doi:10.1016/j.gaceta.2021.07.020.
 23. Gui H, Gwee S, Koh J, Pang J. Weather Factors Associated with Reduced Risk of Dengue Transmission in an Urbanized Tropical City. *Int J Environ Res Public Health*. 2021;19(1):339. Published 2021 Dec 29. doi:10.3390/ijerph19010339
 24. Chaiyarit J, Sriwongsuk K, Putepapas S, Intarasaksit P. Environmental health factors influencing dengue: a systematic review with thematic categorization. *Int J Environ Health Res*. 2025;1-15. doi:10.1080/09603123.2025.2589371.
 25. Nature Education. Dengue Transmission. Nature.com. Published 2014.
 26. Lozano-Fuentes S, Hayden MH, Welsh-Rodriguez C, et al. The dengue virus mosquito vector *Aedes aegypti* at high elevation in Mexico. *Am J Trop Med Hyg*. 2012;87(5):902-909. doi:10.4269/ajtmh.2012.12-0244
 27. Sieghart L, Rogers D. Bangladesh: The Challenges of Living in a Delta Country. World Bank Blogs. May 19, 2015.
 28. Islam MA, Sato T. Influence of Terrestrial Precipitation on the Variability of Extreme Sea Levels along the Coast of Bangladesh. *Water*. 2021;13(20):2915. doi:10.3390/w13202915.
 29. Miah D. Sustainable river management in Bangladesh: Challenges and ways forward. The Climate Watch. December 1, 2024.
 30. Raza W, Mahmud I, Hossain R. Bangladesh: Finding it difficult to keep cool. Washington, DC: World Bank; 2021. doi:10.1596/36534
 31. Hasan MN, Khalil I, Chowdhury MAB, et al. Two decades of endemic dengue in Bangladesh (2000–2022): trends, seasonality, and impact of temperature and rainfall patterns on transmission dynamics. *J Med Entomol*. 2024;61(2):345-353. doi:10.1093/jme/tjae001.

32. Bonna AS, Pavel SR, Mehjabin T, Ali M. Dengue in Bangladesh. *IJID One Health*. 2023;1:100001. doi:10.1016/j.ijidoh.2023.100001
33. Najmus Sakib SM. Heavy rains may double dengue cases in Oct. *The Financial Express*. October 4, 2025.
34. TBS Report. Why Bangladesh seeing so much rain in October? *The Business Standard*. October 11, 2025.
35. Islam J, Frentiu FD, Devine GJ, Bambrick H, Hu W. A state-of-the-science review of long-term predictions of climate change impacts on dengue transmission risk. *Environ Health Perspect*. 2025;133(5):56002. doi:10.1289/EHP14463
36. Rahman MS, Shiddik MAB. Explainable artificial intelligence for predicting dengue outbreaks in Bangladesh using eco-climatic triggers. *Glob Epidemiol*. 2025;10:100210. doi:10.1016/j.gloepi.2025.100210
37. Islam MT, Kamal ASMM, Islam MM, Hossain S. Time series patterns of dengue and associated climate variables in Bangladesh and Singapore (2000–2020): A comparative study of statistical models to forecast dengue cases. *Int J Environ Health Res*. 2025;35(9):2289-2299. doi:10.1080/09603123.2024.2434206
38. Hossain MS, Noman AA, Mamun SMAA, Mosabbir AA. Twenty-two years of dengue outbreaks in Bangladesh: Epidemiology, clinical spectrum, serotypes, and future disease risks. *Trop Med Health*. 2023;51(1):37. doi:10.1186/s41182-023-00528-6
39. Khan S, Akbar SM, Mahtab MA, et al. Bangladesh records persistently increased number of dengue deaths in recent years: Dissecting the shortcomings and means to resolve. *IJID Regions*. 2024;12:100395. doi:10.1016/j.ijregi.2024.100395
40. Alam KE, Ahmed MJ, Chalise R, et al. Time series analysis of dengue incidence and its association with meteorological risk factors in Bangladesh. *PLoS One*. 2025;20(8):e0323238. doi:10.1371/journal.pone.0323238
41. Islam MA, Hasan MN, Tiwari A, et al. Correlation of dengue and meteorological factors in Bangladesh: A public health concern. *Int J Environ Res Public Health*. 2023;20(6):5152. doi:10.3390/ijerph20065152
42. Islam J, Hu W. Rapid human movement and dengue transmission in Bangladesh: A spatial and temporal analysis based on different policy measures of COVID-19 pandemic and Eid festival. *Infect Dis Poverty*. 2024;13(1):99. doi:10.1186/s40249-024-01267-4
43. Ogieuhi IJ, Ahmed MM, Jamil S, et al. Dengue fever in Bangladesh: Rising trends, contributing factors, and public health implications. *Trop Dis Travel Med Vaccines*. 2025;11(1):26. doi:10.1186/s40794-025-00251-6
44. Siddique AB, Hasan M, Ahmed A, Rahman MH, Sikder MT. Youth's climate consciousness: Unraveling the dengue-climate connection in Bangladesh. *Front Public Health*. 2024;12:1346692. doi:10.3389/fpubh.2024.1346692
45. Rahman MS, Karamehic-Muratovic A, Baghbanzadeh M, et al. Climate change and dengue fever knowledge, attitudes and practices in Bangladesh: A social media-based cross-sectional survey. *Trans R Soc Trop Med Hyg*. 2020;115(1):85-93. doi:10.1093/trstmh/traa093
46. Hossain MdI, Alam NE, Akter S, et al. Knowledge, awareness and preventive practices of dengue outbreak in Bangladesh: A countrywide study. *PLoS One*. 2021;16(6):e0252852. doi:10.1371/journal.pone.0252852
47. Rahman MM, Khan SJ, Tanni KN, et al. Knowledge, attitude, and practices towards dengue fever among university students of Dhaka City, Bangladesh. *Int J Environ Res Public Health*. 2022;19(7):4023. doi:10.3390/ijerph19074023
48. Rahman MM, Tanni KN, Roy T, et al. Knowledge, attitude and practices towards dengue fever among slum dwellers: A case study in Dhaka City, Bangladesh. *Int J Public Health*. 2023;68:1605364. doi:10.3389/ijph.2023.1605364
49. Rahman MS, Amrin M, Chowdhury AH, Suwanbamrung C, Karamehic-Muratovic A. Knowledge and beliefs about climate change and emerging infectious diseases in Bangladesh: Implications for one health approach. *J Health Popul Nutr*. 2025;44(1):360. doi:10.1186/s41043-025-01112
50. Banik R, Islam MS, Mubarak M, Rahman M, Gesesew HA, Ward PR, Sikder MT. Public knowledge, belief, and preventive practices regarding dengue: Findings from a community-based survey in rural Bangladesh. *PLoS Negl Trop Dis*. 2023;17(12):e0011778. doi:10.1371/journal.pntd.0011778
51. Chowdhury NF, Haque MJ, Jahan MS, Rashid MAM, Mostafa MG, Rashid F. Knowledge, beliefs, and preventive practices regarding dengue among rural communities in Bangladesh. *KYAMC J*. 2024;15(3). doi:10.3329/kyamcj.v15i3.75223

52. Chowdhury SMMH, Rashid MA, Trisha SY, Ibrahim M, Hossen MS. Dengue investigation research in Bangladesh: Insights from a scoping review. *Health Sci Rep.* 2025;8(3):e70568. doi:10.1002/hsr2.70568
53. Pure E, Husna ALA, Rokony S, Thowai AS, Moulee ST, Jahan A, Khatun A, Sarkar M, Bibi S, Tabassum TT, Nurunnabi M. Knowledge, attitude, and practices regarding dengue infection: A community-based study in rural Cox's Bazar. *J Commun Dis.* 2025;57(1):121-130. doi:10.24321/0019.5138.202516
54. Mohiuddin AK. Dengue protection and cure: Bangladesh perspective. *Eur J Sustain Dev Res.* 2019;4(1):em0104. doi:10.29333/ejosdr/6260
55. Al-Amin HM, Johora FT, Irish SR, et al. Insecticide resistance status of *Aedes aegypti* in Bangladesh. *Parasit Vectors.* 2020;13(1):622. doi:10.1186/s13071-020-04503-6
56. Al-Amin HM, Gyawali N, Graham M, et al. Insecticide resistance compromises the control of *Aedes aegypti* in Bangladesh. *Pest Manag Sci.* 2023;79(8):2846-2861. doi:10.1002/ps.7462
57. Hossain S, Islam MdM, Hasan MdA, Chowdhury PB, Easty IA, Tusar MdK, Rashid MB, Bashar K. Association of climate factors with dengue incidence in Bangladesh, Dhaka City: A count regression approach. *Heliyon.* 2023;9(5):e16053. doi:10.1016/j.heliyon.2023.e16053
58. Paul R, Fincher C. Bangladesh sees worst single-day surge in dengue cases and deaths this year. Reuters. 2025 Sep 21.
59. Rahman A. Dengue deaths up 150%, cases double compared to last year. Bonik Barta. 2025 Sep 22.
60. UNB. 8 more dead, 778 hospitalised as Bangladesh fails to curb dengue. United News of Bangladesh. 2025 Nov 23.
61. Paul R. Dengue cases surge across Bangladesh as experts call for urgent action. Reuters. 2025 Oct 7.
62. DGHS/UNICEF. Dengue Dynamic Dashboard for Bangladesh. Health Emergency Operation Center & Control Room, Directorate General of Health Services. Accessed 2025 Dec 8.
63. News Desk. Dengue: Four more die, 920 hospitalised in 24Hrs. Daily Sun. 2025 Nov 18.
64. Asaduzzaman M, Khan EA, Hasan MN, et al. The 2023 dengue fatality in Bangladesh: Spatial and demographic insights. *IJID Regions.* 2025;15:100654. doi:10.1016/j.ijregi.2025.100654
65. Hossain MS, Noman AA, Mamun SMAA, Mosabbir AA. Twenty-two years of dengue outbreaks in Bangladesh: Epidemiology, clinical spectrum, serotypes, and future disease risks. *Trop Med Health.* 2023;51(1):37. doi:10.1186/s41182-023-00528-6
66. Barr JS, Martin LE, Tate AT, Hillyer JF. Warmer environmental temperature accelerates aging in mosquitoes, decreasing longevity and worsening infection outcomes. *Immun Ageing.* 2024;21:61. doi:10.1186/s12979-024-00465-w
67. Perdomo HD, Khorramnejad A, Cham NM, Kropf A, Sogliani D, Bonizzoni M. Prolonged exposure to heat enhances mosquito tolerance to viral infection. *Commun Biol.* 2025;8:168. doi:10.1038/s42003-025-07617-8
68. Jacobo J. Mosquitoes found in Iceland for 1st time as temperatures in the region rise. ABC News. 2025 Oct 22.
69. Galeana-Pizaña JM, Cruz-Bello GM, Caudillo-Cos CA, Jiménez-Ortega AD. Impact of deforestation and climate on spatio-temporal spread of dengue fever in Mexico. *Spat Spatio-Temporal Epidemiol.* 2024;50:100679. doi:10.1016/j.sste.2024.100679
70. Andrade AC, Falcão LA, Borges MA, Leite ME, Espírito Santo MM. Are land use and cover changes and socioeconomic factors associated with the occurrence of dengue fever? A case study in Minas Gerais State, Brazil. *Resources.* 2024;13(3):38. doi:10.3390/resources13030038
71. Chaves LS, Conn JE, López RV, Sallum MA. Abundance of impacted forest patches <5 km² is a key driver of the incidence of malaria in Amazonian Brazil. *Sci Rep.* 2018;8(1):7077. doi:10.1038/s41598-018-25344-5
72. Nawar N, Sorker R, Chowdhury FJ, Mostafizur Rahman Md. Present status and historical changes of urban green space in Dhaka City, Bangladesh: A remote sensing driven approach. *Environ Chall.* 2022;6:100425. doi:10.1016/j.envc.2021.100425
73. IIED. Hot Cities: Dhaka. International Institute for Environment and Development, London. 2024 Jun.
74. World Bank. Bangladesh faces health and economic risks from rising temperature. Press Release. 2025 Sep 16.
75. Banu S, Hu W, Guo Y, Hurst C, Tong S. Projecting the impact of climate change on dengue transmission in Dhaka, Bangladesh. *Environ Int.* 2014;63:137-142. doi:10.1016/j.envint.2013.11.002

76. Quam J, Campbell S. South Asia: Urban Geography I – Dhaka. In: *The Eastern World: Daily Readings on Geography*. College of DuPage Digital Press; 2022.
77. UNB. Dhaka world's 2nd largest city with 36.6 million: UN. *The Daily Star*. 2025 Nov 26.
78. Sultana A, Islam A, Hosna A, Tahsin A, Islam A. The impact of urbanization on the proliferation of *Aedes aegypti* (Diptera: Culicidae) mosquito population in Dhaka Mega City, Bangladesh. *Bangladesh J Zool*. 2024;52(2):201-215. doi:10.3329/bjz.v52i2.77460
79. Kamal AS, Al-Montakim MdN, Hasan MdA, et al. Relationship between urban environmental components and dengue prevalence in Dhaka City – An approach of spatial analysis of satellite remote sensing, hydro-climatic, and census dengue data. *Int J Environ Res Public Health*. 2023;20(5):3858. doi:10.3390/ijerph20053858
80. Roy S, Biswas A, Shawon MT, Akter S, Rahman MM. Land use and meteorological influences on dengue transmission dynamics in Dhaka City, Bangladesh. *Bull Natl Res Cent*. 2024;48:32. doi:10.1186/s42269-024-01188-0
81. Inspira Advisory and Consulting Limited. Challenges of slum living in Bangladesh: A closer look at WASH inequities in Bangladesh's slums. *Inspira-bd.com*. 2023 Jul 17.
82. Paulson W, Kodali NK, Balasubramani K, et al. Social and housing indicators of dengue and chikungunya in Indian adults aged 45 and above: Analysis of a nationally representative survey (2017–18). *Arch Public Health*. 2022;80(1):125. doi:10.1186/s13690-022-00868-5
83. UNICEF. 230 tons of fecal waste end up in open water bodies in Dhaka daily – UNICEF and WaterAid call for stronger sanitation management. *UNICEF Bangladesh*. 2025 Feb 25.
84. Alam HMN. Dhaka's drains, dengue, and denial. *The Daily Star*. 2025 Jul 10.
85. Islam J, Asif MH, Rahman S, Hasan M. Exploring mosquito hazards in Bangladesh: Challenges and sustainable solutions. *IUBAT Rev*. 2024;7(2):1–29. doi:10.3329/iubatr.v7i2.78792
86. Nature India. Poor access to tap water linked to dengue risk. 2021. doi:10.1038/nindia.2021.27
87. UNB. A roundtable discussion on 'Solid Waste Management – Challenges and Solutions for Bangladesh'. *United Nations Bangladesh*. 2024 Oct 3.
88. Sasi MS, Lal N. The impact of solid waste management practices on vector-borne disease risk in Thiruvananthapuram. *Int J Multidiscip Res*. 2024;6(4):1-10. doi:10.36948/ijfmr.2024.v06i04.26724
89. Rosser JI, Openshaw JJ, Lin A, et al. Seroprevalence, incidence estimates, and environmental risk factors for dengue, chikungunya, and Zika infection amongst children living in informal urban settlements in Indonesia and Fiji. *BMC Infect Dis*. 2025;25(1):51. doi:10.1186/s12879-024-10315-1
90. World Health Organization. Climate change, pollution and health: Impact of chemicals, waste and pollution on human health. Executive Board EB154/24. Geneva: WHO; 2023 Dec 18.
91. Lu HC, Lin FY, Huang YH, Kao YT, Loh EW. Role of air pollutants in dengue fever incidence: Evidence from two southern cities in Taiwan. *Pathog Glob Health*. 2022;117(6):596-604. doi:10.1080/20477724.2022.2135711
92. Mailepessov D, Ong J, Aik J. Influence of air pollution and climate variability on dengue in Singapore: A time-series analysis. *Sci Rep*. 2025;15(1):13467. doi:10.1038/s41598-025-97068-2
93. Ju X, Zhang W, Yimaer W, et al. How air pollution altered the association of meteorological exposures and the incidence of dengue fever. *Environ Res Lett*. 2022;17(12):124041. doi:10.1088/1748-9326/aca59f
94. Thongtip S, Sapbamrer P, Chaichanan P, Chiablam S, Pimonsree S. Association of meteorology and air quality with dengue fever incidence in upper northern Thailand. *EnvironmentAsia*. 2025;18(1):164-173. doi:10.14456/ea.2025.13
95. Mohammad AKH, Che Dom N, Do Camalxaman S, Syed Ismail SN. Correlational analysis of air pollution index levels on dengue surveillance data: A retrospective study in Melaka, Malaysia. *J Sustain Sci Manag*. 2020;15(8):1-9. doi:10.46754/jssm.2020.12.012
96. Carneiro MAF, Alves BCA, Gehrke FS, et al. Environmental factors can influence dengue reported cases. *Rev Assoc Med Bras*. 2017;63(11):957-961. doi:10.1590/1806-9282.63.11.957
97. Tewari P, Ma P, Gan G, et al. Non-linear associations between meteorological factors, ambient air pollutants and major mosquito-borne diseases in Thailand. *PLoS Negl Trop Dis*. 2023;17(12):e0011763. doi:10.1371/journal.pntd.0011763

98. Maquart P-O, Froehlich Y, Boyer S. Plastic pollution and infectious diseases. *Lancet Planet Health*. 2022;6(10):e842-e845. doi:10.1016/s2542-5196(22)00198-x
99. Afroze CA, Ahmed MN, Azam MN, Jahan R, Rahman H. Microplastics pollution in Bangladesh: A decade of challenges, impacts, and pathways to sustainability. *Integr Environ Assess Manag*. 2025 Aug 11. doi:10.1093/inteam/vjaf108
100. Meijer LJJ, van Emmerik T, van der Ent R, Schmidt C, Lebreton L. More than 1000 rivers account for 80% of global riverine plastic emissions into the ocean. *Sci Adv*. 2021;7(18):eaaz5803. doi:10.1126/sciadv.aaz5803
101. Environment and Social Development Organization (ESDO). Huge use of poly bag: 78 thousand tons of waste in a year. ESDO. 2021 Jun 5.
102. Chowdhury SI. Urban per capita plastic use 9 kg, 24 kg in Dhaka. *New Age*. 2021 Dec 20.
103. Li JH, Liu XH, Liang GR, et al. Microplastics affect mosquito from aquatic to terrestrial lifestyles and are transferred to mammals through mosquito bites. *Sci Total Environ*. 2024;917:170547. doi:10.1016/j.scitotenv.2024.170547
104. Shopon HU-R. Dhaka: Unplanned city faces a grand spectacle of risks (Article in Bengali). *Deutsche Welle*. 2025 Jul 23.
105. Hassan A. Building faults overlooked if officials are appeased. *Prothomalo*. 2024 Mar 9.
106. Tribune Desk. Potential Aedes breeding grounds found in 70% DNCC homes. *Dhaka Tribune*. 2020 Jul 4.
107. Staff Correspondent. Greetings and promises on our 15th anniversary. Aedes reproduction--High rises mainly responsible. *Daily Sun*. 2021 May 6.
108. TBS Report. Construction work will be halted if Aedes larvae found on site: Mayor Taposh. *The Business Standard*. 2024 Apr 25.
109. Staff Correspondent. Dengue infection: 13 Dhaka wards at high risk. *The Daily Star*. 2025 Jun 19.
110. Kamal M, Sultana R, Julkarnayeen M. Dengue crisis prevention and control: Governance challenges and way forward. *Transparency International Bangladesh*. 2023 Oct 30.
111. Islam MdJ. Dengue rages as TK1,000CR lost to futile mosquito control efforts. *The Business Standard*. 2025 Nov 16.
112. Hossain M, Rakib MS, Hasan MM, Powshi SN, Islam E, Islam NN. The 2023 dengue outbreak in Bangladesh: An epidemiological update. *Health Sci Rep*. 2025;8(5):e70852. doi:10.1002/hsr2.70852
113. TBS Report. Dhaka south increases mosquito control budget amid rising dengue infections, reports revenue growth. *The Business Standard*. 2025 Aug 6.
114. Messina JP, Brady OJ, Golding N, et al. The current and future global distribution and population at risk of dengue. *Nat Microbiol*. 2019;4(9):1508-1515. doi:10.1038/s41564-019-0476-8
115. Haider N, Hasan MN, Onyango J, et al. Global dengue epidemic worsens with record 14 million cases and 9000 deaths reported in 2024. *Int J Infect Dis*. 2025;158:107940. doi:10.1016/j.ijid.2025.107940
116. US CDC. Dengue on the rise: Get the facts. *Centers for Disease Control and Prevention*. 2025 May 29.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.