

Review

Not peer-reviewed version

---

# Autophagy: A Double-Edged Sword in the Aging of *C. elegans*

---

[Tímea Sigmond](#)<sup>\*</sup> and [János Barna](#)

Posted Date: 13 February 2026

doi: 10.20944/preprints202602.0649.v2

Keywords: autophagy; aging; *Caenorhabditis elegans*; lifespan regulation; proteostasis; stress response



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

# Autophagy: A Double-Edged Sword in the Aging of *C. elegans*

Tímea Sigmond <sup>1,\*</sup> and János Barna <sup>1,2,3</sup>

<sup>1</sup> Department of Genetics, Institute of Biology, Eötvös Loránd University, Budapest, Hungary

<sup>2</sup> HUN-REN-ELTE Genetics Research Group, Eötvös Loránd University, Budapest, Hungary

<sup>3</sup> Food and Wine Research Institute, Eszterházy Károly Catholic University, Eger, Hungary

\* Correspondence: sigmond.timea@ttk.elte.hu

## Abstract

Autophagy is a tightly regulated catabolic process essential for cellular homeostasis, stress adaptation, and regeneration. In the nematode *Caenorhabditis elegans*, with its short lifespan, transparent body, and well-defined genetics, the process can be investigated at tissue- and age-specific manner, making it an excellent model to study the connection between autophagy and longevity. While autophagy is indispensable for development and homeostasis, recent studies have revealed that its role in aging is more complex than previously thought. During post-reproductive life, autophagic flux and the degradative capacity of lysosomes decline, resulting in the accumulation of undegraded material and cellular stress. Several studies have demonstrated that the experimental modulation of core autophagy in aged or post-reproductive *C. elegans*, particularly in neurons, can improve proteostasis, preserve tissue integrity, and extend lifespan. Here we review the current results obtained using the genetic model system *Caenorhabditis elegans* that link autophagy to lifespan regulation. We focus on studies that investigate unexpected, context-dependent, or deleterious effects of inhibiting autophagy-related genes during aging. We also discuss how age- and tissue-specific modulation of autophagy could define the most effective strategies for promoting healthy aging. This could provide relevant insights for the therapeutic targeting of autophagy in humans.

**Keywords:** autophagy; aging; *Caenorhabditis elegans*; lifespan regulation; proteostasis; stress response

## 1. Introduction

Macroautophagy, or autophagy, is a conserved catabolic process in eukaryotic cells that delivers misfolded proteins, aggregates, and defective organelles to lysosomes via double-membraned autophagosomes for degradation and recycling [1]. Autophagy occurs at a basal level in most tissues under normal physiological conditions and is rapidly upregulated in response to various intra- and extracellular stressors such as nutrient deprivation, or environmental stress [2]. Dysfunctional autophagy or impaired lysosomal breakdown has been implicated in a wide range of diseases, including inflammation, cardiovascular disorders, metabolic diseases such as diabetes, cancer, and neurodegeneration. Beyond disease, autophagy is also recognized as a critical regulator of organismal lifespan [3].


This conserved cellular machinery operates via well-defined molecular steps. Upon induction, a double-membraned structure called the phagophore (also referred to as the isolation membrane or pre-autophagosomal membrane) forms at multiple sites of the cytoplasm [4,5]. The membrane elongates and engulfs the targeted cytoplasmic components, resulting in autophagosome formation. This structure then fuses with a lysosome to form an autolysosome, where the sequestered material is degraded by acid hydrolases and recycled [6]. Bulk autophagy is a non-selective process that provides the cell with essential macromolecules, especially under nutrient-deprived conditions [7]. In contrast, selective autophagy, such as mitophagy, aggrephagy, or lipophagy involves cargo

recognition by specific receptors and plays a key role in maintaining cellular quality control and homeostasis [8].

Given its short lifespan, genetic tractability, and conserved autophagy machinery, *Caenorhabditis elegans* has emerged as a powerful in vivo model to dissect the role of autophagy in lifespan regulation [9]. Genetic screens in *C. elegans* identified several metazoan-specific autophagy genes (e.g. *epg-2*, *epg-3*, *epg-4*, and *epg-5*), expanding our understanding beyond yeast models [10]. Studies in this organism have also revealed not only the longevity-promoting functions of autophagy, but also unexpected age- and tissue-specific detrimental effects, providing a unique framework to understand the context-dependent nature of autophagy during aging [11–14].

## 2. *C. elegans* as a Model Organism to Study Autophagy

Autophagy is broadly conserved: many mammalian autophagy-related proteins have a single ortholog in *C. elegans*, and genetic studies support the existence of a conserved core machinery [15]. However, some of these components have not yet been characterized experimentally in *C. elegans*, so their roles are inferred solely in their homology with orthologs in other organisms. The core molecular machinery of macroautophagy has been thoroughly reviewed elsewhere [16]; therefore, its mechanism is summarized only schematically in Figure 1.



**Figure 1.** Mechanism and regulation of macroautophagy in *Caenorhabditis elegans*. Autophagy is primarily regulated by the nutrient-sensing kinases mTOR (LET-363) and AMPK (AAK-2). Upon induction, portions of the cytoplasm are sequestered within a double-layered isolation membrane. Closure of this membrane generates an autophagosome, which subsequently matures and fuses with acidic lysosomes to form autolysosomes, where cytoplasmic material is degraded. This process is orchestrated in a hierarchical manner by autophagy-related proteins that assemble into functional complexes, including initiation, nucleation, PI3P-binding, and conjugation systems. Membrane expansion is supported by ATG9-containing vesicles, and additional factors regulate vesicle trafficking and membrane fusion. Transcription of most autophagy-related genes (*atg*) is governed by stress-responsive transcription factors such as HLH-30/TFEB, DAF-16/FOXO, and PHA-4/FOXA.

Several forms of selective autophagy have been described in *C. elegans*, including aggrephagy, which is mediated by the adaptor protein SQST-1/p62, and mitophagy, which is mediated by the

protein DCT-1/BNIP3. These findings highlight the fact that cargo selectivity adds another layer to autophagy-aging interactions [17,18].

Key signaling pathways and processes involved in longevity regulate autophagy in worms. The LET-363/TOR (target of rapamycin) pathway coordinates nutrient sensing, protein synthesis, and autophagy [19,20]. The insulin/IGF-1 pathway influences metabolism, stress resistance, and lifespan [21]. Under conditions of mitochondrial stress or mitochondrial dysfunction, autophagy can be upregulated, emphasizing the integration of stress responses and recycling systems [22]. Dietary restriction (DR) robustly induces autophagy via multiple conserved mechanisms including the suppression of the LET-363/TOR and the insulin/IGF-1 signaling, sirtuin activation, epigenetic modulation, and mitochondrial remodeling [23]. In *C. elegans*, eating-defective mutants, such as *eat-2*, provide a genetic model of dietary restriction [24]. Increased autophagic activity has been observed in these mutants, particularly in intestinal tissues. These findings support a role for autophagy in mediating DR-induced longevity [25]. The helix-loop-helix transcription factor HLH-30, an ortholog of mammalian TFEB, is a central transcriptional regulator of autophagy-related genes in *C. elegans*. Under conditions such as starvation or infection, HLH-30 translocates into the nucleus, where it activates the expression of genes involved in autophagy and lysosomal function [24,26]. Furthermore, the transcription factor PHA-4/FOXA acts in the adult stages to regulate the expression of autophagy-related genes, especially under dietary restriction or TOR suppression [25]. Autophagy gene regulation is also modulated by additional pathways including TGF- $\beta$  (Sma/Mab), Rb (*lin-35*), the unfolded protein response (XBP-1), and mitochondrial stress response (ATFS-1) [27].

To monitor autophagy dynamics in vivo, the transparent *C. elegans* offers robust molecular tools such as the endogenously inserted *gfp::mCherry::lgg-1* reporter, which enables reliable and artifact-free visualization of autophagy progression. This dual-fluorescent system reveals life stage- and stress-dependent regulation of autophagic flux [28].

In addition to mutational inactivation, silencing of autophagy genes is also highly accessible in this animal model. RNA interference (RNAi) can be induced by feeding animals bacteria expressing double-stranded RNA that target specific genes [29]. Sensitivity to RNAi can be enhanced in *rrf-3* mutant backgrounds [30], whereas tissue-specific knockdown is achieved using *sid-1* mutants, defective in systemic RNA interference, complemented with transgenic, cell-type-specific expression of the SID-1 dsRNA receptor [13,31,32]. The auxin-inducible degradation (AID) system provides a recently developed approach in *C. elegans* for conditional, tissue-specific and temporally controlled protein depletion, complementing genetic and RNAi-based perturbations [33]. This ability to monitor and manipulate autophagy in a tissue-specific and temporally resolved manner grants powerful opportunities to elucidate how autophagy contributes to organismal aging.

*C. elegans* has also become a cornerstone of aging research due to its short lifespan and conserved aging hallmarks, including muscle deterioration, intestinal and gonadal atrophy, lipid accumulation, and loss of proteostasis and mobility; features that parallel human aging [34].

Taken together, *C. elegans* offers an unexampled model for studying autophagy contributions in lifespan regulation in vivo because of its short lifespan, genetic tractability, conserved autophagy machinery, and powerful tools for tissue- and age-specific analyses.

### 3. Autophagy as a Conserved Driver of Lifespan Extension in *C. elegans*

Since aging is considered a genetically regulated process, *C. elegans* has become an effective model for investigating the molecular mechanisms underlying lifespan regulation. The first long-lived mutant identified carried the *daf-2(e1370)* allele, which encodes a partial loss-of-function variant of the insulin/IGF-1 receptor [21]. Meléndez and colleagues demonstrated that the autophagy gene *bec-1* (homologous to BECN1/ATG6) is essential for the extended lifespan observed in *daf-2* mutants [35]. This finding was later supported by studies showing that the inactivation of *atg-18*, or the RNAi-mediated silencing of *atg-7* and *atg-12* partially suppresses the longevity phenotype of *daf-2* mutants Table 1.

Autophagy is required for lifespan extension across multiple genetically and environmentally induced longevity pathways in *C. elegans*. Beyond reduced insulin/IGF-1 signaling, dietary restriction, TOR inhibition, and mild mitochondrial dysfunction all extend lifespan in an autophagy-dependent manner. For example, lifespan extension in *let-363/TOR* mutants, *eat-2* dietary-restriction models, and mitochondrial mutants such as *clk-1* or *atp-3* are suppressed by genetic or RNAi-mediated inhibition of core autophagy genes including *bec-1*, *unc-51*, and *atg-18* [11,18,20,25,36,37]. Collectively, the increased autophagic activity is not merely correlated with extended lifespan but is functionally required for longevity across a broad spectrum of long-lived *C. elegans* models Table 1.

Interestingly, many of the cytoprotective mechanisms of autophagy are concentrated in the intestine. Intestine-specific overexpression of the Kruppel-like transcription factor *klf-3*, which regulates cell proliferation and metabolic homeostasis, promotes autophagy by upregulating autophagy-related genes and thereby extends lifespan [38]. In addition, the paracaspase MALT-1, known for its proinflammatory role in mammals, was recently identified as a negative regulator of the early steps of autophagy in *C. elegans*. Loss of *malt-1* enhances lifespan via increased autophagic activity in the intestine, an effect reversed by *atg-13*, *bec-1*, and *lgg-2* knockdown Table 1 [39].

Notably, elevated autophagic activity in the intestine leads to the formation of tubular lysosomes (TLs), a specialized lysosomal network, necessary for the lifespan extension of *eat-2* mutants. Although TL formation alone is not sufficient to extend lifespan, their abundance strongly correlates with improved healthspan. This suggests that TLs play a role in maintaining cellular homeostasis during aging [40,41]. Moreover, pexophagy, the selective degradation of peroxisomes, becomes more active in the intestine of young adults and starved worms. In *prx-11* mutants, where this process is impaired, lifespan is markedly shortened, underscoring the physiological relevance of peroxisome turnover [42]. These findings emphasize that the role of autophagy is highly tissue-dependent and must be studied with appropriate tissue-specific resolution.

Beyond endogenous regulatory circuits, autophagy can be directly modulated through targeted genetic manipulation. Overexpression of the transcription factor HLH-30/TFEB and the selective autophagy receptor SQST-1/p62 has been shown to extend the lifespan of *C. elegans* by enhancing autophagy and improving protein homeostasis Table 1 [17,24,43,44]. In addition, external environmental factors, such as low temperature conditions (under 15 °C) [45], dietary exposure to specific amino acids [46], and in response to stress modulated by microRNA-9 [47], can dynamically modulate autophagy and thereby influence lifespan in *C. elegans*. Mild, transient heat stress during early adulthood can activate the heat-shock response via HSF-1 and stimulate HLH-30/TFEB-dependent autophagy, restoring proteostasis and enhancing lifespan [43]. This hormetic effect exemplifies how temporal stress can beneficially modulate autophagy and counteract age-related proteotoxic damage and illustrating that maintaining optimal autophagic flux is essential for sustaining proteostasis and delaying the aging process [48–51].

**Table 1.** Beneficial and context-dependent effects of autophagy on lifespan regulation in *Caenorhabditis elegans*. The table summarizes autophagy-modulating conditions, regulators, tissue specificity, and animals' age.

Conditions	Activation trigger category	Animal age	Autophagy gene studied	Inactivation/overexpression	Tissue	Ref.
<b>Beneficial effects of autophagy</b>						
TOR pathway inhibition	Endogene activation	Early adulthood	<i>bec-1, unc-51, atg-18</i>	Mutants/RNAi <sup>1</sup>	Multiple tissues	[11,20,54]
Reduced insulin/IGF-1 signaling	Endogene activation	Early adulthood	<i>bec-1, atg-7, lgg-1</i>	Mutants/RNAi <sup>1</sup>	Multiple tissues	[11,26,35]
Mitochondrial dysfunction	Mitochondrial stress	Early adulthood	<i>unc-51, bec-1, atg-18, dct-1</i>	Mutants/RNAi <sup>1</sup>	Multiple tissues	[11,18]

Dietary restriction	Mild stress activation	Early adulthood	Multiple autophagy genes	Mutants/RNAi <sup>1</sup>	Intestine	[11,37]
MALT-1 inhibition	Direct autophagy activation	Early adulthood	<i>hlh-30</i>	Mutants	Intestine	[39]
TFEB/HLH-30 overexpression	Direct transcriptional activation	Early adulthood	<i>hlh-30</i>	Overexpression	Intestine	[44]
SQST-1/p62 overexpression	Activation of selective autophagy	Early adulthood	<i>sqst-1</i>	Overexpression	Intestine, neurons	[24]
Normal homeostasis	Neutral / homeostatic	Lifelong	Core autophagy genes	RNAi <sup>1</sup>	Multiple tissues	[11,52]
<b>Prolonged stress induced overexpression of autophagy</b>						
Proteotoxicity in <i>mec-4(d)</i> mutants	Proteotoxic stress	Adulthood	<i>bec-1, atg-7</i>	RNAi <sup>1</sup>	Neurons	[53]
Constitutive MPK-1 activation	Starvation	Early adulthood	<i>bec-1</i>	RNAi <sup>1</sup>	Pharynx	[54]
<i>sgk-1</i> inhibition	mPTP activation	Adulthood	<i>bec-1, atg-16</i>	RNAi <sup>1</sup>	Intestine	[55]
High glucose diet	Metabolic stress	Adulthood	<i>hlh-30</i>	RNAi <sup>1</sup>	Intestine	[58]
<b>Age-dependent negative effects of autophagy</b>						
Aging-related dysregulation	Neutral / homeostatic	Postreproductive	<i>unc-51, bec-1, atg-7, atg-9</i>	RNAi <sup>1</sup>	Multiple tissues	[62]
Aging-related dysregulation	Proteotoxic stress with aging	Postreproductive	<i>atg-7, bec-1</i>	Tissue-specific RNAi	Neurons	[13]
Aging-related hyperfunction	Excessive autophagy with aging	Postreproductive	<i>bec-1, atg-13, atg-18</i>	RNAi <sup>1</sup>	Intestine	[14]
Cellular toxicity	Progressive lysosomal overload	Late adulthood	Multiple autophagy genes	Mutants	Multiple tissues	[76]

<sup>1</sup>It should be noted that systemic RNAi in *C. elegans* is typically inefficient in neurons because of limited dsRNA transport and processing, which may affect conclusions regarding neuronal autophagy functions.

#### 4. Dysregulation of Autophagy Can Lead to Disadvantageous Effects in *C. elegans*

While autophagy is crucial for cytoprotection under hormetic stress [43], increasing evidence suggests that persistent overactivation can lead to deleterious effects, including cellular damage or death [49,52,53].

Toxic mutations affecting MEC-4, DEG-1, and DEG-3 ion channels induce neurodegeneration through excessive cellular stress and excessive autophagy. Genetic inhibition of key autophagy genes, such as *unc-51*, *bec-1*, and *lgg-1*, suppresses neuronal death, demonstrating that autophagy was found not protective but instead actively contributes to neuronal degeneration [54]. Besides, the mitogen-activated protein kinase MPK-1/MAPK functions in the pharyngeal muscles of *C. elegans*

during starvation [19]. In *mpk-1* mutants, both autophagy levels and lifespan are diminished during regeneration after refeeding; however, MPK-1 direct role in modulating autophagy is not proved. Still, constitutive activation of MPK-1 is associated with excessive autophagy and pharyngeal tissue damage, leading to lethality [55]. Moreover, prolonged opening of the mitochondrial permeability transition pore (mPTP) also leads to mitochondrial dysfunction and cell death [56]. In *C. elegans* lacking SGK-1, main component in the regulation of the cellular ionhomeostasis [57], Zhou et al. (2019) showed that elevated autophagy, in the context of impaired mitochondrial homeostasis and increased mPTP opening, contributes to a shortened lifespan. Suppression of autophagy via RNAi (e.g. *bec-1*, *lgg-1*, and *unc-51*) can restore or partially rescue this phenomenon [56]. Furthermore, under a high-glucose diet, HLH-30-dependent autophagy is overactivated, leading to lipid dysregulation and reduced lifespan. Inhibition of HLH-30 suppresses autophagy and restores longevity Table 1 [58].

Taken together, that chronic autophagy in diverse tissues and stress contexts can lead to the accumulation of cellular damage or even necrotic cell death.

## 5. Antagonistic Pleiotropy and Age-Dependent Reversal of Autophagy Function

Several lines of evidence suggest that autophagy genes can negatively affect the lifespan of *C. elegans*. Loss of *epg-6* (a paralog of *atg-18*) extends lifespan, in contrast to the shortened lifespan observed in *atg-18* mutants, despite both genes contributing to the early steps of autophagosome formation [59]. EPG-6 functions as a PI3P effector that regulates the progression from omegasomes to autophagosomes [60,61], whereas ATG-18 plays broader, pleiotropic roles, including metabolic regulation and dauer fat accumulation [59]. These observations reflect functional differences between distinct autophagy-related genes. Several subsequent studies demonstrated that even the same autophagy genes can exert opposing effects on lifespan depending on the timing of their inactivation.

The first direct evidence demonstrating that autophagy genes with an overall protecting role can become detrimental later in life was provided by Hashimoto et al. (2009) [62]. Although lifelong autophagy inactivation reduces lifespan, RNAi silencing of certain autophagy genes, such as *unc-51*, *bec-1*, *atg-7*, and *atg-9*, in young adults slightly enhances it [13,62,63]. This effect is independent of longevity pathways, such as dietary restriction, insulin/IGF-1 signaling, and reduced mitochondrial respiration. The TOR signaling pathway is suggested to contribute to this lifespan extension partially upon adult-specific autophagy gene silencing Table 1 [62].

These observations raised an important mechanistic question: how can inhibition of the same process reduce lifespan in one context, yet extend it in another? One possible mechanism involves age-related hyperfunction of intestinal autophagy. In *C. elegans*, yolk proteins are synthesized by autophagy conversion of intestinal tissue, partly regulated by the transcription factor PHA-4 [64]. Mechanistically, this reflects a failure to downregulate reproduction-associated autophagy after reproductive maturity. The excessive vitellogenesis drives progressive intestinal atrophy and accumulation of pseudocoelomic lipoprotein pools (PLPs) [14]. Importantly, post-reproductive inhibition of autophagy by RNA interference targeting early acting autophagy genes (e.g., *atg-13*, *bec-1*), suppresses yolk overproduction, alleviates intestinal degeneration, and significantly extends lifespan [14]. These findings demonstrate that late-life intestinal autophagy is a detrimental process rather than a protective recycling mechanism. This supports the theory of antagonistic pleiotropy, which states that mechanisms that enhance early-life reproductive fitness can also drive late-life degeneration [13]. Notably, the antagonistic effects of autophagy are not restricted to the intestine Table 1.

Furthermore, neuron-specific knockdown of core autophagy genes, such as *atg-7* and *bec-1*, in adult *C. elegans* was also shown to extend lifespan by up to ~30% [13]. The inhibition of neuronal autophagy improved motor functions and maintained muscle structure in a non-cell-autonomous manner Table 1. These findings suggest that, in late life, sustained neuronal autophagy may represent a maladaptive process whose attenuation promotes longevity and functional health, highlighting the

tissue- and age-specific antagonistic roles of autophagy with potential relevance for neurodegenerative disease intervention [13].

## 6. Spatiotemporal and Compensatory Regulation of Autophagy During Aging

Building on the observations described above, autophagy emerges as a process with strongly context-dependent effects on aging. In the nematode *Caenorhabditis elegans* autophagy supports development and cellular homeostasis; however, its chronic dysregulation, whether by overactivation or inhibition, can lead to cellular dysfunction, atrophy, tissue degeneration, and reduced lifespan [12,65–67]. Lifespan optimization likely depends on the spatiotemporal control of autophagy. Inhibition of the early steps of autophagy in older animals has been shown to improve overall health and longevity [12,65–67]. However, targeting pleiotropic autophagy genes that play roles in multiple cellular processes, such as *lgg-1* [68,69] can also have negative effects [14].

Among the different tissues, the intestine emerges as a central hub for coordinating systemic aging processes in *C. elegans*. As a multifunctional organ responsible for metabolism, innate immunity, and stress responses, the intestine integrates various longevity pathways [70]. Several long-lived mutants, including those affecting insulin signaling, dietary restriction, or mitochondrial respiration, require intestinal autophagy for lifespan extension [37]. Age-related intestinal pathologies, such as yolk overproduction and lipoprotein accumulation, further highlight the gut as a primary site of aging-related deterioration [14]. This observation naturally leads to the concept that tissue-specific modulation of autophagy, especially in the intestine, may be a promising strategy for enhancing longevity in *C. elegans*.

In certain contexts, autophagy inhibition triggers compensatory mechanisms to preserve proteostasis and organismal health, suggesting the existence of backup systems that safeguard longevity. For example, neurons under proteotoxic or mitochondrial stress, and impaired lysosomal activity increase exopher biogenesis, extruding aggregates and organelles from the cell, which correlates with better neuronal function [64]. Suppression of autophagy genes acting in early steps of the process, such as *unc-51*, *atg-13*, *atg-7*, *atg-4.1*, and *lgg-1*, leads to enhanced exopher biogenesis in neurons. These exophers facilitate the removal of toxic protein aggregates, including aggregation-prone proteins such as polyglutamine-containing species, thereby reducing proteotoxic stress and contributing to improved neuronal function and longevity [71].

Besides neurons, pharyngeal muscles also exhibit a protective ‘safety mechanism’ that suppresses protein aggregation when core degradation pathways are compromised [72]. Chaperone-mediated autophagy (CMA) has recently been identified in *C. elegans* and shows strong cross-regulation with macroautophagy under starvation conditions in different cell-types [73]. However, other mechanisms, such as the canonical endosomal microautophagy have not been established in *C. elegans* [74]. Compensatory activation of alternative clearance mechanisms may reflect another adaptive response to declining autophagic efficiency with age, mitigating the deleterious effects of persistent or dysfunctional autophagy.

Declining lysosomal capacity, particularly in neuronal cells, has profound consequences for cellular homeostasis and is closely linked to the development of neurodegenerative diseases. In aged or chronically stressed cells, lysosomal function becomes compromised, characterized by impaired acidification, defective transport, and reduced fusion of autophagosomes and endosomes with lysosomes. Consequently, autophagic flux is disrupted, leading to the accumulation of protein aggregates, damaged organelles, and other toxic cellular components [75]. Recent work by Murley et al. demonstrated that macroautophagy can itself induce lysosomal damage even in the young, arrested larvae that impedes quiescent cell re-entry, highlighting a context-dependent detrimental role of autophagy beyond its canonical protective functions. This confirms the notion that autophagy’s impact on organismal health and aging depends critically on temporal and cellular state [76]. These findings highlight that the effects of autophagy modulation on lifespan are highly context-dependent and shaped by compensatory pathways, tissue identity, and stress conditions.

Zhong et al. (2025) recently demonstrated that in *C. elegans* neurons, HLH-30 actively expands lysosomal degradative capacity in early adulthood, and that this capacity declines with age. Loss of HLH-30 leads to impaired proteostasis and accelerated dendritic degeneration, highlighting lysosomal dysfunction as a key driver of age-associated neuronal decline [77]. These results support the notion that preserving lysosomal competence, rather than uniformly enhancing autophagy, is critical for maintaining tissue health during aging.

It is worth noting that most of the findings mentioned in this review were derived primarily from lifespan measurements. Although lifespan often correlates with healthspan, it does not always accurately capture the functional and physiological aspects of aging. In *C. elegans*, several well-established approaches enable direct assessment of healthspan, including the quantification of lipofuscin accumulation as a marker of cellular aging, measurements of locomotion and paralysis, pharyngeal pumping assays, and behavioral or neuromuscular tests that monitor tissue functionality [78–80]. Incorporating such parameters can refine our understanding of how autophagy influences not only lifespan but also the quality of aging in this model.

## 7. Relevance to Human Disease

The dual role of autophagy has also been observed in human diseases. The metaphor of the “double-edged sword” refers to the paradoxical role of autophagy in cancer, acting both as a tumor suppressor and a survival mechanism for malignant cells [81–83]. Similar contradictory roles have been found in liver disorders, muscle degeneration, and infectious diseases [84]. In neurodegenerative disorders including Alzheimer’s and Parkinson’s diseases, defects in autophagosome–lysosome fusion and the accumulation of autolysosomal intermediates represent key pathogenic drivers, mirroring the age-related decoupling of autophagic flux described in *C. elegans* [75,85]. One therapeutic strategy involves activating autophagy precisely at the point of blockage using small molecules [86–90], while another considers the directed inhibition of autophagy to treat age-related diseases.

Accumulating evidence from mammalian model systems indicates that autophagy is not universally protective and can exert detrimental effects under specific pathological conditions. In models of metabolic and cardiovascular disease, excessive or dysregulated autophagy has been shown to promote tissue damage rather than confer protection. For example, in diabetic cardiomyopathy, hyperactivation of autophagy contributes to cardiomyocyte dysfunction and ventricular remodeling, while pharmacological inhibition of autophagy ameliorates disease progression [91]. Similarly, in pancreatic  $\beta$  cells, excessive autophagy and ferritinophagy drive ferroptotic cell death and metabolic dysfunction [92]. Moreover, recent studies in the central nervous system demonstrate that hyperactivation of secretory autophagy promotes neuroinflammation and neurodegeneration by enhancing cytokine release and pyroptotic cell death, contributing to Alzheimer’s disease pathology [93]. Together, these findings indicate that, as observed in *C. elegans*, autophagy in mammals can shift from a cytoprotective to a pathogenic process depending on cellular context, stress intensity, and lysosomal competence.

Human studies also support the concept that autophagy dysfunction is highly tissue-dependent. Impaired autophagy manifests in distinct pathological outcomes depending on cellular context, including proteotoxic stress in neurons, metabolic dysregulation in hepatocytes, and contractile dysfunction in skeletal muscle, rather than a uniform proteostasis collapse across tissues [94,95]. All these findings support the main conclusion drawn from nematode research: safe and effective modulation of autophagy is likely to require both spatiotemporal precision and careful calibration, rather than uniform activation or inhibition across the whole organism [49,94]. Furthermore, hormetic interventions such as intermittent fasting, mild heat stress, or phytochemical-induced stress responses have shown to modestly enhance autophagy and improve proteostasis in mammals, echoing the beneficial effects of moderate autophagic activation observed in *C. elegans* [84]. These findings suggest that fine-tuning autophagic activity, rather than maximizing it, may be the most promising strategy for promoting human healthspan.

## 8. Conclusions

Together, these findings support a view of autophagy as a context-dependent regulator of aging rather than a uniformly beneficial process. Studies in *C. elegans* highlight how its effects on longevity depend on timing, tissue specificity, and physiological state, providing a framework to guide future efforts to modulate autophagy. Emerging genetic tools, such as auxin-inducible degron (AID)-based protein depletion, offer unprecedented opportunities to manipulate autophagy with high temporal and spatial precision, thereby improving our ability to distinguish beneficial versus detrimental autophagy activity during aging.

**Author Contributions:** Conceptualization, writing—original draft preparation, T.S.; writing—review and editing, J.B.; visualization, T.S.; supervision, J.B.; funding acquisition, J.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research has been supported by the National Research Development and Innovation Office (NKFIH) through the OTKA Grants FK131944. J.B. is also supported by the HUN-REN-ELTE Genetics Research Group (01062).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** This study is a review article and does not contain any original data.

**Acknowledgments:** The author is grateful to colleagues for insightful discussions and constructive feedback that contributed to the development of this review.

**Conflicts of Interest:** The authors declare no competing financial interest.

## Abbreviations

The following abbreviations are used in this manuscript:

AID	Auxin-inducible degradation
<i>atg</i>	Autophagy-related genes
<i>C. elegans</i>	<i>Caenorhabditis elegans</i>
DR	Dietary restriction
FOXA	Forkhead box A
HLH-30	Helix-loop-helix protein 30
IGF-1	Insulin growth factor 1
PHA-4	Pharynx Defective protein 4
PLPs	Pseudocoelomic lipoprotein pools
RNAi	RNA interference
SID	Systemic RNA interference defective
TFEB	Transcription factor EB
TOR	Target of Rapamycin
TLs	Tubular lysosomes

## References

1. Mizushima, N. A brief history of autophagy from cell biology to physiology and disease. *Nat. Cell Biol.* **2018**, *20*, 521–527. <https://doi.org/10.1038/s41556-018-0092-5>
2. Mizushima, N., Komatsu, M. Autophagy: renovation of cells and tissues. *Cell.* **2011** *11*;147(4):728-41. doi: 10.1016/j.cell.2011.10.026.
3. Levine, B.; Kroemer, G. Biological functions of autophagy genes: A disease perspective. *Cell* **2019**, *176*, 11–42. <https://doi.org/10.1016/j.cell.2018.09.048>

4. Hollenstein, D.M.; Kraft, C. Autophagosomes are formed at a distinct cellular structure. *Curr. Opin. Cell Biol.* **2020**, *65*, 50–57. <https://doi.org/10.1016/j.ceb.2020.02.012>
5. Mizushima, N. Autophagy: Process and function. *Genes Dev.* **2007**, *21*, 2861–2873. <https://doi.org/10.1101/gad.1599207>
6. Ohsumi, Y. Historical landmarks of autophagy research. *Cell Res.* **2014**, *24*, 9–23. <https://doi.org/10.1038/cr.2013.169>
7. Kuma, A.; et al. The role of autophagy during the early neonatal starvation period. *Nature* **2004**, *432*, 1032–1036. <https://doi.org/10.1038/nature03029>
8. Gatica, D.; Lahiri, V.; Klionsky, D.J. Cargo recognition and degradation by selective autophagy. *Nat. Cell Biol.* **2018**, *20*, 233–242. <https://doi.org/10.1038/s41556-018-0037-z>
9. Palmisano, N.J.; Meléndez, A. Autophagy in *C. elegans* development. *Dev. Biol.* **2019**, *447*, 103–125. <https://doi.org/10.1016/j.ydbio.2018.04.009>
10. Tian, Y.; et al. *C. elegans* screen identifies autophagy genes specific to multicellular organisms. *Cell* **2010**, *141*, 1042–1055. <https://doi.org/10.1016/j.cell.2010.04.034>
11. Tóth, M.L.; et al. Longevity pathways converge on autophagy genes to regulate life span in *Caenorhabditis elegans*. *Autophagy* **2008**, *4*, 330–338. <https://doi.org/10.4161/auto.5618>
12. Hars, E.S.; et al. Autophagy regulates ageing in *C. elegans*. *Autophagy* **2007**, *3*, 93–95. <https://doi.org/10.4161/auto.3636>
13. Wilhelm, T.; et al. Neuronal inhibition of autophagy extends lifespan. *Genes Dev.* **2017**, *31*, 1561–1572. <https://doi.org/10.1101/gad.301648.117>
14. Ezcurra, M.; et al. Intestinal autophagy in aging. *Curr. Biol.* **2018**, *28*, 2544–2556.e5. <https://doi.org/10.1016/j.cub.2018.06.035>
15. Alberti, A.; Michelet, X.; Djeddi, A.; Legouis, R. The autophagosomal protein LGG-2 acts synergistically with LGG-1 in dauer formation and longevity in *C. elegans*. *Autophagy* **2010**, *6*, 622–633. <https://doi.org/10.4161/auto.6.5.12252>
16. Palikaras, K.; Tavernarakis, N. In vivo mitophagy monitoring in *Caenorhabditis elegans* to determine mitochondrial homeostasis. *Bio Protoc.* **2017**, *7*, e2215. <https://doi.org/10.21769/BioProtoc.2215>
17. Kumsta, C.; et al. The autophagy receptor p62/SQST-1 promotes proteostasis and longevity in *C. elegans*. *Nat. Commun.* **2019**, *10*, 5648. <https://doi.org/10.1038/s41467-019-13540-4>
18. Palikaras, K.; Lionaki, E.; Tavernarakis, N. Coordination of mitophagy and mitochondrial biogenesis during ageing in *C. elegans*. *Nature* **2015**, *521*, 525–528. <https://doi.org/10.1038/nature14300>
19. Blackwell, T.K.; Sewell, A.K.; Wu, Z.; Han, M. TOR signaling in *Caenorhabditis elegans* development, metabolism, and aging. *Genetics* **2019**, *213*, 329–360. <https://doi.org/10.1534/genetics.119.302504>
20. Vellai, T.; et al. Influence of TOR kinase on lifespan in *C. elegans*. *Nature* **2003**, *426*, 620. <https://doi.org/10.1038/426620a>
21. Kenyon, C.; et al. A *C. elegans* mutant that lives twice as long as wild type. *Nature* **1993**, *366*, 461–464. <https://doi.org/10.1038/366461a0>
22. Lan, J.; et al. Translational regulation of non-autonomous mitochondrial stress response promotes longevity. *Cell Rep.* **2019**, *28*, 1050–1062.e6. <https://doi.org/10.1016/j.celrep.2019.06.078>
23. Panowski, S.H.; et al. PHA-4/Foxa mediates diet-restriction-induced longevity of *C. elegans*. *Nature* **2007**, *447*, 550–555. <https://doi.org/10.1038/nature05837>
24. O'Rourke, E.J.; Ruvkun, G. MXL-3 and HLH-30 link lipolysis and autophagy to nutrient availability. *Nat. Cell Biol.* **2013**, *15*, 668–676. <https://doi.org/10.1038/ncb2741>
25. Visvikis, O.; et al. Innate host defense requires TFEB-mediated transcription. *Immunity* **2014**, *40*, 896–909. <https://doi.org/10.1016/j.immuni.2014.05.002>
26. Hansen, M.; et al. Autophagy in lifespan extension by dietary restriction in *C. elegans*. *PLoS Genet.* **2008**, *4*, e24. <https://doi.org/10.1371/journal.pgen.0040024>
27. Guo, B.; et al. Genome-wide screen identifies regulators of autophagy in *C. elegans*. *EMBO Rep.* **2014**, *15*, 705–713. <https://doi.org/10.1002/embr.201338310>
28. Bördén, K.; Vellai, T.; Sigmond, T. Developing endogenous autophagy reporters in *C. elegans*. *Int. J. Mol. Sci.* **2025**, *26*, 10178. <https://doi.org/10.3390/ijms262010178>

29. Conte, D.; et al. RNA interference in *C. elegans*. *Curr. Protoc. Mol. Biol.* **2015**, *109*, 26.3.1–26.3.30. <https://doi.org/10.1002/0471142727.mb2603s109>
30. Maher, K.N.; et al. Large-scale gene knockdown in *C. elegans*. *J. Vis. Exp.* **2013**, *79*, e50693. <https://doi.org/10.3791/50693>
31. Calixto, A.; et al. Enhanced neuronal RNAi in *C. elegans*. *Nat. Methods* **2010**, *7*, 554–559. <https://doi.org/10.1038/nmeth.1463>
32. Whangbo, J.S.; et al. SID-1 domains important for dsRNA import. *G3* **2017**, *7*, 3887–3899. <https://doi.org/10.1534/g3.117.300308>
33. Vicencio, J.; et al. Engineering the auxin-inducible degron system. *Nat. Commun.* **2025**, *16*, 10848. <https://doi.org/10.1038/s41467-025-66347-x>
34. Son, H.G.; et al. Age-dependent biomarkers in *C. elegans*. *Aging Cell* **2019**, *18*, e12853. <https://doi.org/10.1111/ace1.12853>
35. Meléndez, A.; et al. Autophagy genes are essential for dauer development. *Science* **2003**, *301*, 1387–1391. <https://doi.org/10.1126/science.1087782>
36. Long, X.; et al. TOR deficiency causes developmental arrest. *Curr. Biol.* **2002**, *12*, 1448–1461. [https://doi.org/10.1016/S0960-9822\(02\)01091-6](https://doi.org/10.1016/S0960-9822(02)01091-6)
37. Gelino, S.; et al. Intestinal autophagy improves healthspan. *PLoS Genet.* **2016**, *12*, e1006135. <https://doi.org/10.1371/journal.pgen.1006135>
38. Hsieh, P.N.; et al. Conserved KLF-autophagy pathway. *Nat. Commun.* **2017**, *8*, 914. <https://doi.org/10.1038/s41467-017-00899-5>
39. Vérièpe-Salerno, J.; et al. MALT-1 shortens lifespan. *Autophagy Rep.* **2023**, *2*, 2277584. <https://doi.org/10.1080/27694127.2023.2277584>
40. Villalobos, T.V.; et al. Tubular lysosome induction. *Nat. Aging* **2023**, *3*, 1091–1106. <https://doi.org/10.1038/s43587-023-00470-6>
41. Sigmond, T.; Vellai, T. Lysosomal alteration links food limitation to longevity. *Nat. Aging* **2023**, *3*, 1048–1050. <https://doi.org/10.1038/s43587-023-00483-1>
42. Dolese, D.A.; et al. Tubular lysosomes link pexophagy and aging. *Autophagy* **2021**, 1–12. <https://doi.org/10.1080/15548627.2021.1990647>
43. Kumsta, C.; et al. Hormetic heat stress induces autophagy. *Nat. Commun.* **2017**, *8*, 14337. <https://doi.org/10.1038/ncomms14337>
44. Lapierre LR, De Magalhaes Filho CD, McQuary PR, Chu CC, Visvikis O, Chang JT, Gelino S, Ong B, Davis AE, Irazoqui JE, Dillin A, Hansen M. The TFEB orthologue HLH-30 regulates autophagy and modulates longevity in *Caenorhabditis elegans*. *Nat Commun.* **2013** 4:2267. doi: 10.1038/ncomms3267.
45. Edwards, C.; et al. Amino acid-mediated lifespan extension. *BMC Genet.* **2015**, *16*, 8. <https://doi.org/10.1186/s12863-015-0167-2>
46. Yang, J.; et al. miR-34 regulates lifespan via atg9. *Age* **2013**, *35*, 11–22. <https://doi.org/10.1007/s11357-011-9324-3>
47. Hansen, M.; Rubinsztein, D.C.; Walker, D.W. Autophagy and longevity. *Nat. Rev. Mol. Cell Biol.* **2018**, *19*, 579–593. <https://doi.org/10.1038/s41580-018-0033-y>
48. Madeo, F.; et al. Essential role for autophagy in lifespan extension. *J. Clin. Invest.* **2015**, *125*, 85–93. <https://doi.org/10.1172/JCI73946>
49. Castillo-Quan, J.I.; et al. Genetics of longevity. *Adv. Genet.* **2015**, *90*, 1–101. <https://doi.org/10.1016/bs.adgen.2015.06.002>
50. Gelino, S.; Hansen, M. Autophagy and aging. *J. Clin. Exp. Pathol.* **2012**, Suppl. 4. <https://doi.org/10.4172/2161-0681.S4-006>
51. Maiuri, M.C.; Kroemer, G. Autophagy in stress and disease. *Cell Death Differ.* **2015**, *22*, 365–366. <https://doi.org/10.1038/cdd.2014.236>
52. Levine, B.; Kroemer, G. Autophagy in aging and disease. *Cell Death Differ.* **2009**, *16*, 1–2. <https://doi.org/10.1038/cdd.2008.139>
53. Tóth, M.L.; et al. Autophagy genes and neurodegeneration. *J. Cell Sci.* **2007**, *120*, 1134–1141. <https://doi.org/10.1242/jcs.03401>

54. Kang, C.; You, Y.; Avery, L. Dual roles of autophagy in starvation. *Genes Dev.* **2007**, *21*, 2161–2171. <https://doi.org/10.1101/gad.1573107>
55. Zhou, B.; et al. Mitochondrial permeability and lifespan. *Cell* **2019**, *177*, 299–314.e16. <https://doi.org/10.1016/j.cell.2019.02.013>
56. Sakai, N.; et al. TORC2 signaling in learning. *PLoS ONE* **2017**, *12*, e0177900. <https://doi.org/10.1371/journal.pone.0177900>
57. Franco-Juárez, B.; et al. High glucose induces autophagy. *Aging* **2018**, *10*, 2657–2667. <https://doi.org/10.18632/aging.101577>
58. Takacs, Z.; et al. ATG-18 and EPG-6 in lifespan control. *Cells* **2019**, *8*, 236. <https://doi.org/10.3390/cells8030236>
59. Lu, Q.; et al. EPG-6 regulates autophagosome formation. *Dev. Cell* **2011**, *21*, 343–357. <https://doi.org/10.1016/j.devcel.2011.06.024>
60. Polson, H.E.J.; et al. WIPI2 regulates LC3 lipidation. *Autophagy* **2010**, *6*, 506–522. <https://doi.org/10.4161/auto.6.4.11863>
61. Hashimoto, Y.; et al. Suppression of autophagy extends lifespan. *Genes Cells* **2009**, *14*, 717–726. <https://doi.org/10.1111/j.1365-2443.2009.01306.x>
62. Byrne, J.; Wilhelm, T.; Richly, H. Neuronal autophagy and longevity. *Aging* **2017**, *9*, 1953–1954. <https://doi.org/10.18632/aging.101297>
63. Perez, M.F.; Lehner, B. Vitellogenins in *C. elegans*. *Front. Physiol.* **2019**, *10*, 1067. <https://doi.org/10.3389/fphys.2019.01067>
64. Mizunuma, M.; et al. mTORC2-SGK-1 in longevity. *Aging Cell* **2014**, *13*, 869–878. <https://doi.org/10.1111/accel.12248>
65. Meléndez, A.; et al. Monitoring autophagy in aging. *Methods Enzymol.* **2008**, *451*, 493–520. [https://doi.org/10.1016/S0076-6879\(08\)03229-1](https://doi.org/10.1016/S0076-6879(08)03229-1)
66. Palmisano, N.J.; et al. RAB-10 promotes autophagy. *Autophagy* **2017**, *13*, 1742–1753. <https://doi.org/10.1080/15548627.2017.1356976>
67. Jenzer, C.; et al. Autophagy in apoptosis. *Autophagy* **2019**, *15*, 228–241. <https://doi.org/10.1080/15548627.2018.1512452>
68. Jenzer, C.; et al. Human GABARAP rescues autophagy. *Autophagy* **2014**, *10*, 1868–1872. <https://doi.org/10.4161/auto.29745>
69. McGhee, J.D. The *C. elegans* intestine. *WormBook* **2007**, 1–36. <https://doi.org/10.1895/wormbook.1.133.1>
70. Yang, Y.; et al. ATG-16.2 mediates neuronal autophagy. *Nat. Aging* **2024**, *4*, 198–212. <https://doi.org/10.1038/s43587-023-00548-1>
71. Jung, R.; et al. Tissue-specific resistance to aggregation. *PLoS Biol.* **2023**, *21*, e3002284. <https://doi.org/10.1371/journal.pbio.3002284>
72. Gallardo-Campos, M.; et al. KFERQ-selective autophagy. *PLoS ONE* **2025**, *20*, e0330339. <https://doi.org/10.1371/journal.pone.0330339>
73. Mukherjee, A.; et al. Selective microautophagy in *Drosophila*. *Autophagy* **2016**, *12*, 1984–1999. <https://doi.org/10.1080/15548627.2016.1208887>
74. Nixon, R.A. Autophagy in neurodegeneration. *Nat. Med.* **2013**, *19*, 983–997. <https://doi.org/10.1038/nm.3232>
75. Murley, A.; et al. Macroautophagy and lysosomal damage. *Cell* **2025**, *188*, 2670–2686.e14. <https://doi.org/10.1016/j.cell.2025.03.009>
76. Zhong, R.; Richardson, C.E. Lysosomal expansion protects neurons. *PLoS Biol.* **2025**, *23*, e3002957. <https://doi.org/10.1371/journal.pbio.3002957>
77. Bansal, A.; et al. Lifespan vs. healthspan. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E277–E286. <https://doi.org/10.1073/pnas.1412192112>
78. Huang, C.; et al. Physiological predictors of lifespan. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 8084–8089. <https://doi.org/10.1073/pnas.0400848101>
79. Podshivalova, K.; et al. Aging and late-life decline. *Cell Rep.* **2017**, *19*, 441–450. <https://doi.org/10.1016/j.celrep.2017.03.062>

80. Kondapuram, S.K.; et al. Targeting autophagy in cancer. *JCMT* **2019**. <https://doi.org/10.20517/2394-4722.2018.105>
81. Thorburn, A.; et al. Autophagy and cancer therapy. *Mol. Pharmacol.* **2014**, *85*, 830–838. <https://doi.org/10.1124/mol.114.091850>
82. Shintani, T.; Klionsky, D.J. Autophagy in health and disease. *Science* **2004**, *306*, 990–995. <https://doi.org/10.1126/science.1099993>
83. Wong, S.Q.; et al. Autophagy in aging. *Hum. Genet.* **2020**, *139*, 277–290. <https://doi.org/10.1007/s00439-019-02031-7>
84. Stavoe, A.K.H.; Holzbaur, E.L.F. Autophagy in neurons. *Annu. Rev. Cell Dev. Biol.* **2019**, *35*, 477–500. <https://doi.org/10.1146/annurev-cellbio-100818-125242>
85. Djajadikerta, A.; et al. Autophagy induction as therapy. *J. Mol. Biol.* **2020**, *432*, 2799–2821. <https://doi.org/10.1016/j.jmb.2019.12.035>
86. Rahman, M.A.; Rhim, H. Autophagy in neurodegeneration. *BMB Rep.* **2017**, *50*, 345–354. <https://doi.org/10.5483/bmbrep.2017.50.7.069>
87. Kovács, T.; et al. AUTEN-99 protects against neurodegeneration. *Sci. Rep.* **2017**, *7*, 42014. <https://doi.org/10.1038/srep42014>
88. Papp, D.; et al. AUTEN-67 and neuroprotection. *Autophagy* **2016**, *12*, 273–286. <https://doi.org/10.1080/15548627.2015.1082023>
89. Billes, V.; et al. AUTEN-67 in Huntington's disease. *J. Huntingtons Dis.* **2016**, *5*, 133–147. <https://doi.org/10.3233/JHD-150180>
90. Madonna, R.; et al. Empagliflozin and autophagy. *Cardiovasc. Res.* **2023**, *119*, 1175–1189. <https://doi.org/10.1093/cvr/cvad009>
91. Wei, S.; et al. Arsenic-induced autophagy. *J. Hazard. Mater.* **2020**, *384*, 121390. <https://doi.org/10.1016/j.jhazmat.2019.121390>
92. Hartmann, J.; et al. Secretory autophagy and neurodegeneration. *Nat. Commun.* **2024**, *15*, 2635. <https://doi.org/10.1038/s41467-024-46953-x>
93. Kaushik, S.; Cuervo, A.M. Proteostasis and aging. *Nat. Med.* **2015**, *21*, 1406–1415. <https://doi.org/10.1038/nm.4001>
94. Rubinsztein, D.C.; Mariño, G.; Kroemer, G. Autophagy and aging. *Cell* **2011**, *146*, 682–695. <https://doi.org/10.1016/j.cell.2011.07.030>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.